

Appendix 2 (as supplied by authors): Cross-sectional association between suicide by someone personally known and suicidality outcomes in youth, assuming those who reported “I don’t know” regarding exposure to suicide were not exposed, OR (95% CI)

	Unadjusted	Partially Adjusted	Fully Adjusted
<u>Ages 12-13</u>			
Ideation			
Exposure in Past Year	2.88 (1.75-4.75)	2.92 (1.75-4.88)	3.58 (1.99-6.47)
Exposure > Year Ago	1.64 (1.12-2.42)	1.66 (1.13-2.44)	1.56 (0.99-2.45)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	2.20 (1.14-4.24)	2.14 (1.09-4.22)	2.70 (1.28-5.70)
Exposure > Year Ago	2.38 (1.45-3.90)	2.37 (1.44-3.90)	2.33 (1.29-4.20)
No Exposure	Reference	Reference	Reference
<u>Ages 14-15</u>			
Ideation			
Exposure in Past Year	2.45 (1.67-3.59)	2.26 (1.54-3.31)	2.46 (1.56-3.88)
Exposure > Year Ago	2.16 (1.59-2.94)	1.95 (1.43-2.67)	1.96 (1.36-2.84)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	3.64 (2.33-5.68)	3.26 (2.08 – 5.11)	3.36 (1.96-5.76)
Exposure > Year Ago	2.39 (1.63-3.51)	2.10 (1.41 – 3.12)	1.94 (1.21-3.11)
No Exposure	Reference	Reference	Reference
<u>Ages 16-17</u>			
Ideation			
Exposure in Past Year	1.75 (1.14-2.70)	1.74 (1.13-2.69)	1.73 (1.05-2.83)
Exposure > Year Ago	1.99 (1.43-2.77)	1.80 (1.28- 2.52)	1.69 (1.12-2.56)
No Exposure	Reference	Reference	Reference
Attempt			
Exposure in Past Year	2.46 (1.41-4.31)	2.44 (1.37-4.34)	2.69 (1.35-5.36)
Exposure > Year Ago	2.36 (1.50-3.71)	1.99 (1.23-3.22)	1.96 (1.07-3.59)
No Exposure	Reference	Reference	Reference

*Partially adjusted models account for gender and socioeconomic status. For ages 14-15 and 16-17, fully adjusted models account for gender, socioeconomic status, prior depression/anxiety, prior monthly drinking episodes, and prior drug use. For ages 12-13, fully adjusted models account for gender, socioeconomic status, and prior depression/anxiety.