

Appendix 3 (as supplied by authors): Prospective association between exposure to suicide and suicidality outcomes two years later, assuming those who reported “I don’t know” regarding exposure to suicide were not exposed, OR (95% CI)

	Unadjusted	Partially Adjusted	Fully Adjusted
<u>Exposure to suicide of a schoolmate</u>			
<u>Ages 12-13</u>			
Ideation	1.09 (0.48-2.47)	1.22 (0.54-2.75)	0.65 (0.22-1.91)
Attempt	1.87 (0.64-5.45)	2.15 (0.74-6.28)	0.95 (0.18-4.97)
<u>Ages 14-15</u>			
Ideation	1.32 (0.86-2.01)	1.31 (0.86-2.00)	1.44 (0.91-2.30)
Attempt	1.74 (0.98-3.06)	1.70 (0.96-3.01)	1.80 (0.94-3.46)
<u>Exposure to suicide of someone personally known by the respondent</u>			
<u>Ages 12-13</u>			
Ideation	1.20 (0.60-2.38)	1.26 (0.63-2.52)	1.24 (0.59-2.62)
Attempt	1.42 (0.58-3.50)	1.51 (0.60-3.76)	1.62 (0.57-4.61)
<u>Ages 14-15</u>			
Ideation	1.48 (0.93-2.36)	1.42 (0.90-2.24)	1.39 (0.82-2.35)
Attempt	2.03 (1.06-3.90)	1.87 (0.98-3.57)	1.77 (0.83-3.80)

*Partially adjusted models account for gender and socioeconomic status. For ages 14-15 and 16-17, fully adjusted models account for gender, socioeconomic status, prior depression/anxiety, prior monthly drinking episodes, and prior drug use. For ages 12-13, fully adjusted models account for gender, socioeconomic status, and prior depression/anxiety.