

Title: Nutrient Re-Supplementation Arrests Bio-Oil Accumulation in *Phaeodactylum tricornutum*

Contributors: Valenzuela, J.^{1,2}, R.P. Carlson^{2,3,6}, R. Gerlach^{2,3,6}, K.E. Cooksey⁴, B.M. Peyton^{2,3,6}, B. Bothner^{1,6}, and M.W. Fields^{2,4,5,6*}

Affiliations: ¹Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT

²Center for Biofilm Engineering, Montana State University, Bozeman, MT

³Department of Chemical and Biological Engineering, Montana State University, Bozeman, MT

⁴Department of Microbiology, Montana State University, Bozeman, MT

⁵Adjunct Research Fellow, National Center for Genome Resources, Santa Fe, NM

⁶Thermal Biology Institute, Montana State University, Bozeman, MT

Journal: *Applied Microbiology and Biotechnology*

Corresponding Author:

Dr. M.W. Fields
matthew.fields@biofilm.montana.edu

Figure S1. Nile Red specific fluorescence (Nile Red fluorescence intensity/cell) in *P. tricornutum*. Graphs start at 90 hours to show time-points of initial nitrate depletion (92.5 hours or 98 hours). A) Specific Nile Red fluorescence in control conditions and nutrient replete conditions: Control (×), N replete (●), P replete (▲), N+P replete (■). B) Specific Nile Red fluorescence in control conditions and nutrient re-supplemented conditions: Control (×), N re-supplemented (●), P supplemented (▲), N+P supplemented (■).

