

SDC Table 1. Baseline clinical characteristics of HIV-infected persons, stratified by inclusion/exclusion status*

	Excluded from Study (n=8874)	Included in Study (n=10,841)
Age, years	46 (40, 52)	46 (40, 52)
Race, %		
White	39%	41%
Black	46%	49%
Other	15%	15%
Female, %	2.0%	2.3%
Comorbid Conditions, %		
Hypertension	33%	39%
Diabetes	6.0%	7.5%
Dyslipidemia	6.6%	15%
Smoking	11%	19%
Hepatitis C Virus	6.2%	16%
Measurements		
CD4+ Count, cells/mm ³	257 (42, 477)	220 (51, 410)
HIV Viral Load (1000 copies/mL)	19 (1, 110)	63 (16, 248)
Systolic blood pressure, mmHg	129 (116, 140)	127 (114, 139)
Diastolic blood pressure, mmHg	78 (70, 85)	77 (70, 84)
Body mass index, kg/m ²	25 (22, 28)	24 (22, 27)
Total cholesterol, mg/dL	167 (138, 195)	167 (140, 197)
Triglycerides, mg/dL	128 (96, 219)	138 (95, 211)
Low density lipoprotein, mg/dL	107 (81, 134)	98 (76, 124)
High density lipoprotein, mg/dL	36 (29, 46)	37 (29, 47)
Glucose, mg/dL	95 (86, 108)	94 (86, 106)
Albumin, g/dL	3.8 (3.3, 4.1)	3.9 (3.5, 4.2)
eGFR, mL/min/1.73m ²	91 (74, 111)	97 (82, 114)
eGFR <60mL/min/1.73m ²	1.2%	5.4%
Proteinuria	38%	33%

*Continuous variables reported as median (IQR). Proteinuria defined by urinalysis protein 30 mg/dL or greater.