

SDC Table 5. Association of era* of antiretroviral use and cumulative tenofovir with risk of kidney disease outcomes

Outcome	Tenofovir x Era Interaction	Era	Tenofovir Effect (per year of use)	P Value	Overall Tenofovir Effect (From Table 2)
Proteinuria					
Era x Tenofovir	p = 0.014	Early	2.15 (1.26, 3.68)	0.0051	1.34 (1.25-1.45)
		Mid	1.48 (1.28, 1.71)	<.0001	
		Late	1.23 (1.16, 1.31)	<.0001	
Rapid Decline					
Era x Tenofovir	p = 0.0018	Early	1.69 (0.96, 3.00)	0.071	1.11 (1.03-1.18)
		Mid	1.34 (1.18, 1.53)	<.0001	
		Late	1.04 (0.98, 1.12)	0.20	
Chronic Kidney Disease					
Era x Tenofovir	p = 0.12	Early	3.34 (1.23-9.04)	0.018	1.33 (1.18-1.51)
		Mid	1.60 (1.25-2.05)	0.0002	
		Late	1.35 (1.21-1.50)	<.0001	

*Early = <2003, mid = 2003-04, late = 2005-07. All estimates based on multivariable adjusted time-dependent Cox models described in Table 2.