

Supplementary Data

SUPPLEMENTARY TABLE S1. INCIDENCE AND NUMBERS OF CASES OF THE FOLLICULAR VARIANT OF PAPILLARY THYROID CANCER (MORPHOLOGY CODE 8340) COMPARED TO THE PAPILLARY AND FOLLICULAR GROUPS WITHOUT 8340 OVER STUDY PERIOD, 1980–2009

Year of diagnosis	Follicular variant of papillary		Follicular without 8340		Papillary without 8340	
	IR	Count	IR	Count	IR	Count
1980	0.87	180	0.90	172	2.01	410
1981	0.88	184	0.76	152	2.28	478
1982	0.87	180	0.75	144	2.46	511
1983	0.72	157	0.91	183	2.62	557
1984	0.67	145	0.79	162	2.87	621
1985	0.73	161	0.85	173	2.98	649
1986	0.76	165	0.84	177	3.23	701
1987	0.69	154	0.90	190	2.99	665
1988	0.71	164	0.82	179	2.90	656
1989	0.78	177	0.88	193	3.25	748
1990	0.94	214	0.76	168	3.36	772
1991	0.93	215	0.73	162	3.27	780
1992	1.11	268	0.83	194	3.39	821
1993	1.10	264	0.82	195	3.22	792
1994	1.37	337	0.85	205	3.47	858
1995	1.50	374	0.74	181	3.57	905
1996	1.52	389	0.90	224	3.67	940
1997	1.59	407	0.77	193	3.94	1019
1998	1.78	465	0.80	206	3.95	1038
1999	1.85	486	0.82	213	4.14	1098
2000	1.95	520	0.79	209	4.44	1190
2001	2.13	577	0.83	221	4.80	1304
2002	2.40	654	0.84	228	5.43	1490
2003	2.36	651	0.93	254	5.73	1577
2004	2.60	723	0.98	269	5.92	1654
2005	2.72	767	1.07	301	6.47	1826
2006	2.23	644	1.10	315	7.30	2098
2007	2.83	826	1.02	295	7.83	2271
2008	2.88	847	1.08	316	8.43	2475
2009	3.20	951	1.14	336	9.33	2765

Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population.