



Breast milk and cognitive development – resolving the nature vs. nurture conflict: A systematic review

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-003259
Article Type:	Research
Date Submitted by the Author:	20-May-2013
Complete List of Authors:	Walfisch, Asnat; Hospital for Sick Children, Motherisk Program Sermer, Corey; Hospital for Sick Children, Motherisk Program Cressman, Alex; Hospital for Sick Children, Motherisk Program Koren, Gideon; Hospital for Sick Children, Motherisk Program
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Paediatrics, Reproductive medicine, Public health
Keywords:	Developmental neurology & neurodisability < PAEDIATRICS, PERINATOLOGY, PUBLIC HEALTH, REPRODUCTIVE MEDICINE

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3 Breast milk and cognitive development – resolving the nature
4 vs. nurture conflict: A systematic review
5
6
7
8

9
10 **Contributing Authors:**

11 Asnat Walfisch MD¹, Corey Sermer², Alex Cressman MSc¹, and Gideon Koren MD
12 FACMT FRCPC¹
13

14
15
16
17 **Author Affiliations:**

18
19 ¹ The Motherisk Program, Hospital for Sick Children; Department of Pediatrics and
20 Pharmacology, University of Toronto, Toronto, Ontario, Canada
21

22
23 ²University of Western Ontario, London, Ontario, Canada
24
25
26

27
28 **Corresponding author (and reprint requests):**

29 Gideon Koren, MD, FACMT, FRCPC

30
31 Director of Motherisk Program

32
33 Hospital for Sick Children, 555 University Avenue, M5G 1X8

34
35 Toronto, Ontario, Canada

36
37 Phone: 416-813-6780

38
39 Fax: 416-813-7562

40
41 E-mail: gidiup_2000@yahoo.com
42

43
44 **Word count:**

45
46 Abstract: 259 words

47
48 Manuscript (excluding references): 2,221

49
50 Running title: Breast Milk and Cognitive Development – Walfisch et al
51
52
53
54
55
56
57
58
59
60

ABSTRACT:

Objectives: The association between breastfeeding and child cognitive development is conflicted by studies reporting positive and null effects. Consequently, debate remains whether such a relationship is confounded by factors associated with breastfeeding, specifically maternal socioeconomic class, IQ and education.

Methods: To try to resolve this debate, we conducted a systematic review of the literature investigating the association between breastfeeding and cognitive outcomes of healthy infants born at term.

Results: Eighty-four studies met our inclusion criteria with 34 rated as high quality, 26 as moderate and 24 as low quality. Critical assessment of accepted studies revealed the following associations: 14 (null), 25 (positive), 15 (null after adjusting for confounders), 14 (positive - diminished after adjusting for confounders), and 16 with mixed results with positive associations in sub-group analysis. Directionality of effect did not correlate with study quality, however studies showing a decreased effect after multivariate analysis were of superior quality compared to other study groupings (i.e. 12/14 high quality). Further, studies that showed null or diminished effect after multivariate analysis corrected for significantly more confounders (7.9 ± 3.4) as compared to those that found no change following adjustment (4.8 ± 4.1) ($p = 0.001$).

Conclusion: The results of this systematic review support a conclusion that much of the reported effect of breastfeeding on child neurodevelopment may be due to confounding. Although it is unlikely that additional work will change the current synthesis, future studies in this field should attempt to rigorously control for all important confounders. Alternatively, study designs using sibling cohorts discordant for breastfeeding may yield more robust conclusions.

1
2
3 **Article summary:**
4

5 **Article focus:**
6

- 7
- 8 • Although most published data supports the association between breast-feeding
9 and IQ of the offspring, debate remains whether this is a causal relationship or
10 an association with favorable parental characteristics.
11
 - 12 • We conducted a systematic review of the literature investigating the
13 association between breastfeeding and cognitive outcomes of healthy term
14 infants.
15
16
17
18
19

20
21 **Key messages:**
22

- 23
- 24 • Over eighty studies addressing this issue were published with conflicting
25 results.
26
 - 27 • Studies, where the initial positive effect of breastfeeding on IQ disappeared or
28 diminished after multivariate analysis, controlled for almost twice as many
29 confounders than studies showing no such change.
30
31
 - 32 • Much of the reported effect of breastfeeding on child cognitive abilities is due
33 to maternal cognitive and socioeconomic effects.
34
35
36
37
38

39 **Strengths and limitations of this study:**
40

- 41
- 42 • The significant heterogeneity in study design and rigor precluded the conduct
43 of a formal meta-analysis.
44
45
46
47

48 **Funding**
49

50 This research received no specific grant from any funding agency in the public,
51 commercial or not-for-profit sectors.
52
53
54
55
56
57
58
59
60

INTRODUCTION:

Breastfeeding confers a range of nutritional and immunological advantages upon infants including reduction in childhood illness,¹⁻⁵ diabetes,⁶⁻⁷ and obesity.⁸

The potential of breast milk to enhance cognitive development has been the focus of numerous studies since Hoefler and Hardy's initial observation in 1929.⁹ It is generally agreed that children who breast-feed are more intelligent, however debate remains whether this is a causal relationship or merely an association with favorable parental socioeconomic class and IQ. The beneficial effects of breastfeeding on the child's neurodevelopment are hypothesized, by some, to be mediated by long-chain polyunsaturated fatty acids (LCPUFAs)¹⁰ which are present in human milk, but not in cow's milk or most infant formulas.¹¹ However, a recent systematic review of all randomized trials where mother's diet was supplemented with PUFA's during pregnancy has failed to confirm such an effect.¹²

The pendulum of opinion has swung back and forth with different investigators showing inconsistent results depending upon study design and rigor. The Achilles heel of most of these studies, and the probable explanation for the conflicting results, is the difficulty in controlling for confounders that may affect child development. Furthermore, the ability to clarify this relationship is hindered by ethical considerations, which preclude randomized controlled trials, given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

Well-established confounders in breastfeeding research include demographic and IQ differences between mothers who breastfeed and those who chose not to.¹³ Parents who score high on a range of cognitive abilities have children with above average IQ scores.¹³ In parallel, advantage in mother's IQ more than doubles the odds of

1
2
3 breastfeeding.¹³ Thus, some of the published data demonstrates the disappearance of
4
5 the breast-feeding effect on child's cognition after correction for maternal IQ.
6

7
8 In an attempt to partially overcome these sources of bias, a few randomized trials
9
10 have been published, with randomization to breastfeeding promotion¹⁴⁻¹⁵ or in preterm
11
12 infants.¹⁶ In the breastfeeding promotion intervention trial (PROBIT Trial) by Kramer
13
14 and colleagues, IQ scores and academic performance tests were more favorable in the
15
16 intervention group, however, statistically significant differences were only shown for
17
18 some of the sub-scores.¹⁵
19

20
21 Systematic reviews examining the impact of breastfeeding on cognitive abilities have
22
23 reached conflicting results.^{13, 17-19} The Meta-analysis by Jain et al¹⁸ suggests that less
24
25 than 25% of studies in this area have adjusted for socio-demographic confounders.
26

27
28 There is a paucity of literature critically assessing the current published evidence
29
30 within this field. In trying to address these challenges, the objective of the present
31
32 work was to conduct a systematic review of published studies investigating the
33
34 association between breast-feeding and neurodevelopmental outcome of healthy
35
36 infants born at term.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS:

The study was conducted based on a prospectively prepared protocol, using the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines.²⁰

Literature search:

Searches were conducted in the following databases (all from inception to December 2012): MEDLINE(R) with Daily Update, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, Health Technology Assessment, NHS Economic Evaluation Database, Embase and PsycINFO using the OvidSP interface, and on Science Citation Index Expanded (SCI-EXPANDED), Social Sciences Citation Index (SSCI), Conference Proceedings Citation Index- Science (CPCI-S), and Conference Proceedings Citation Index - Social Science & Humanities (CPCI-SSH) using the Web of Knowledge interface.

A search strategy was developed based on the MEDLINE, EMBASE and PsycINFO database subject headings and the 'Used for' synonyms listed in the scope notes for the terms "breast feeding", "human milk", "breast milk", "infant formula", "artificial milk", "cognition", "intelligence" and "intelligent quotient". Reference lists of identified studies, textbooks, previously published systematic reviews, and review articles were also searched. No language restrictions were applied and studies in languages other than English were translated for incorporation into the study.

Study selection:

Both prospective and retrospective studies (RCTs, non-randomised controlled clinical trials, cohort studies, longitudinal studies, and case-control studies) were included if:

1. One of the study aims was to address the question of breastfeeding and cognitive

1
2
3 development;

- 4
5 2. The authors used reliable validated methods to evaluate cognitive development
6
7 [e.g. Bayley scales of infant development, Wechsler Intelligence Scale for
8
9 Children Revised (WISC-R)];
10
11 3. Prospective or retrospective documentation of use and duration of breastfeeding
12
13 was used; and
14
15 4. The authors focused on healthy term infants and not those at increased biological
16
17 risk for developmental delays (e.g. prematurity, intrauterine growth restriction).
18
19

20 Studies were excluded if:

- 21
22 1. The study group included preterm or small for gestational age babies.
23
24 2. Evaluation of the cognitive development was done using only a non-reliable or
25
26 subjective tool (such as school grades, or maternal report).
27
28 3. Dietary patterns and breastfeeding were not evaluated since birth.
29
30

31
32 Titles and abstracts were reviewed for possible exclusion by two reviewers (A.W. and
33
34 C.S.). If both reviewers excluded a citation, it was eliminated from further review. If
35
36 at least one reviewer included the citation or if there was insufficient information to
37
38 make a determination from the title and abstract, the full article was obtained for
39
40 review. Full text articles were reviewed by three authors (A.W., C.S. and A.C.) for
41
42 suitability for inclusion. Disagreements regarding study eligibility were resolved by
43
44 consensus.
45
46

47 Study quality grading

48
49 Quality assessment of individual studies was performed by two authors (A.W and
50
51 C.S.) using the three category summary grading system (A, B, C) suggested by Ip et
52
53 al.¹⁹ Their system defines a generic grading system that is applicable to each type of
54
55
56
57
58
59
60

1
2
3 study design including randomized controlled trials, cohort, and case-control studies
4
5 as follows:

6
7 **A (good)**

8
9 A study that adheres mostly to the commonly held concepts of high quality including
10 the following: clear description of the population, setting, interventions and
11 comparison groups; clear description of the comparison groups; appropriate
12 measurement of outcomes; appropriate statistical and analytic methods and reporting;
13 no reporting errors; less than 20 percent dropout; clear reporting of dropouts; and
14 appropriate consideration and adjustment for potential confounders.
15
16
17
18
19
20
21

22
23 **B (fair/moderate)**

24
25 Category B studies do not meet all the criteria in category A because they have some
26 deficiencies, but none of which are likely to cause major biases. The study may have
27 suboptimal adjustment for potential confounders. The study may also be missing
28 information, making it difficult to assess limitations and potential problems.
29
30
31
32

33
34 **C (poor)**

35
36 Category C studies either did not consider potential confounders or did not adjust for
37 them appropriately. These studies may have serious shortcomings in design, analysis
38 or reporting; have large amounts of missing information, or discrepancies in
39 reporting.
40
41
42
43
44

45 **Data extraction:**

46
47 Extracted data were compiled in an evidence table. The table includes a description of
48 the studies that addressed the key question according to the inclusion/exclusion
49 criteria. The table provides information about study design, target population and
50 sample size, description of breast-feeding exposure and method of categorization,
51 nature of the comparison group, cognitive development assessment tool used and
52
53
54
55
56
57
58
59
60

1
2
3 participants' age, summary of the results prior to adjustment for confounders, a list of
4
5 all confounders adjusted for, differences in IQ between the groups after adjustment
6
7 for possible confounders (if available), and study quality grading according to the
8
9 scale described above.¹⁹
10

11
12 Statistical analysis:
13

14 Comparison of studies based on their results or quality was performed by Chi Square
15
16 or ANOVA as appropriate.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS:

The flow of the literature search is displayed in Figure 1. Of the 1,696 potentially relevant citations identified, 84 studies met the *a priori* inclusion criteria for this systematic review (Table 1).^{9,13,15, 21-101} Overall agreement between reviewers on the inclusion of studies was 100%.

Out of these 84 publications, 34 were rated as high quality (grade A), 26 as moderate (grade B) and 24 as low quality (grade C). Overall, there were 14 studies showing no association between IQ and breastfeeding, 25 positive studies, 15 initially positive studies that became negative after accounting for confounders, and 14 studies where the initial positive effect was diminished but remained statistically significant after accounting for confounders. The remaining 16 studies showed mixed results (e.g. association only observed in the female sub-group or only in carriers of a specific allele).

In general, the directionality of the results did not correlate with the studies' quality. However, the studies showing decrease in the effect after multivariate analysis were of superior quality compared to the rest (i.e. 12 of 14 had a quality score of A).

Different studies corrected in their analyses for different potential confounders, ranging from zero to 16 total confounders (Table). Confounders commonly considered in these studies were socioeconomic status, maternal education, birth weight, gestational age, birth order and gender. Some considered quality and quantity of stimulation of the child to be crucial confounders but did not consider maternal or paternal intelligence and other important factors. Studies that showed null or diminished effect in their multivariate analysis controlled for significantly more confounders (7.9 ± 3.4) as compared to those that found no change following adjustment for confounders (4.8 ± 4.1) ($p = 0.001$). Furthermore, many of the studies

1
2
3 did not have a clear definition of breastfeeding or cumulative breast milk exposure. A
4
5 large variety of cognitive assessment tools were used and study outcomes were
6
7 measured anywhere from 8 days of age into adulthood. The significant heterogeneity
8
9 in study design and rigor precluded the conduct of a formal meta-analysis.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

DISCUSSION:

The continuing debate of whether breastfeeding imparts direct advantage on child cognition, or whether this is merely an association with favorable familial socioeconomic status and cognition, is not purely theoretical. From a public health perspective, if breastfeeding has biological effects on a child's IQ - this will be one of very few cost-effective means to significantly improve a child's neurodevelopment. If, on the other hand, there is no such effect, in the case where breastfeeding is either impossible or not sought by the mother, this will allow these women to rest assured that their choice will not have long-term developmental consequences.

In the case of other comparable therapeutic dilemmas, these conflicts is typically resolved through randomized controlled trials (RCT), which are not ethically feasible in this case given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

The closest comparison to a formal RCT in reducing selection bias would be sibling-pair analysis, when cognition of breastfed infants is compared to that of their siblings who were formula fed. This design ensures similar socioeconomic and maternal characteristics. Importantly, the few studies that have followed this design were negative or positive.^{13, 40}

The second closest design to RCT was employed in the PROBIT study by Kramer and colleagues,¹⁴⁻¹⁵ who cluster randomized women in Eastern Europe to receive or not receive formal education about the advantages of breastfeeding. This study did show favorable effects, but it has been argued that the mothers randomized for the breastfeeding promotion arm might have been influenced not only in providing higher rates of breastfeeding, but also by improving other positive health behaviors.

1
2
3 Our analysis reveals that there are over eighty studies addressing this issue and that
4 their results divide almost evenly between positive and negative associations. The
5 quality of “positive” or “negative” studies did not differ, except for higher quality on
6 average in studies that showed an apparent decrease in effect after multivariate
7 analysis.
8
9

10
11
12
13
14 We have shown that studies, where the initial positive effect of breastfeeding on IQ
15 disappeared or substantially diminished after multivariate analysis, controlled for
16 almost twice as many confounders than studies showing no such change. When
17 compared to a meta-analysis conducted 14 years ago,¹⁷ it appears that many more new
18 studies did attempt to control for confounding measures of socioeconomic status and
19 parental education, amongst others.
20
21

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Given that more tight control of confounders resulted in greater likelihood of disappearance of breastfeeding effect, it can be argued that the remaining positive effect reflects residual uncontrolled bias, as shown by Der and colleagues in their large study.¹³ In that study, before adjustment, breastfeeding was associated with an increase of around 4 points in mental ability. *Post hoc* analysis revealed that adjustment for maternal intelligence accounted for most of this effect – where full adjustment for a range of relevant confounders yielded a small (0.52) and non-significant effect size (95% confidence interval -0.19 to 1.23). In our systematic review, a similar effect was recorded by a total of 15 studies, and in addition 14 studies showed substantially diminished effect after adjustment.

Another factor that needs to be seriously considered in our review is the existence of bias against the null hypothesis. The likelihood of studies not detecting a significant effect in pregnancy to be submitted and published in the peer review literature is

1
2
3 substantially lower than that of positive studies¹⁰²⁻¹⁰³. This can create a distorted
4
5 balance that may seriously affect the conclusions on effects of interventions.
6

7
8 In conclusion, this systematic review suggests that much of the reported effect of
9
10 breastfeeding on child cognitive abilities is due to maternal cognitive and
11
12 socioeconomic effects. When considered together with the fact that a recent
13
14 systematic review failed to corroborate a biological effect of milk PUFA on brain
15
16 development, it is likely that breastfeeding does not, by itself, directly affect child IQ.
17
18 Although it is unlikely that additional studies will change substantially the current
19
20 synthesis, future studies in this field should attempt to rigorously control for all
21
22 important confounders even if they are difficult to obtain (e.g. parental IQ).
23
24 Alternatively, study designs using sibling cohorts discordant for breastfeeding may
25
26 yield more robust conclusions to further clarify this dilemma.
27
28
29
30
31

32 Author's contribution:

33
34 Asnat Walfisch has participated in all phases of this study including literature search,
35
36 data collection and review, quality grading, interpretation of the results, and has
37
38 written a part of the manuscript.

39
40 Corey Sermer has participated in all phases of this study including literature search,
41
42 data collection and review, quality grading, and revision of the manuscript.

43
44 Alex Cressman has participated in the literature search, data collection and review,
45
46 translation of foreign language manuscripts, and has taken a significant part in the
47
48 manuscript preparation and revision.

49
50 Gideon Koren participated in all phases of this study. He initiated the study and
51
52 supervised actively throughout its conduct. Specifically he was involved in the data
53
54 interpretation, statistical analysis, and has written and revised a substantial part of the
55
56 manuscript.
57
58
59
60

References

1. Ryan AS, Wenjun Z, Acosta A. Breastfeeding continues to increase into the new millennium. *Pediatrics*. 2002 Dec;110:1103-9.
2. Dell S, To T. Breastfeeding and asthma in young children: Findings from a population based study. *Archives of Pediatric and Adolescent Medicine* 2001;155:1261-1265.
3. Bhandari N, Bahl R, Mazumdar S, Martines J, Black RE, Bhan MK. Effect of community-based promotion of exclusive breastfeeding on diarrhoeal illness and growth: A cluster randomised trial. *Lancet* 2003;361:1418-1423.
4. Galton Bachrach VR, Schwarz E, Bachrach LR. Breastfeeding and the risk of hospitalisation for respiratory disease in infancy. A meta-analysis. *Archives of Pediatric and Adolescent Medicine* 2003;157:237-243.
5. Howie PW, Forsyth JS, Ogston SA, Clark A, Florey CD. Protective effect of breastfeeding against infection. *BMJ* 1990;300:11-16.
6. Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskeine E, Samuelsson U. Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood. *Diabetes Metabolism Research Reviews* 2004;20:150-157.
7. Young TK, Martens PJ, Taback SP, Sellers EAC, Dean HJ, Cheang M, Flett B. Type 2 diabetes mellitus in children prenatal and early infancy risk factors among native Canadians. *Archives of Pediatrics and Adolescent Medicine* 2002;156:651-655.
8. Owen CG, Martin RM, Whincup PH, Davey Smith G, Cook DG. Effect of infant feeding on the risk of obesity across the life course: A quantitative review of the published evidence. *Pediatrics* 2005;115:1367-1377.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
9. Hoefler C, Hardy MC. Later development of breast fed and artificially fed infants. *J Am Med Assoc.* 1929;92:615–20.
 10. Koletzko B, Lien E, Agostoni C, Böhles H, Campoy C, Cetin I, Decsi T, Dudenhausen JW, Dupont C, Forsyth S, Hoesli I, Holzgreve W, Lapillonne A, Putet G, Secher NJ, Symonds M, Szajewska H, Willatts P, Uauy R, World Association of Perinatal Medicine Dietary Guidelines Working Group. The roles of long chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *J Perinat Med* 2008;36:5–14.
 11. Sauerwald TU, Demmelmair H, Koletzko B. Polyunsaturated fatty acid supply with human milk. *Lipids.* 2001;36:991-6.
 12. Lo A, Sienna J, Mamak E, Djokanovic N, Westall C, Koren G. The effects of maternal supplementation of polyunsaturated Fatty acids on visual, neurobehavioural, and developmental outcomes of the child: a systematic review of the randomized trials. *Obstet Gynecol Int* 2012;2012:591531.
 13. Der G, Batty GD, Deary IJ. Effect of breast feeding on intelligence in children: prospective study, sibling pairs analysis, and meta-analysis. *BMJ* 2006;333(7575):945.
 14. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzikovich I, Shapiro S, et al for the PROBIT Study Group. Promotion of Breastfeeding Intervention Trial (PROBIT): A randomised controlled trial in the Republic of Belarus. *JAMA* 2001;285:413-420.
 15. Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L, Igumnov S, Fombonne E, Bogdanovich N, Ducruet T, Collet JP, Chalmers B, Hodnett E, Davidovsky S, Skugarevsky O, Trofimovich O, Kozlova L,

- 1
2
3 Shapiro S; Promotion of Breastfeeding Intervention Trial (PROBIT) Study
4
5 Group. Breastfeeding and child cognitive development: new evidence from a
6
7 large randomized trial. *Arch Gen Psychiatry* 2008;65(5):578-584.
8
9
10 16. Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and
11
12 subsequent intelligence quotient in children born preterm. *Lancet*
13
14 1992;339:261-4.
15
16 17. Anderson JW, Johnstone BM, Remley DT. Breastfeeding and cognitive
17
18 development: a meta-analysis. *Am J Clin Nutr* 1999;70:525-35.
19
20 18. Jain A, Concato J, Leventhal JM. How good is the evidence linking
21
22 breastfeeding and intelligence? *Pediatrics* 2002;109:1044-53.
23
24 19. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, Trikalinos T, Lau J.
25
26 Breastfeeding and maternal and infant health outcomes in developed
27
28 countries. *Evid Rep Technol Assess* 2007;153:1-186.
29
30 20. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting
31
32 systematic reviews and meta-analyses of studies that evaluate healthcare
33
34 interventions: explanation and elaboration. *BMJ*. 2009; 339:b2700.
35
36 21. Amanda M, Singh RP. A study of intelligence of children in relation to infant-
37
38 feeding practices and nutrition. *Indian Journal of Psychometry & Education*
39
40 1992;23:23-28.
41
42 22. Andres A, Cleves MA, Bellando JB, Pivik RT, Casey PH, Badger TM.
43
44 Developmental status of 1-year-old infants fed breast milk, cow's milk
45
46 formula, or soy formula. *Pediatrics*. 2012;129:1134-40.
47
48 23. Angelsen NK, Vik T, Jacobsen G, Bakketeig LS. Breast feeding and cognitive
49
50 development at age 1 and 5 years. *Arch Dis Child* 2001;85:183–188.
51
52 24. Auestad N, Halter R, Hall RT, Blatter M, Bogle ML, Burks W, Erickson JR,
53
54
55
56
57
58
59
60

Breast Milk and Cognitive Development – Walfisch et al

- 1
2
3 Fitzgerald KM, Dobson V, Innis SM, Singer LT, Montalto MB, Jacobs JR,
4
5 Qiu W, Bornstein MH. Growth and development in term infants fed long-
6
7 chain polyunsaturated fatty acids: a double-masked, randomized, parallel,
8
9 prospective, multivariate study. *Pediatrics*. 2001;108(2):372-81.
- 10
11
12 25. Auestad N, Scott DT, Janowsky JS, Jacobsen C, Carroll RE, Montalto MB,
13
14 Halter R, Qiu W, Jacobs JR, Connor WE, Connor SL, Taylor JA, Neuringer
15
16 M, Fitzgerald KM, Hall RT. Visual, cognitive, and language assessments at 39
17
18 months: a follow-up study of children fed formulas containing long-chain
19
20 polyunsaturated fatty acids to 1 year of age. *Pediatrics*. 2003;112(3 Pt 1):e177-
21
22 83.
- 23
24
25 26. Bartels M, van Beijsterveldt CEM, Boomsma I. Breastfeeding, Maternal
26
27 Education and Cognitive Function: A Prospective Study in Twins. *Behav*
28
29 *Genet* 2009;39:616–622.
- 30
31
32 27. Bauer G, Ewald ES, Hoffman J, Dubanoski R. Breast feeding and cognitive
33
34 development in three-year-old children. *Arch Dis Child* 1991;64:763–4.
- 35
36
37 28. Birch EE, Garfield S, Castañeda Y, Hughbanks-Wheaton D, Uauy R, Hoffman
38
39 D. Visual acuity and cognitive outcomes at 4 years of age in a double-blind,
40
41 randomized trial of long-chain polyunsaturated fatty acid-supplemented infant
42
43 formula. *Early Human Development* 2007;83:279–284.
- 44
45
46 29. Bon N. Relationship between breastfeeding and subsequent IQ of child.
47
48 *Bulletin de Psychologie* 1980;33:935-947.
- 49
50
51 30. Bouwstra H, Dijck-Brouwer DAJ, Boehm G, Boersma ER, Muskiet FAJ,
52
53 Hadders-Algra M. Long-chain polyunsaturated fatty acids and neurological
54
55 developmental outcome at 18 months in healthy term infants. *Acta Pædiatrica*,
56
57 2005; 94: 26–32.
- 58
59
60

- 1
2
3 31. Burruchaga MS, Sanz JIR, Zubizarreta JIP, Fernandez JB, Crespo PS.
4
5 Intellectual development in the second year of life in healthy breast-fed
6
7 children vs. formula-fed children. *An Esp Pediatr* 2000; 52: 530-536.
8
9
10 32. Caspi A, Williams B, Kim-Cohen J, Craig IW, Milne BJ, Poulton R,
11
12 Schalkwyk LC, Taylor A, Werts H, Moffitt TE. Moderation of breastfeeding
13
14 effects on the IQ by genetic variation in fatty acid metabolism.
15
16 *Proc.Natl.Acad.Sci.U.S.A.* 2007;104:18860-18865.
17
18
19 33. Clark KM, Castillo M, Calatroni A, Walter T, Cayazzo M, Pino P, Lozoff B.
20
21 Breastfeeding and Mental and Motor Development at 5 1/2 Years. *Ambul*
22
23 *Pediatr.* 2006 ; 6(2): 65–71.
24
25
26 34. Daniels MC, Adair LS. Breast-feeding influences cognitive development in
27
28 Filipino children. *J Nutr* 2005;135:2589–2595.
29
30
31 35. de Andraca,I.; Pino,P.; de la Parra,A.; Rivera,F.; Castillo,M. Risk factors for
32
33 psychomotor development among infants born under optimal biological
34
35 conditions. *Rev Saude Publica* 1998;32:138-147.
36
37
38 36. De Andraca,I.; Salas,MI; Lopez,C.; Cayazzo,MS; Icaza,G. Effect of breast
39
40 feeding an psychosocial variables upon psychomotor development of 12
41
42 months old infants. *Arch Latinoam Nutr* 1999;49:223-231.
43
44
45 37. Di Girolamo D, Braibanti P, Mamone P. Feeding methods and cognitive
46
47 development in the first 3 years of life. *Rassegna di Psicologia* 1993;10:33-49.
48
49
50 38. Eickmann SH, de Lira PI, Lima Mde C, Coutinho SB, Teixeira Mde L,
51
52 Ashworth A. Breast feeding and mental and motor development at 12 months
53
54 in a low-income population in northeast Brazil. *Paediatr Perinat Epidemiol*
55
56 2007;21:129-137.
57
58
59 39. Elwood PC, Pickering J, Gallacher JEJ, Hughes J, Davies D. Long term effect
60

Breast Milk and Cognitive Development – Walfisch et al

- of breast feeding: Cognitive function in the Caerphilly cohort. *J.Epidemiol.Community Health* 2005;59:130-133.
40. Evenhouse E, Reilly S. Improved estimates of the benefits of breastfeeding using sibling comparisons to reduce selection bias. *Health Serv Res* 2005;40:1781-1802.
41. Fergusson DM, Beautrais AL, Silva PA. Breast-feeding and cognitive development in the first seven years of life. *Social Science and Medicine* 1982;16:1705-1708.
42. Florey Du CV, Leech AM, Blackhall A. Infant feeding and mental and motor development at 18 months of age in first born singletons. *Int J Epidemiol* 1995;24:S21-S26.
43. Foroushani AR, Mohammad K, Mahmoodi M, Siassi F. Effect of breastfeeding on cognitive performance in a British birth cohort. *Eastern Mediterranean Health Journal* 2010;16:202-208.
44. Gale CR, Martyn CN. Breastfeeding, dummy use, and adult intelligence. *Lancet* 1996;347:1072-1075.
45. Gale CR, Marriott LD, Martyn CN, Limond J, Inskip HM, Godfrey KM, Law CM, Cooper C, West C, Robinson SM. Breastfeeding, the use of docosahexaenoic acid-fortified formulas in infancy and neuropsychological function in childhood. *Arch Dis Child* 2010;95:174-179.
46. Ghys A, Bakker E, Hornstra G, Van den Hout M. Red blood cell and plasma phospholipid arachidonic and docosahexaenoic acid levels at birth and cognitive development at 4 years of age. *Early Hum Dev* 2002;69:83-90.
47. Gibson-Davis CM, Brooks-Gunn J. Breastfeeding and verbal ability of 3-year-olds in a multicity sample. *Pediatrics* 2006;118: e1444-e1445.

- 1
2
3 48. Gomez-Sanchiz M, Canete R, Rodero I, Baeza JE, Gonzalez JA. Influence of
4 breast-feeding and parental intelligence on cognitive development in the 24-
5 month-old child. *Clin Pediatr* 2004;43:753-761.
6
7
8
9
10 49. GREENE LC, LUCAS A, LIVINGSTONE MBE, HARLAND PSEG,
11 BAKER BA. Relationship between Early Diet and Subsequent Cognitive
12 Performance during Adolescence. *Biochem Soc Trans* 1995;23:S376-s376.
13
14
15
16 50. Gurka KK, Kellams AL, Locasale-Crouch J, Gurka MJ. Breastfeeding and
17 cognition: The role of confounders. *Breastfeeding Medicine*. Conference: 15th
18 Annual International Meeting of the Academy of Breastfeeding Medicine -
19 Breastfeeding: A Bridge to the Gold Standard San Francisco, CA United
20 States. Conference Start: 20101027 Conference End: 20101030.Conf
21 (TRUNCATED) 2010;5:326.
22
23
24
25
26
27
28
29
30 51. Hart S, Boylan LM, Carroll S, Musick YA, Lampe RM. Brief Report: Breast-
31 fed One-Week-Olds Demonstrate Superior Neurobehavioral Organization. *J*
32 *Pediatr Psychol* 2003;28:529-533.
33
34
35
36 52. Holme A, MacArthur C, Lancashire R. The effects of breastfeeding on
37 cognitive and neurological development of children at 9 years. *Child Care*
38 *Health Dev* 2010;36:583-590.
39
40
41
42
43 53. Horwood LJ, Fergusson DM. Breastfeeding and later cognitive and academic
44 outcomes. *Pediatrics* 1998;101:E9.
45
46
47
48 54. Innis SM, Nelson CM, Lwanga D, Rioux FM, Waslen P. Feeding formula
49 without arachidonic acid and docosahexaenoic acid has no effect on
50 preferential looking acuity or recognition memory in healthy full- term infants
51 at 9 mo of age. *Am J Clin Nutr* 1996;64:40-46.
52
53
54
55
56 55. Jacobson SW, Chiodo LM, Jacobson JL. Breastfeeding effects on intelligence
57
58
59
60

Breast Milk and Cognitive Development – Walfisch et al

- 1
2
3 quotient in 4- and 11-year-old children. *Pediatrics* 1999;103:E71.
4
5
6 56. James PF. Breastfeeding and intelligence. *Lancet* 1992;339:614.
7
8 57. Jedrychowski W, Perera F, Jankowski J, Butscher M, Mroz E, Flak E, Kaim I,
9
10 Lisowska-Miszczuk I, Skarupa A, Sowa A. Effect of exclusive breastfeeding
11
12 on the development of children's cognitive function in the Krakow
13
14 prospective birth cohort study. *Eur J Pediatr* 2012;171:151–158.
15
16 58. Jiang M, Foster EM, Gibson-Davis CM. Breastfeeding and the Child
17
18 Cognitive Outcomes: A Propensity Score Matching Approach *Matern Child*
19
20 *Health J* 2011;15:1296–1307.
21
22 59. Johnson DL, Swank PR, Howie VM, Baldwin CD, Owen M. Breast feeding
23
24 and children's intelligence. *Psychol Rep* 1996;79:1179-1185.
25
26 60. Keim SA, Daniels J, Siega-Riz AM, Dole N. Long-Chain Polyunsaturated
27
28 Fatty Acids from Breastmilk and Formula and Infant Cognitive Development.
29
30 *Am J Epidemiol* 2009;169:S11.
31
32 61. Lawlor DA, Najman JM, Batty GD, O'Callaghan MJ, Williams GM, Bor W.
33
34 Early life predictors of childhood intelligence: Findings from the Mater-
35
36 University study of pregnancy and its outcomes. *Paediatr Perinat Epidemiol*
37
38 2006;20:148-162.
39
40 62. Lucas A, Stafford M, Morley R, Abbott R, Stephenson T, MacFadyen U,
41
42 EliasJones A, Clements H. Efficacy and safety of long-chain polyunsaturated
43
44 fatty acid supplementation of infant-formula milk: a randomised trial. *Lancet*
45
46 1999; 354:1948-1954.
47
48 63. Maimaitiming T, Wang YP. Feeding and intellectual development in infants
49
50 and children from Uygur, Kazak, and Han nationality: A comparison of 454
51
52 cases. *Journal of Clinical Rehabilitative Tissue Engineering Research*
53
54
55
56
57
58
59
60

- 1
2
3 2007;11:10710-10712.
4
5 64. Makrides M, Neumann MA, Simmer K, Gibson RA . A critical appraisal of
6
7 the role of dietary long-chain polyunsaturated fatty acids on neural indices of
8
9 term infants: A randomized, controlled trial. *Pediatrics* 2000;105:32-38.
10
11 65. Malloy MH, Berendes H. Does breast-feeding influence intelligence quotients
12
13 at 9 and 10 years of age?. *Early Hum Dev* 1998;50:209-217.
14
15 66. Martin NW, Benyamin B, Hansell NK, Montgomery GW, Martin NG, Wright
16
17 MJ, Bates TC. Cognitive Function in Adolescence: Testing for Interactions
18
19 Between Breast-Feeding and FADS2 Polymorphisms. *J Am Acad Child*
20
21 *Adolesc Psychiatry* 2011;50(1):55–62.
22
23 67. McCrory C, Layte R. The effect of breastfeeding on children's educational test
24
25 scores at nine years of age: Results of an Irish cohort study. *Social Science*
26
27 *and Medicine* 2011;72:1515-1521.
28
29 68. Morales E, Bustamante M, Gonzalez JR, Guxens M, Torrent M, et al. (2011)
30
31 Genetic Variants of the FADS Gene Cluster and ELOVL Gene Family,
32
33 Colostrums LC-PUFA Levels, Breastfeeding, and Child Cognition. *PLoS*
34
35 *ONE* 6(2): e17181.
36
37 69. Morris SS, Grantham-McGregor SM, Lira PIC, Assuncao AMS, Ashworth A.
38
39 Effect of breastfeeding and morbidity on the development of low birthweight
40
41 term babies in Brazil. *Acta Paediatr* 1999;88:1101-1106.
42
43 70. Morrow-Tlucak M, Haude RH, Ernhart CB. Breastfeeding and cognitive
44
45 development in the first 2 years of life. *Social Science and Medicine*
46
47 1988;26:635-639.
48
49 71. Mortensen EL, Michaelsen KF, Sanders SA, Reinisch JM. The association
50
51 between duration of breastfeeding and adult intelligence. *J Am Med Assoc*
52
53
54
55
56
57
58
59
60


- 1
2
3 2002;287:2365-2371.
4
5 72. Mukerji M, Sharma K. Mental development as a function of maternal
6 economic status, literacy, occupation and feeding pattern. *Psycho-Lingua*
7 1993;23:77-81.
8
9 73. Nassar MF, Younis NT, El-Arab SE, Fawzi FA. Neuro-developmental
10 outcome and brain-derived neurotrophic factor level in relation to feeding
11 practice in early infancy. *Maternal and Child Nutrition* 2011;7:188-197.
12
13 74. NELSON CM, INNIS SM, WALSEN P, WHITFIELD M. Prospective
14 Measures of Visual and Cognitive-Development in Term Gestation Breast-Fed
15 and Formula-Fed Infants to 18 Months of Age. *Pediatr Res* 1995;37:A315.
16
17 75. Niemela A, Jarvenpaa AL. Is breastfeeding beneficial and maternal smoking
18 harmful to the cognitive development of children?. *Acta Paediatrica,*
19 *International Journal of Paediatrics* 1996;85:1202-1206.
20
21 76. Oddy WH, Kendall GE, Blair E, De Klerk NH, Stanley FJ, Landau LI, Silburn
22 S, Zubrick S. Breast feeding and cognitive development in childhood: A
23 prospective birth cohort study. *Paediatr Perinat Epidemiol* 2003;17:81-90.
24
25 77. Oddy WH, Li J, Whitehouse AJO, Zubrick SR, Malacova E. Breastfeeding
26 duration and academic achievement at 10 years. *Pediatrics* 2011;127:e137-
27 e145.
28
29 78. Paine BJ, Makrides M, Gibson RA. Duration of breast-feeding and Bayley's
30 Mental Developmental Index at 1 year of age. *J Paediatr Child Health*
31 1999;35:82-85.
32
33 79. Pollock JI. Long-term associations with infant feeding in a clinically
34 advantaged population of babies. *Dev Med Child Neurol* 1994;36:429-440.
35
36 80. Quinn PJ, O'Callaghan M, Williams GM, Najman JM, Andersen MJ, Bor W.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 The effect of breastfeeding on child development at 5 years: A cohort study. *J*
4 *Paediatr Child Health* 2001;37:465-469.
5
6
7 81. Rao MR, Hediger ML, Levine RJ, Naficy AB, Vik T. Effect of breastfeeding
8 on cognitive development of infants born small for gestational age. *Acta*
9 *Paediatrica, International Journal of Paediatrics* 2002;91:267-274.
10
11 82. Ribas-Fito N, Julvez J, Torrent M, Grimalt JO, Sunyer J. Beneficial effects of
12 breastfeeding on cognition regardless of DDT concentrations at birth. *Am J*
13 *Epidemiol* 2007;166:1198-1202.
14
15 83. Richards M, Hardy R, Wadsworth MEJ. Long-term effects of breast-feeding
16 in a national birth cohort: Educational attainment and midlife cognitive
17 function. *Public Health Nutr* 2002;5:631-635.
18
19 84. Richards M, Wadsworth M, Rahimi-Foroushani A, Hardy R, Kuh D, Paul A.
20 Infant nutrition and cognitive development in the first offspring of a national
21 UK birth cohort. *Dev Med Child Neurol* 1998;40:163-167.
22
23 85. Rodgers B. Feeding in infancy and later ability and attainment: A longitudinal
24 study. *Developmental Medicine & Child Neurology* 1978;20:421-426.
25
26 86. Rogan WJ, Gladen BC. Breast-feeding and cognitive development. *Early Hum*
27 *Dev* 1993;31:181-193.
28
29 87. Santiago Burruchaga M, Ruiz Sanz JI, Pijoan Zubizarreta JI, Benito Fernandez
30 J, Sanjurjo Crespo P. Intellectual development in the second year of life in
31 healthy breast-fed children vs. Formula-fed children. *An Esp Pediatr*
32 2000;52:530-536.
33
34 88. Silva AAM, Mehta Z, O'Callaghan FJK. Duration of breast feeding and
35 cognitive function: Population based cohort study. *Eur J Epidemiol*
36 2006;21:435-441.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 89. Silva PA, Buckfield P, Spears GF. Some maternal and child developmental
4 characteristics associated with breast feeding: a report from the Dunedin
5 Multidisciplinary Child Development Study. *Aust Paediatr J* 1978;14:265-268.
6
7
8
9
10 90. Sloan S, Stewart M, Dunne L. The effect of breastfeeding and stimulation in
11 the home on cognitive development in one-year-old infants. *Child Care in*
12 *Practice* 2010;16:101-110.
13
14
15 91. Slykerman RF, Thompson JMD, Becroft DMO, Robinson E, Pryor JE, Clark
16 PM, Wild CJ, Mitchell EA. Breastfeeding and intelligence of preschool
17 children. *Acta Paediatrica, International Journal of Paediatrics* 2005;94:832-
18 837.
19
20
21
22
23
24
25 92. Steer CD, Smith GD, Emmett PM, Hibbeln JR, Golding J. FADS2
26 polymorphisms modify the effect of breastfeeding on child IQ. *PLoS ONE*
27 2010;5(7).
28
29
30
31
32 93. Taylor B, Wadsworth J. Breast feeding and child development at five years.
33 *Dev Med Child Neurol* 1984;26:73-80.
34
35
36 94. Temboury MC, Otero A, Polanco I, Arribas E. Influence of breast-feeding on
37 the infant's intellectual development. *J Pediatr Gastroenterol Nutr* 1994;18:32-
38 36.
39
40
41
42 95. Thorsdottir I, Gunnarsdottir I, Kvaran MA, Gretarsson SJ. Maternal body
43 mass index, duration of exclusive breastfeeding and children's developmental
44 status at the age of 6 years. *Eur J Clin Nutr* 2005;59:426-431.
45
46
47
48
49 96. Tozzi AE, Bisiacchi P, Tarantino V, Chiarotti F, D'Elia L, De Mei B, Romano
50 M, Gesualdo F, Salmaso S. Effect of duration of breastfeeding on
51 neuropsychological development at 10 to 12 years of age in a cohort of
52 healthy children. *Developmental Medicine & Child Neurology* 2012;54:843-
53
54
55
56
57
58
59
60

- 1
2
3 848.
4
5 97. Veena SR, Krishnaveni GV, Srinivasan K, Wills AK, Hill JC, Kurpad AV,
6
7 Muthayya S, Karat SC, Nalinakshi M, Fall CHD. Infant feeding practice and
8
9 childhood cognitive performance in South India. Arch Dis Child 2010;95:347-
10
11 354.
12
13 98. Whitehouse AJO, Robinson M, Li J, Oddy WH. Duration of breast feeding
14
15 and language ability in middle childhood. Paediatr Perinat Epidemiol
16
17 2011;25:44-52.
18
19 99. Wigg NR, Tong S, McMichael AJ, Baghurst PA, Vimpani G, Roberts R. Does
20
21 breastfeeding at six months predict cognitive development?. Aust N Z J Public
22
23 Health 1998;22:232-236.
24
25 100. Zaini MZ, Lim CT, Low WY, Harun F. Effects of nutritional status on
26
27 academic performance of Malaysian primary school children. Asia Pac J
28
29 Public Health 2005;17:81-87.
30
31 101. Zhou SJ, Baghurst P, Gibson RA, and Makrides M. Home
32
33 environment, not duration of breast-feeding, predicts intelligence quotient of
34
35 children at four years Nutrition 2007;23:236–241.
36
37 102. Koren G. Bias against the null hypothesis in maternal-fetal
38
39 pharmacology and toxicology. Clin Pharmacol Ther. 1997 Jul;62(1):1-5.
40
41 103. Koren G, Nickel C. Perpetuating fears: bias against the null hypothesis
42
43 in fetal safety of drugs as expressed in scientific citations. J Popul Ther Clin
44
45 Pharmacol. 2011;18(1):e28-32.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1
Flow diagram:



Author and Reference	Study Design	Target Population	Breast milk exposure	Cognitive development assessment tool and participant age	Results	Confounders adjusted for	Difference in IQ after adjustment	Study Quality Grading
Amanda et al 1992 (21)	Prospective cohort. Feeding method collected retrospectively	84 school aged children from two schools in India	Exclusive breast milk for more than 4 months vs. less than 4 months or formula fed	General mental ability test for children from 7-11 years (Srivastava and Saxena 1988-89)	There were significantly more breast fed (>4 m) children in the higher IQ category (IQ>109)	None.	Not provided	C
Andres et al 2012 (22)	Longitudinal study	391 healthy infants enrolled in the Beginnings Study	Breast fed vs. Soy fed vs. milk-based formula fed	Assessed at ages 3, 6, 9, and 12 months using the Bayley Scales of Infant Development second edition, from which the Mental Developmental Index (MDI) and Psychomotor Development Index (PDI) were derived	BF infants scored slightly higher than formula-fed infants on the Mental Developmental Index (MDI) score at ages 6 and 12 months (P < 0.05). Confounders included in the model.	Socioeconomic status, mother's age and IQ, gestational age, gender, birth weight, head circumference, race, age, and diet history	BF infants scored 1-2 points higher than formula-fed infants on the Mental Developmental Index (MDI) score at ages 6 and 12 months (P<0.05)	A
Angelsen et al 2001(23)	Prospective cohort	345 children	< 3 months 3-6 months > 6 months	Bayley Scales of Infant Development (BSID) at 13 months. Wechsler Preschool and Primary Scales of Intelligence (WPPSI-R), and Peabody Developmental Motor Scales (PDMS) at age 5y	Shorter duration of breast-feeding was associated with lower scores on mental developmental tests both at 13 months and at 5 years of age. Unadjusted difference of 8 points	Maternal age, education, smoking, and Raven score (IQ)	Maternal age, education, and intelligence were significant confounder. When analyzing performance IQ and verbal IQ separately, the median IQ value, was not statistically different when adjusting for maternal Raven score	B
Auestad et al 2001 (24)	Prospective randomized longitudinal study comparing different formula types and non randomized breast feeding group.	294 children from four sites in the United states	Breast feeding until age 12m, versus 3 different types of formulas (±AA, DHA)	Bayley scale for infant development at 6 and 12 months Fagan test of infant intelligence at 6 and 9 months MacArthur communicative development inventories at 9 and 14 months	No difference in any of the parameters checked between any study groups.	None.	No difference in any of the parameters checked between any of the study groups.	A

Auestad et al 2003 (25)	Follow up study of Auestad 2001: Prospective randomized longitudinal study comparing different formula types and a non randomized breast feeding group.	157 children from the original 294 children from four sites in the United states	Breast feeding until age 12m, versus 3 different types of formulas (\pm AA,DHA)	At 39 months, standard tests of IQ (Stanford Binet IQ), receptive vocabulary (Peabody Picture Vocabulary Test-Revised), and expressive vocabulary (mean length of utterance).	No difference in any of the parameters checked between any study groups.	None	No difference in any of the parameters checked between any study groups.	A
Bartels et al 2009 (26)	Retrospective cohort. Breast feeding status prospectively assessed	672 monozygotic male twin pairs 637 dizygotic male twin pairs 860 monozygotic female twin pairs 647 dizygotic female twin pairs 679 male-female twin pairs 598 female-male twin pairs	<Two weeks >Two weeks	Dutch CITO-elementary test at age 12 years.	Breast-fed children of highly educated mothers score on average 7.6 point higher on a standardized test of cognitive abilities than formula-fed children of mothers with a low education.	Maternal education, income.	A significant positive effect of breastfeeding on cognitive abilities above the expected positive effect of maternal education. Exact numbers not provided.	B
Bauer et al 1991 (27)	Prospective cohort	50 children from Honolulu, Hawaii	Breastfeeding as a continuous variable over time.	The McCarthy Scales of Children's Abilities at age 3 years.	The duration of breastfeeding was significantly correlated with scores on the scales, General, cognitive, Quantitative and Memory	Socio-economic status, gender, and pesticide exposure	Remained significant, numbers not provided	C
Birch et al 2007 (28)	Single-center, double-blind, randomized clinical trial comparing different formula types and non randomized breast feeding group.	52 healthy term infants enrolled for DHA and ARA supplementation and 32 breast-fed infants served as controls	Assigned diets were fed exclusively through 17 weeks of age. In the breast feeding group the average duration of breastfeeding was 43 \pm 9 weeks.	The Wechsler Preschool and Primary Scale of Intelligence, Revised (WPPSI-R) was used to assess intelligence at 4 years of age.	The control formula and DHA-supplemented groups had Verbal IQ scores poorer than the breast-fed group. There was no difference in performance or full-scale IQ between all groups.	None.	No adjustment done.	B

Bon 1980 (29)	Retrospective cohort	954 children from France	Exclusive bottle fed Vs. Breastfed for <15 days Vs. Breastfed between 15 days and 2 months; Vs. Breastfed more than two months.	The PM-47 Non-Verbal test, at 6-8.5 years of age.	Higher scores for girls who were breast-fed versus not breast-fed. No difference in boys.	Could not be assessed from the text.	Not performed.	C
Bouwstra et al 2005(30)	A prospective, double-blind, randomized control study comparing different formulas. Non randomized breast fed group as control.	A control formula - n=169, an LC-PUFA supplemented n=146 Breastfed group n=159	Supplementation - 2 mo. All formula-fed infants received control formula between 2 and 6 mo.	The Bayley Scales of Infant Development (BSIDII) at 18 months.	Bayley's MDI and PDI result values did not differ significantly between the three groups.	Parity, HOME score, parental Education	No difference between the groups.	B
Burruchaga et al 2000 (31)	Prospective cohort	39 children born at term and from homogeneous sociocultural status in Spain.	Exclusive breast milk for at least 2 months, vs. exclusive bottle-fed.	The Bayley Scales of Infant Development at 22 months.	Bayley's MDI and PDI result values did not differ significantly between the groups.	Maternal education, head circumference, maternal occupation, birth order, smoking status.	No difference between the groups.	C
Caspi et al 2007 (32)	Retrospective cohort	858 children from the New Zealand (Dunedin) birth cohort 1848 children from the British (E-risk study) twin birth cohort	No breastfeeding Vs. breastfeeding	New Zealand cohort: IQ measured at ages 7, 9, 11, 13 years using Wechsler Intelligence Scale for Children-Revised. IQ scores combined for an overall score. British cohort: revised short form of Wechsler Preschool and Primary Scale of Intelligence-age 5	Difference in IQ test scores between breastfed children and those not breastfed was 5.6 and 6.3 IQ points in the Dunedin and E-risk cohorts, respectively. Benefit mediated by a specific genotype. Only in children carrying the C allele.	Genetic variation in fatty acid metabolism (rs174575) Socioeconomic status, Maternal cognitive ability, Gestational age, Birth weight	Children not carrying the C allele did not benefit from breastfeeding.	A

1 2 3 4 5 6 7 8 9 10 11 12 13 14	Clark et al 2006 (33)	Prospective cohort	784 Chilean children	< 2months > 2 > 8 months > 8 months	At age 5.5 years. Wechsler Preschool and Primary Scales of Intelligence – Revised (WPPSI-R) Preschool Language Scale – 3rd Edition (PLS-3) the Broad Cognitive Abilities Standard Scale (BCA) of the Bateria Woodcock-Muñoz-Revisada (Bateria-R)	Poorer outcomes on the cognitive and language assessments were found for both the short and long extremes of breastfeeding as the sole milk source. The highest scores were observed in children who received breast milk as the sole milk source for 2–8 months.	Gender, birth weight, child's age at testing, maternal education, IQ, depression, age, father absence, paternal education, HOME score, socioeconomic status, nutritional status, iron deficiency anemia at 12 months, and iron supplementation.	Difference in IQ not significant.	B
15 16 17 18 19 20 21 22	Daniels et al 2005 (34)	Prospective cohort	1984 Filipino children	Any breastfeeding during: 0-6m 6-12m 12-18m 18-24m Over 24m	Philippines Nonverbal Intelligence Test of fluid abilities at ages 8.5 and 11.5 y	Poor education and suboptimal living conditions among BF mothers were strong negative confounders, causing inverse associations between BF and cognitive ability. Increased duration of any BF was of small significant benefit for cognitive development at both ages.	Parental education, paternal presence in home, maternal age, parity, alcohol during pregnancy, preterm status of child, mother reads, child's gender, baths (n/wk), dietary variety at 2 y, household income, non-income-producing assets, electricity in home, and environmental hygiene score.	1.6 points among normal birthweight breast-fed for 12 to 18 mo vs. less than 6 mo.	A
23 24 25 26 27	De Andraca et al 1998 (35)	Prospective cohort. Sub analysis of an RCT concerning iron fortified formulas	788 infants, 4-6 months of age.	<30 days vs. >30 days. All children in the study were breastfed for average of 75 days.	Bayley Scale of Infant Development – MDI and PDI at 12 months of age.	Breast-feeding for more than 30 days was associated with significantly lower scores (2.5 points less in MDI and 2.3 in PDI). Probably due to low SES.	SES, Education and occupation of parents, alcohol abuse, HOME, maternal intelligence by WAIS, stressful events.	No adjustment was done for the specific association of breastfeeding and Bayley scale.	C
28 29 30 31 32	De Andraca et al 1999 (36)	Prospective cohort	138 mother-infant dyad	Exclusive breast feeding for 6 months or more vs weaning before 45 days of age	Bayley Scale of Infant Development at 12 months of age	No difference between the groups in MDI and PDI	None; study reports similar family characteristics in each group.	No difference.	C
33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49	Der et al 2006 (13)	Database analysis of a prospective study, sibling pairs analysis, and metaanalysis	5,475 children 332 pairs of sibling discordant for breastfeeding status 545 discordant for duration of breastfeeding	Breastfeeding vs. no breastfeeding. Breastfeeding history obtained mostly within a year of birth	Peabody individual achievement test (PIAT) was administered to children between 5 and 14y	Unadjusted effect of breastfeeding +4.7 compared to non-breastfeeding.	Maternal IQ, education, age, family poverty, HOME stimulation score, and birth order.	After adjustment the difference became non-significant.	A

Di girolamo et al 1993 (37)	Prospective cohort. Breastfeeding data collected retrospectively	80 children	Full breast-feeding for at least 4 months vs. no breast feeding since 2 weeks of age.	Bayley Scales of Infant Development at 8-30 months	Breast-feeding group had a higher average MDI (of 6 points) compared to the bottle fed group, but no difference in PDI.	Parental education and age, gender, birth order.	Remained significant. Data not provided.	C
Eickmann et al 2007 (38)	Prospective cohort	191 Brazilian infants	'Predominantly breast fed' 'partially breast fed' and 'non-breast fed'	Bayley Scales of Infant Development II at 12 months of age.	Full breast-feeding at 1 month was associated with a small significant benefit in mental development compared with partial or no breast-feeding. No additional advantage in mental development was found with longer durations of full breastfeeding.	Adjusted for family income, possession of TV and fridge, flush toilet, maternal work and years of schooling, number of children under 5 years, home stimulation index, smoking during pregnancy, birth weight, infant's sex, hemoglobin, weight-for-age.	Full breast feeding at 1 month was associated with +3.0 points, P = 0.02 compared with partial or no breast feeding.	C
Elwood et al 2005 (39)	Prospective cohort. Breastfeeding data collected retrospectively	779 men from Caerphilly, South Wales, UK	Artificially fed vs. Breastfed, duration unknown vs. Breastfed <3m vs. breastfed >3m	Men aged 60–74 years. 1. The national adult reading test (NART) 2. The AH4 13 verbal and mathematical reasoning 3. The choice reaction time (CRT) for hand-brain reaction speed.	In the normal birth weight group the mean cognitive function was similar in both groups. In the men whose birth weight had been below the median, having been artificially fed was associated with significantly lower results in 2 of the 3 tests.	Age, social class, education, birth order and family size, father's social class, father's unemployment.	In the normal birthweight group, the adjusted mean cognitive function was similar in both groups.	B
Evenhouse et al 2005 (40)	Database analysis of a prospective study, sibling pairs analysis. Data from the National Longitudinal Study of Adolescent Health.	2,734 sibling pairs	No breastfeeding < 3 months 3-6 months 6-9 months 9-12 months 12-24 months >24 months	Add Health's abbreviated version of the Peabody Picture Vocabulary Test (PVT), normed for age and sex.	Persistent positive correlation between breastfeeding and cognitive ability. Unadjusted Difference in PVT 4.9 percentile points when comparing never breastfed to any breastfeeding in the full sample.	Birth weight, gender, birth order, parental investment, environmental characteristics	1.68 percentile points higher for ever breast fed to never breastfed.	A

Fergusson et al 1982 (41)	Prospective cohort	Birth cohort of children from the Dunedin Multidisciplinary Child Development Study: 1037 children assessed at age 3y 991 at age 5y 954 at age 7 y.	Child bottle- fed; Breast-fed for up to 4 months; Breast- fed for 4 months or longer.	Measures of intelligence at 3, 5 and 7 years. 3y-Peabody Picture Vocabulary Test 5y-Stanford Binet Intelligence Scale 7y-Weschler Child Intelligence Scale	Children who were breast-fed for 4 months or longer had scores which were 3.84 (average) points higher than bottle-fed infants (on scales with a standard deviation of 10).	Maternal intelligence (SRA verbal scale), maternal education, childhood experiences. Maternal training in child rearing, family socioeconomic status. Birth weight and gestational age.	A small but statistically significant benefit in the test scores of breastfed vs. bottle- fed infants (mean = 1.89).	A
Florey et al 1995 (42)	Retrospective cohort	592 first born singletons in Dundee	Breast fed on discharge from the hospital Vs. Bottle-fed.	Age 18 months. Bayley Scales of Infant Mental and Motor Development.	Higher mental development was significantly related to breast-feeding on discharge from hospital.	Partner's social class, mother's education, height, alcohol and cigarette consumption, placental weight and the child's sex, birth weight and gestational age at birth.	After adjustment, the difference in Bayley mental development index between the groups was between 3.7 and 5.7 units.	C
Foroushani et al 2010 (43)	Longitudinal cohort. Breastfeeding data collected retrospectively at age 2y.	5362 singletons born in 1946 in England, Wales and Scotland	No breastfeeding 1-3 months 4+ months	Age 8 – sentence completion, reading and vocabulary. Age 11 – Verbal, reading and vocabulary. Age 15 – Verbal, reading and vocabulary. Age 26 – reading Age 43 – visual and memory.	Children who were breastfed longer scored higher on verbal tests. Tests at older age (26, 43) were not significantly different.	Birth weight, childhood illness, home conditions, parents' age and education, child's behavioral scores, parents interest in the child's development, school type.	Mean score 1.5 points higher for ages 8,11,15. Not significant at older ages.	C
Gale et al 1996 (44)	Prospective cohort	994 men and women, born between 1920 and 1930 in Hertfordshire, UK	Exclusive breast feeding, exclusive bottle feeding, mixed feeding	65-75 years old. AH4 IQ test, taken on a computer.	Participants who had been exclusively breastfed had slightly higher IQ scores compared to the 2 other groups.	The use of a dummy in infancy, number of older siblings, father's occupational class, and mother's age at the participant's birth.	After adjustment, no association was found between adult intelligence, and method of infant feeding.	B
Gale et al 2010 (45)	Prospective cohort	241 children born to the Southampton Women's Survey, UK.	Breast feeding Fortified formula feeding Unfortified formula feeding	Age 4 – Wechsler Pre-School and Primary Scale of Intelligence (3rd ed.), sentence repetition and verbal fluency measured by the NEPSY	In unadjusted analysis children who were breastfed or fed with a fortified formula had significantly higher scores.	Maternal IQ and education, social class, on benefits, age at birth, birthweight.	After adjustment the differences in IQ between groups became non-significant.	A

Ghys et al 2002 (46)	Prospective cohort. Data on breast feeding collected retrospectively	128 full term children	Breast-feeding was expressed as the number of months of breast-feeding regardless of possible additional formula feeding.	At 4 years: Dutch adaptation of the Kaufman Assessment Battery for Children (K-ABC), the Groningen Developmental Scale (GOS). And items of the motor scale of the McCarthy Scales of Children's Mental Abilities.	Duration of breast-feeding showed significant correlations with cognitive development (Pearson correlation coefficient 0.26)	Plasma and RBC DHA and AA, maternal intelligence, birth weight, duration of breast-feeding and paternal educational attainment, smoking during pregnancy.	In the regression analysis the correlation disappeared.	B
Gibson-Davis et al 2006 (47)	Longitudinal birth cohort study. Breast-feeding information collected retrospectively at 1 year.	1645 American-born mothers and their babies.	Breastfeeding for at least 1 month vs none.	At three years of age: Peabody Picture Vocabulary Test-Third Edition	In unadjusted mean comparisons, breastfed children had Peabody Picture Vocabulary Test scores that were 6.6 points higher than children who were not breastfed.	An extensive set of demographic characteristics, including mother's Peabody Picture Vocabulary Test and the Home Observation for Measurement of the Environment score. Mothers were categorized into 1 of 3 educational-status groups.	After adjusting for demographic characteristics and maternal verbal ability, the coefficient dropped to 1.72.	A
Gómez-Sanchiz et al 2004 (48)	Prospective cohort, information on feeding collected retrospectively.	238 healthy babies born at term, non-IUGR.	Formula fed, Breast-fed up to 4 months, Breast-fed for more than 4 months.	Bayley Infant Development Scale at 24 months of age	Infants breast-fed for longer than 4 months scored higher on the mental development scale than those breast-fed for less time.	Socio-demographic and neonatal variables including parental IQ score.	The results of multiple linear regression analysis showed that infants breast-fed for longer than 4 months scored 4.3 points more than those breast-fed for less time.	A

Greene et al 1995 (49)	Retrospective cohort	432 subjects. 208 males, 224 females	Breast-fed vs. non-breast fed. And Breast fed for up to 12 weeks vs. more than 12 weeks.	Age: 11-16 years. Raven's Standard progressive matrices test and subsets of the Primary Mental Ability Test namely verbal meaning, reasoning and number facility.	The breastfed children showed a highly significant advantage over the non-breastfed children for all measures of IQ assessed, ranging from a 4.3 point advantage in Raven's IQ to a 6.0 point advantage in Primary mental abilities IQ.	Birth weight, gestational age, birth rank, child's sex, social class, mother's age and mother's educational level.	Following adjustment, the beneficial effect of breastfeeding (yes vs no) was statistically non-significant. A 6 point advantage in verbal IQ and 5.4 point advantage in reasoning IQ were observed for subjects breastfed for > 12 weeks, compared with less.	C
Gurka et al 2010 abstract (50)	Prospective cohort	1,050 children, from the National Institute of Child Health and Development Study of Early Child Care and Youth Development	Never; 0-6 months; longer than 6 months	Age: 4 years old. Standardized (mean.100; SD.15) cognitive outcomes.	Significant positive associations were observed between breastfeeding and cognitive outcomes before adjusting for other factors.	Maternal age, education, observed quality of the home environment, mother's attitude regarding modernity of parenting, and maternal verbal IQ	No difference.	A
Hart et al 2003 (51)	Prospective cohort	83 healthy full term infants	Exclusively breastfed vs exclusively non-breastfed	Brazelton Neonatal Behavioral Assessment Scale (BNBAS) at mean age of 8.95 days	Breast-fed infants surpassed formula-fed infants on the items of the orientation, motor, range of state, and state refulation dimensions of the BNBAS. Breast-fed infants also exhibited fewer abnormal reflexes, signs of depression, and withdrawal.	Socioeconomic status	After adjustment for SES only, the differences remained significant.	B

Hofer et al 1929 (9)	Retrospective cohort	383 children	Artificially fed, breast-fed for three months or less, from four to nine months, and from ten to twenty months.	Age – 7-13 years. Stanford Revision of the Binet-Simon intelligence test and the Pintner-Patterson performance scale (a nonverbal intelligence test), and by a group educational test, the Stanford achievements test.	Artificially fed were inferior in all standardized measurements to those breast fed from four to nine months, and, with one exception, to those breast fed three months or less. Artificially fed equaled or excelled those breast fed from ten to twenty months.	None; similar paternal IQ	Adjustment was not done although they mention paternal IQ to be similar between the groups.	C
Holme et al 2010 (52)	A secondary analysis of data from follow-up study of a RCT of an intervention to reduce smoking in pregnancy	1218 children	Not breastfed, Breastfed up to 2 months, 2-4 months, over 4 months. Also, any breastfeeding vs none	British Ability Scales (Total IQ, Visual IQ, and verbal IQ), and Quick Neurological Screening Test (QNST) at age 9 years.	Before adjustment, breastfeeding was significantly associated with higher total, verbal and visual IQ scores, and increasing duration was significantly correlated with IQ scores. Breastfeeding was associated with a crude total IQ increase of 5.49 points, which was reduced to 1.78 points on analysis with breastfeeding as a binary variable (yes/no – still significant)	Maternal demographics (including education, race, and age), smoking history, work patterns, depression, social support, neonatal details and ill health in the child	Total IQ became non-significant after adjustment	A
Horwood et al 1998 (53)	Longitudinal study – children studied at birth, 4 months, 1 year, at annual intervals to age 16, and 18 yrs	1064 children born in New Zealand in 1977	Not breastfed, breastfed for <4 months, breastfed for 4-7 months, breastfed for >=8 months	Revised Wechsler Intelligence Scale for Children (total IQ) – age 8 and 9 Teachers rating of school performance (reading and mathematics) – age 8 and 12 Progressive Achievement Test of Reading Comprehension – ages 10 and 12 Progressive Achievement Test of Mathematics – age 11 Tests of scholastic abilities – age 13 High School Outcomes	Increasing duration of breastfeeding was associated with consistent and statistically significant increases in cognitive abilities, and children who were breastfed for >=8 months had mean test scores that were between 0.35 and 0.59 SD units higher (more than 5 points) than children who were bottle fed.	Maternal age, maternal education, family socioeconomic status, averaged standard of living, averaged family income, maternal smoking during pregnancy, gender, birth order, birth weight	Upon adjustment associations were reduced and children who were breastfed for >=8 months had scores that were 0.11-0.3 SD units (less than 3 points) higher than children who were bottle fed.	A

Innis et al 1996 (54)	Retrospective cohort	433 full term infants born in 1994 in Vancouver	Never breastfed, breastfed less than 1 month, 1-3 months, 3-6 months, 6-8 months, more than 8 months, and mixed feeding (breast and formula milk)	Visual acuity measured using acuity card procedure with Teller Acuity Cards. Cognitive development measured using the Fagan Test of Infant Intelligence, (version 4.1) at 39 +/- 1 weeks of age	There were no differences in visual acuity or novelty preference among the infants when they were stratified by incidence or duration of breastfeeding.	None.	No difference.	B
Jacobson et al 1999 (55)	Prospective Longitudinal study	323 predominantly white, middle-class children born from 1980-81, at age 4, and 280 children at age 11, from 2 cohorts of similar demographic information *most exposed to PCB during pregnancy	Breastfed vs. not breastfed	McCarthy Scales of Children's Abilities and the Peabody Picture Vocabulary Test-Revised at the age of 4. Wechsler Intelligence Scale for Children-III, Wide Range Achievement Test-Revised, and the Woodcock Word, Passage, and Reading Comprehension test at the age of 11.	At both the ages of 4, and 11, breastfeeding was significantly related to higher IQ scores.	Social class, education, maternal IQ, parenting skills (Home observation for measurement of the environment (HOME))	The relationship was reduced to non-significance after adjustment for maternal IQ and parenting skills (HOME)	B
James et al, letter 1992 (56)	Prospective cohort	38 full-term children (taken as a sample from an extended Pembrokeshire farming family, UK)	Bottle-fed babies vs. breast-fed babies	IQ test At age 16 (no details)	No difference in IQ	No adjustment	No difference	C
Jedrychowski et al 2012 (57)	Longitudinal study	468 term babies in Krakow, Poland.	Complementary breastfeeding (including none), Vs. exclusive breastfeeding up to 3 months, 4-6 months, or longer than 6 months.	In the first 3 years of the follow-up the Bayley Mental Scales of Infant Development-second edition (BSID-II) were used. At the age of 6 and 7, the Wechsler intelligence test for children (WISC-R) was administered.	Children on mixed breastfeeding achieved lower total IQ scores at each of IQ check-ups compared with those who were exclusively breastfed.	Maternal education, baby's gender, parity, and weight gain in pregnancy.	Children breastfed exclusively for >6 months increased by 3.8 points (95% CI, 2.11-5.45).	B

Jiang et al 2011 (58)	Longitudinal study	3,271 children and their mothers from the US participating in the Child Development Supplement of the Panel Study of Income Dynamics	Yes or no ever breast feeding and, never breastfed; less than 6 months, 7–12 months, and more than 12 months.	Woodcock Johnson Psycho-Educational Battery-Revised (WJ-R) test score and Wechsler Intelligence Scale for Children-Revised (WISC-R) test score at 3 and 6 years of age.	Breastfed children had higher scores on WJ-R and WISC-R tests.	Child's age, race and ethnicity, sex, number of siblings, whether the child was first born to the mother, whether the child was born preterm, whether the child was born SGA, HOME scale, maternal IQ, age, education, health status, insurance, marital status, working, income.	Three out of the five effects remain significant; the effect sizes are smaller, with only one effect size being larger than one-fifth of a standard deviation. Longer spells of breastfeeding are uncorrelated with increases in the measures of achievement.	A
Johnson et al 1996 (59)	Longitudinal study	204 Euro-American full-term infants were followed up to the age of 3 from Galveston (Texas) area	Breastfed vs. not breastfed AND Duration of breastfeeding	At age 3 years; Stanford-Binet Fourth Edition and Peabody Picture Vocabulary Test-Revised.	Breastfeeding added significantly to the prediction of the Composite IQ Comprehension factor, Vocabulary, Absurdities, Memory for Sentences, and Peabody Picture Vocabulary Test-Revised. Duration of breastfeeding only added to the prediction of Vocabulary scores.	Socioeconomic status, HOME scores (parenting skills), mother's intelligence, mother's smoking behavior, gender, and birth order of the child	Breast feeding was associated with 4.6-point higher mean in children's Intelligence.	A
Keim et al 2009-abstract (60)	Prospective cohort	347 children	Exclusively breastfed vs formula fed	Mullen Scales of Early Learning at 1 yr of age	Infants exclusively breastfed demonstrated better visual reception, fine motor and overall cognitive development at 12 months than formula fed infants (4-6 points).	Preterm birth, smoking, race/ethnicity, education	Differences were weakened after adjustment. No numbers provided in abstract.	C

Kramer et al 2008 (15)	Cluster-randomized trial.	13,889 Belarussian children born between June 1996 and December 1997	Experimental group vs control group: Experimental group was encouraged to breastfeed and had greater levels of breastfeeding at 3, 6, 9 and 12 months (exclusive breastfeeding also 7 fold higher at 3 months)	Wechsler Abbreviated Scales of Intelligence and teacher evaluations of academic performance in reading, writing, mathematics, and other subjects at 6.5 yrs of age.	The experimental group had higher means on all of the Wechsler Abbreviated Scales of Intelligence measures, with cluster-adjusted mean differences of 7.5 points for verbal IQ, 2.9 points for performance IQ, and 5.9 points for full-scale IQ. Teachers' academic ratings were significantly higher in the experimental group for both reading and writing.	Maternal (and paternal) IQ, as well as all other demographic and confounding variables should be distributed randomly between the treatment groups and should not confound the treatment effect.	Cluster-adjusted mean differences of 7.5 points for verbal IQ, 2.9 points for performance IQ, and 5.9 points for full-scale IQ.	A
Lawlor et al 2006 (61)	Prospective cohort	3794 woman who delivered a singleton baby between 1981-84 in Brisbane, Australia	Never, <4 months, >= 4 months	Peabody Picture Vocabulary Test at age of 5 Raven's standard progressive matrices (Raven's SPM) and the Wide Range Achievements Test version 3 (WRAT3) at 14 years	Univariate analysis of breastfeeding vs IQ showed a significant difference between the breastfeeding groups (higher scores associated with longer breastfeeding up to 8.6 points difference).	Gender, maternal age, maternal ethnicity, maternal education, paternal education, family income, gravidity, maternal smoking, fetal distress, duration of the first and second stage of labour, mod of delivery, apgar scores at 1 and 5 min, birthweight for sex and gestational age (z score), height for age and sex (z score), BMI for age and sex (z score)	Significance remained with mean difference in IQ of 6.8 between never and over 4 months of breastfeeding.	A
Lucas et al 1999 (62)	Prospective double-blind, randomized controlled study (different formulas, breastfed control group not randomized)	447 healthy full-term children born in the UK between 1993-1995.	Breastfed for at least 6 weeks vs formula fed.	Bayley Scales of Infant Development II at 18 months.	No differences in overall developmental scores at 9 months or 18 months or in any subscale quotient at 9 months were found.	Sex, centre, maternal age, maternal education, maternal marital status, and social class	No difference with adjustment for potential confounding factors.	A
Maimaitiming et al 2007 (63)	Retrospective cohort	442 infants and children inhabited by Uygur, Han, or Kazak nationality in West China	Breast fed vs mixed feeding	Denver Developmental Screening Test at age up to 3.	There were no differences in scores between breastfeeding vs mixed feeding groups.	none	No difference.	C

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Makrides et al 2000 (64)	Prospective randomized double blind controlled. Different formulas. Breastfed group was used as control, but not randomized.	68 formula fed infants and 46 breastfed infants born at term of appropriate weight	Formula fed vs breastfed	Infant VEP acuity at 16 and 34 weeks Bayley's Scales of Development at 1 year and 2 years.	At 1 yr of age, MDI scores of breastfed and formula-fed infants were not different, but at 2 years of age, MDI scores of breastfed infants were higher than formula fed infants. PDI scores were similar.	Home screening questionnaire scores, occupational prestige, parental education, gender, birth weight, maternal smoking, birth order, feeding mode.	MDI score was higher at age 2 in the breastfed group. 95% CI 3-16.8.	A
Malloy et al 1997 (65)	Retrospective cohort	518 children born in 1978-79 in Washington DC who were exposed to chloride deficient formulas.	No breastfeeding (176) vs any time length of breastfeeding (342)	Weschler Intelligence Scale-Revised at 9 or 10 years old	Breastfed scored significantly higher on full scale IQ; Further analyses limited to those exclusively breastfed for the first 60 days failed to demonstrate any significant relationship.	Maternal education, paternal education and annual income	Following adjustment, the difference was not significant.	B
Martin et al 2011 (66)	Retrospective cohort	1431 children (twin siblings) from the greater Brisbane area	Exclusively formula fed, any breastfeeding between birth and 3 months, exclusively breastfed for 3 to 6 months, exclusively breastfed for 6 months or more	FSIQ (full scale IQ) assessed using the Multi-dimensional Aptitude Battery (MAB) at 16 years of age	Breastfeeding was significantly associated with FSIQ scores. No effect of duration of breastfeeding on FSIQ was found.	Socioeconomic status, paternal education, maternal education, gestational age, and birth weight.	The effect was no longer significant after adjustment.	A
McCrory et al 2011 (67)	Retrospective cross-sectional study	8568 school children in Ireland born between 1997 and 1998.	Breastfed vs not breastfed AND Never breastfed, <=5 weeks, 6-15 weeks, 16-25 weeks, 26+ weeks	Age 9 years. Vocabulary component of the Drumcondra Primary Reading Test-Revised and part 1 of the Drumcondra Primary Mathematics Test-Revised.	In unadjusted analysis, children who were breastfed scored 8.67 percentage points higher on reading and 7.42 percentage points higher on math. Evidence of dose-response relationship was weak.	Gender, birth weight, gestation period, NICU, primary and secondary carer's social class, primary carer's education level, household income, mothers age at birth, Irish/Non-Irish, number of children's books in the home.	After adjustment, remained significant but weakened: 3.24 and 2.23 percentage points in reading and math respectively	A

<p>Morales et al 2011 (68)</p>	<p>Retrospective cohort analyzing polymorphisms in genes encoding enzymes involved in LC-PUFA synthesis</p>	<p>Two population-based birth cohorts n = 400 mother-child pairs from INMA-Sabadell; and n = 340 children from INMA-Menorca</p>	<p>Different types of formula fed vs breastfed</p>	<p>Mental development was assessed at age 14 months using the Bayley Scales of Infant Development, first Edition – MDI only, and at age 4 years old by the Spanish version of the McCarthy Scales of Children’s Abilities (MCSA)</p>	<p>Children with variants associated with lower synthesis of LC-PUFA had higher scores when breastfed, while those with greater capacity to synthesize these fatty acids had higher scores regardless of breastfeeding practices.</p>	<p>Sex, child age, psychologist, quality of neuropsychological test, maternal education, breastfeeding, and use of gas stove at home.</p>	<p>Not being breastfed conferred an 8- to 9-point disadvantage in cognition among children with low FADS1 activity and in a 5-8 point disadvantage in cognition among children with low ELOVL5 activity.</p>	<p>A</p>
<p>Morris et al 1999 (69)</p>	<p>Prospective cohort</p>	<p>102 normal birthweight children born in Brazil at term.</p>	<p>Number of breastfeeds per day (recorded daily).</p>	<p>Bayley Scales of Infant Development at 6 months and 12 months (MDI and PDI).</p>	<p>Breastfeeding frequency over the first 4 weeks of life, but not later, was significantly associated with mental development (MDI) at 6 months of age.</p>	<p>Socioeconomic data: family income, a household resources index, a housing quality index, a water and sanitation index, and maternal and paternal literacy</p>	<p>An average of one extra breastfeed per day resulted in an increase of approximately one-quarter of a point. The effect was no longer apparent at 12 months of age.</p>	<p>B</p>
<p>Morrow-Tlucak et al 1988 (70)</p>	<p>Prospective cohort</p>	<p>229 children born in Ohio between 1981-1982</p>	<p>No breastfeeding, breastfeeding =<4 months, breastfeeding >4 months</p>	<p>Bayley Scales of Infant Development – Mental Development Index (MDI) at 6 months, 1 year, and 2 years of age</p>	<p>Significant differences in MDI scores between the three groups (longer breastfeeding = higher scores) at both 1 year and 2 years of age (no significant difference at 6 months).</p>	<p>Parent education (mean of both parents), maternal attitude (Authoritarian Family Ideology), maternal intelligence (PPVT-R), cigarette use, maternal age, rage, marital status, Home Observation for Measurement of the Environment (HOME) at age 1, HOME at age 2, exact age at time of testing</p>	<p>With covariate control, a small but significant relationship between duration of breastfeeding and Bayley MDI at 1 and 2 years was detected. Infants breastfed for 4 months or more scored on average 9 points higher compared to the bottle-fed infants.</p>	<p>A</p>

Mortensen et al 2002 (71)	Prospective longitudinal birth cohort	Mixed sample: 973 men and women All-male sample: 2280 men	Divided to five groups: < 1 month 2-3 months 4-6 months 7-9 months > 9 months	Wechsler Adult Intelligence Scale (WAIS) at a mean age of 27.2 years in the mixed-sex sample. Børge Priens Prøve (BPP) test at a mean age of 18.7 years in the all-male sample.	Duration of breast-feeding was associated with significantly higher scores on the Verbal, Performance, Full Scale WAIS IQs and BPP test.	Parental social status Parental education Single mother status Mother's height, age, and weight gain during pregnancy cigarette consumption during the third trimester number of pregnancies Gestational age birth weight Birth length Indexes of pregnancy Delivery complications	The results remained significant with 4.6 points higher IQ for those breast-fed over 9 months compared to less than 1 month.	B
Mukerji et al 1993 (72)	Cross-sectional study	100 children aged 0-3 years.	No breast-feeding or mixed. Exclusive breast-feeding for 4-9 months. Exclusive breastfeeding for more than 9 months	Developmental Screening Test (DST)	In the 4-9 months exclusive breast fed group 100% had average or above average IQ. In the two other groups this number was significantly lower.	None.	No adjustments were made.	C
Nassar et al 2011 (73)	Cross-sectional study	42 healthy infants	Breastfed, artificially fed, mixed feeding	Bayley scale of infant development-second edition (BSID-II) between 4-6 months of age.	No difference in MDI or PDI. There was significant difference only in total behavior rating scale (TBRS) and motor quality percentile rank.	No significant difference between groups in terms of age, sex, and socioeconomic standard (indirectly adjusted for with multiple regression)	Significant increase in mean adjusted TBRS and motor quality percentile rank.	B
Nelson et al 1995 (74)	Prospective cohort?	Term gestation infants	Breast fed or formula fed for 3 months	Teller Acuity Cards at 14d, 3, 4, 8, and 18 months Fagan Test of Infant Intelligence at 8, 10 and 12 months Bayley Scales of Infant Development at 4, 8, and 18 months	There were no significant differences between breast-fed and formula-fed infants in visual acuity at 14d, 3, 8, or 18 months, or recognition memory or the Bayley PDI or MDI at any age		Meeting abstract only available	C

<p>Niemela et al 1996 (75)</p>	<p>Prospective follow-up</p>	<p>726 children born between 1985-86 in Finland</p>	<p>Breastfed <5 months, breastfed >5 months (matched pairwise based on maternal education and sex)</p>	<p>Non-verbal Columbian Mental Maturity Scale (CMM), visual integration using the Beery test and active vocabulary by naming of pictures at 56 months</p>	<p>Children breastfed for 5 months or more attained higher scores in developmental tests (sig difference found in relation to the general cognitive capacity and visual motor integration). No evidence of any interaction between verbal development and breastfeeding to 5 months or more.</p>	<p>Groups matched pairwise based on sex and maternal education.</p>	<p>Maternal education and parental status correlated with all cognitive scores. Sex and breastfeeding correlated with scores of the general cognitive capacity and visual motor integration tests in multiple linear regressions.</p>	<p>B</p>
<p>Oddy et al 2003 (76)</p>	<p>Prospective cohort</p>	<p>1401 children at first follow up and 1283 children at second follow up from the Western Australian Pregnancy Cohort Study following 2860 children in Perth, Australia</p>	<p>Never breastfed, fully breastfed 0-4 months, 4-6 months, more than 6 months</p>	<p>Peabody Picture Vocabulary Test-Revised (PPVT-R) for receptive English vocabulary – verbal intelligence at 6 yrs Performance subtest by the Wechsler Intelligence Scale for Children – Third Edition (WISC-III) – Block Design Test at age 8yrs</p>	<p>On average, children breast fed for more than 6 months had mean verbal IQ scores that were 6.44 points higher and Block Design scores that were 1.13 points higher than children never breastfed (small but significant differences).</p>	<p>Gestational age, maternal age, maternal education, parental smoking and older siblings (all covariates that were significantly correlated with verbal IQ and the performance subtest or breastfeeding)</p>	<p>breast feeding for >6 months was associated with an increase in verbal IQ of 3.56 points. The Performance subtest was weakened and was no longer a significant difference after adjustment.</p>	<p>A</p>
<p>Oddy et al</p>	<p>Prospective</p>	<p>980 children from the</p>	<p>Breastfed less</p>	<p>Western Australian Monitoring</p>	<p>Continuous breastfeeding was</p>	<p>Gender, maternal age, maternal</p>	<p>Results were</p>	<p>A</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

2011 (77)	cohort	Western Australian Pregnancy Cohort Study following 2860 children in Perth, Australia	than 4 vs more than 4 months And Breastfed less than 6 vs more than 6 months	Standards in Education (WAMSE) scores in: math, reading, writing, and spelling at age 10yrs	significantly associated with an increase in scores with each additional month of breastfeeding for math, reading, writing, and spelling.	education, family income, marital status, parent looks at book with child at age 5, maternal country of birth	attenuated when adjusted for confounders. Significant interactions were found in math and spelling, revealing that boys were more likely to have improved academic scores if breastfed for a longer period.	
Paine et al 1999 (78)	Retrospective cohort	96 healthy full term Caucasian children from the Adelaide area in Australia	Duration of exclusive breastfeeding	Bayley Scales of Infant Developmental at age 10-14 months	Duration of exclusive breast-feeding significantly predicted mental development scores for boys, but not for girls. Duration of breast-feeding did not predict psychomotor development scores.	Duration of breast-feeding, parents occupational prestige, parents' education level and smoking habits, number of siblings, birth order, HSQ score, gestational age, birthweight, age of testing, maternal age and gender were considered as possible independent variables – independent variables with p < 0.02 included in the model – gender, maternal age, birthweight, and duration of breastfeeding.	None.	C
Pollock et al 1994 (79)	Prospective cohort	3838 children from the 1970 British	Wholly breast fed for more than	Human figure drawing score, copying design score and English	Significant difference found between groups at age 5 for the	Age father left full time education, age mother left full	All results adjusted for all	A

		Births Survey at full term and healthy birth weight	3 months vs. wholly bottle fed	Picture Vocabulary Test score at age 5. Dichotomised outcomes: Pictorial Language Test, Friendly Maths test, Edinburgh Reading test, Spelling Test, British Ability Scales (Word definitions, similarities, matrices, total) and Continuous outcomes: Pictorial Language test, Friendly Maths test, Edinburgh Reading test, Spelling test, and British Ability Scales (Word definitions, similarities, matrices, total) at age 10	English Picture Vocabulary test. Significant difference found for both dichotomized outcomes and continuous outcomes for the British Ability Scales and dichotomized outcomes for the Pictorial Language test at age 10.	time education, highest educational qualifications of mother, mother's smoking behavior during pregnancy, antenatal labour preparation class attendance, infant's place of birth	other independent predictors of breast-feeding: English Picture Vocabulary test: aOR 1.50. Dichotomized outcomes: Pictorial Language test aOR 1.49 Word definitions aOR 1.55 Similarities aOR 1.64 Total aOR 1.64 Continuous outcomes: Word definitions mean diff 3.5 Matrices mean diff 2.6 Similarities mean difference 3.0 Total mean difference 3.1	
Quinn et al 2001 (80)	Prospective cohort	3880 healthy children from the Mater Hospital-University	Never breastfed, < 3 weeks, 3-7 weeks, 7 weeks –	Peabody Picture Vocabulary Test Revised (PPVT-R) at 5 yrs. of age.	Significantly increasing scores were found between duration of breastfeeding and the PPVT-R	Birthweight, poverty, maternal education, maternal age, time in daycare/preschool, number of	After adjusting for a wide range of biological and	A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

		of Queensland Study of Pregnancy project	4 months, 4-6 months, still breast feeding at 6 months		scores.	children in household at 5 years, English speaking background for mother & father, and degree of infant stimulation	social factors, the adjusted mean for those breastfed for 6 months or more was 8.2 points higher for females and 5.8 higher for males than compared to those never breastfed (this was significant).	
Rao et al 2002 (81)	Prospective cohort	299 children born in Norway/Sweden at appropriate size for gestational age (comparison group)	<12 weeks Vs. >12 weeks of breastfeeding AND Duration of breastfeeding as a continuum	Bayley Scales of Infant Development at 13 months of age Norwegian version of the Wechsler Preschool and Primary Scales of Intelligence – Revised at 5 yrs. of age Peabody Development Motor Scale measured at 5 yrs. of age	There were statistically significant differences in IQ between the 2-breastfeeding groups.	Site of enrolment, maternal education, maternal IQ, maternal smoking, admission to a neonatal intensive care unit, kindergarten attendance, gender and asymmetric intrauterine growth retardation	3.7 points for total IQ and 4.1 points for performance IQ. Results remained unaltered when adjusted for confounding variables.	B
Ribas-Fito et al 2007 (82)	Prospective cohort	391 children born in Spain between 1997-1999	0-2 weeks breastfeeding 2-20 weeks breastfeeding 20+ weeks breastfeeding	Spanish version of the McCarthy Scales of Children's Abilities (general cognitive scale, verbal scale, perceptual-performance scale, memory scale, quantitative scale, motor scale subsets) at 4 yrs. of age.	Children with longer periods of breastfeeding performed significantly better on the McCarthy cognitive scale (except motor).	Gender, academic trimester at examination, psychologist, maternal social class, maternal education, and maternal use of alcohol and tobacco during pregnancy	After adjustment for confounders, significance remained for the general cognitive scale, and a trend remained in other subsets.	A
Richards et al 2002 (83)	Retrospective cohort	1741 people from the MRC National Survey of Health and	Never Up to 2 months 2-7 months	National Adult Reading Test (NART), word-list learning task (verbal memory) and timed letter	Educational Attainment: Odds of obtaining higher qualifications by the age of 26	Early background variables: Sex, father's social class, mother's education, birth order,	Effect of BF on education was strengthened	A

		Development (NSHD), also known as the British 1946 birth cohort	7+ months	search task (mental speed and concentration) at age 53.	were statistically significant, with greater odds for longer breastfeeding. Cognitive Function: Regression coefficients for the unadjusted effect show that breastfeeding was significantly associated with the NART and verbal memory tests.	parental interest in education and material home conditions.	after adjusting for the early background variables and remained significant. No longer significant with further adjustment for cognitive ability at age 15. Cognitive function: Coefficients were reduced after adjusting for the early background variables and were further reduced after adjusting for educational attainment and adult social class Only NART remained sig.	
Richards et al 1998 (84)	Retrospective cohort	511 first born offspring of the British 1946 cohort and their parent	Duration of breastfeeding.	Sentence completion test, reading test and vocabulary test at age 8 yr.	Breastfeeding was positively associated with cognitive function at age 8 in the first offspring of a national birth cohort. This	Social class, parental educational attainment, material home conditions, maternal age at birth, birth order, family size,	Non-significant; after adjusting for social class, maternal	C

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

					association was not evident in the subsample of mothers of these offspring. Association in the offspring cohort became non-significant after adjusting for social class, maternal education, or maternal cognitive performance.	maternal cigarette smoking, parental interest in education, attendance at nursery school, whether offspring cohort members had been taught cognitive skills at age 4, and cognitive test scores of the mothers.	education, or maternal cognitive performance.	
Rodgers et al 1978 (85)	Longitudinal study	1464 children at age 8 and 1398 children at age 15 from the British 1946 cohort	Never bottle-fed vs. never breastfed	Picture intelligence test and word reading test at 8 years of age Non-verbal ability, mathematics, and sentence completion tests at age 15 years of age	A preliminary analysis indicated that low scores were more likely for those who had been bottle-fed than breast-fed. The mean sentence completion scores between the two groups is statistically significant. Every test at both ages was significant except for word reading scores after correction for background factors.	Sex, social group, parental interest in education, material home conditions, sample stratification, father's education, mother's education, family size, birth rank, age at weaning	After correction for confounders, every test at both ages was significant except for word reading scores.	B
Rogan et al 1993 (86)	Prospective cohort	855 newborns being followed in North Carolina were enrolled between 1978-1982 and	Bottle fed 'short breastfeeding' 'medium breastfeeding'	Bayley Mental and Psychomotor Development Index at 6, 12, 18, and 24 months of age. McCarthy General Cognitive, Verbal, Quantitative, Memory,	Bayley Mental and Psychomotor: After adjusting for co-variables, the results at all four time points were similar, differences among the groups were only statistically	Age, race, occupation, education, smoking, drinking, child's sex, birthweight, number of older siblings, prenatal PCB exposure and	Confounders integrated into model; unadjusted results not	B

		followed up to 5 yrs old	'long breastfeeding' 'very long breastfeeding'	Perceptual Performance, and Motor scales at 36, 48, and 60 months of age Report Card Grades from 3 rd , 4 th and 5 th grade (averaged)	sig. at 24 months. McCarthy: All scales showed trends towards higher scores with increasing length of breastfeeding, but the relationship was weakest for the motor scale. Differences after adjustment were only significant at 3 and 4 years (marginally at 5 years) between length of breastfeeding groups Report Cards: Showed slight increase with breastfeeding. Marginally significant after adjustment for English scores and not significant for math.	dichlorodiphenyl dichloroethene exposure , Identity of the examiner.	shown.	
Santiago Burruchaga et al 2000 (87)	Prospective cohort	39 children born at term and from homogenous socio-cultural status	Breastfed for at least 2 months vs. formula fed	Bayleys scale at 2 years of age	No statistically significant differences between groups were found in cognitive function.	Maternal age, level of education, occupation, number of children in the family, smoking.	None; confounders integrated into model.	C
Silva et al 2006 (88)	Longitudinal population based cohort	9367 children from the 1970 British Cohort Study comprising individuals born during April 5-11 in	Never breastfed Less than 1 month 1-3 months More than 3 months	British Ability Scale, the Shortened Edinburgh Reading Test (word recognition), the Friendly Math Tests, and the Pictorial Language Comprehension Test at age of 10 yrs.	Breast-feeding showed a positive association with cognition at 10 years before adjustment.	Socioeconomic class, birth weight, parity, gestational age, maternal age and maternal smoking	Breastfeeding was weakly associated with cognitive function after adjustment	A

		the UK					(standardized coefficient 0.07). However, this effect was much smaller in the structural equation model adjusting for the same variables and did not reach significance level, suggesting that it is of little clinical importance.	
Silva et al 1978 (89)	Retrospective cohort	1037 children from the Dunedin Multidisciplinary Child Development Study	< 1 week 1-4 weeks 5-12 weeks 13-24 weeks 25-36 weeks 37-51 weeks 51+ weeks	Gross motor co-ordination, fine motor co-ordination, verbal comprehension and verbal expression, intelligence, child behavior problems at age 3 yrs.	Comparison of the groups resulted in only one significant difference among 96 comparisons made. No significant differences in age of attainment of milestones, gross or fine motor ability scores, verbal comprehension or expression, ability, intelligence, the incidence of separation problems, hyperactivity, height, weight and head circumference.	Socioeconomic class, general mental ability (IQ), level of education. Pairwise comparison with group matching to account for confounders.	None; pairwise matching said to account for confounders.	B
Sloan et al 2010 (90)	Cross-sectional observational. Breastfeeding data were obtained retrospectively.	137 infants and mothers	Breast fed (defined as more than 1 month) vs. not breast fed (defined as less than a month a not at all)	Bayley Scales of Infant Development (second edition; Bayley, 1993), between 10 and 18 months old.	Mean cognitive scores were significantly higher in breast-fed infants (110) compared with formula-fed infants (105). Breastfeeding duration had a positive linear association with cognitive scores at one year.	Home environment (HOME score) and infant gender	The adjusted standardized beta for breastfeeding was 0.285, which remained significant after adjustment.	C
Slykerman 2005 (91)	Cross-sectional observational. Breastfeeding data were obtained retrospectively.	550 infants, 50% of which were SGA at birth and 50% AGA.	Not at all, less than 6 m, 6 to 12 m, >12 m.	Stanford Binet Intelligence Scale, 4th Edition at age 3.5y-4y	Breastfeeding was not significantly related to intelligence scores in the AGA group.	Examiner administering the intelligence test, gestation, gender, maternal education, marital status, parental occupation, maternal age, parity, and smoking.	No difference in IQ.	B

Steer et al 2010 (92)	Prospective observational. Breastfeeding data were obtained prospectively.	9656 children from the Avon Longitudinal Study of Parents and Children cohort.	Breastfeeding within the first month of life vs. never breastfeeding	Wechsler Intelligence Scale for Children, 8 years old.	Breastfeeding showed a strong association with full-scale IQ with breastfed children scoring 8 points higher IQ on average in unadjusted analyses.	Maternal education, paternal social class, low birth weight, pre-term gestation, home environment, parenting and gender	The breastfeeding effect attenuated to a 3- point advantage after adjustment.	B
Taylor et al 1984 (93)	Longitudinal population based cohort. Breastfeeding data were obtained retrospectively.	13,135 children from The Child Health and Education Study	Never breastfed Less than 1 month 1-3 months More than 3 months	English picture vocabulary test (EPVT) adapted from the American Peabody Picture Vocabulary Test, at 5 years of age.	Children breast-fed for three or more months scoring over one-quarter of a standard deviation above the norm.	The age of the child at testing; the child's sex and birthweight; whether there were older or younger siblings in the home when the study child was five years old; home furnishings and equipment; maternal age at the child's birth; maternal smoking and the social index	Breast feeding remained a significant influence on EPVT scores, but the difference between the groups was small: +0.12 in the standardized EPVT score	A
Temboury et al 1994 (94)	Prospective cohort	229 infants	Breast fed – at least three months Bottle-fed – none or less than 1 month	Bayley's scale at age 18-29 months.	Bottle fed infants had lower IMD scores (index of mental development)	Maternal age, education and social class, job, psychosocial risk, number of children, infants shyness, tantrum, hyperactivity, gender, birthweight, height, place of birth	The result remained significant.	C
Thorsdottir et al 2005 (95)	Longitudinal cohort	85 children in Iceland	Duration of exclusive breastfeeding	The Icelandic developmental inventory at age 6 years. Information collected from the mothers.	Duration of exclusive breastfeeding, in months, was positively related to children's total developmental index.	Maternal and paternal education, and family income, birthweight, maternal BMI.	Total developmental index (B=1.07±0.5, P=0.04)	C
Tozzi et al 2012 (96)	Prospective cohort	1403 children originally included in an Italian clinical trial on acellular Pertussis vaccines	Duration of exclusive breastfeeding both as a continuous variable and as a	An estimated IQ was obtained from scores of the vocabulary, similarities, block design, and coding tests at 10-12 years of age. An estimation of IQ was obtained from the scores of the four WISC-R	Scores of only a few neuropsychological tests were affected by exclusive breastfeeding duration: Mean scores on vocabulary, similarities, the Boston naming test, and	Sex, birthweight, gestational age, mother's age at birth, type of delivery, family composition, parents' education, presence of chronic diseases, current	In the regression analysis, the score on one subcategory of the California verbal learning	A

			categorical variable (<6mo, >6mo).	subtests.	estimated IQ improved with the duration of breastfeeding, whereas performance in one of the subtests for writing praxis decreased with breastfeeding duration.	prescription of antihistamines or antiepileptic drugs, and the amount of thimerosal to which children were exposed through vaccines.	test was negatively associated with breastfeeding for longer than 6 months. No difference was detected in any of the other test scores included in the analysis	
Veena et al 2010 (97)	Longitudinal cohort	514 children from the Mysore Parthenon birth cohort in south India	Six categories from <3 to ≥18 months	Kaufman Assessment Battery at 9- to 10-year-old children	Within this cohort, in which prolonged breastfeeding was the norm (90% breastfed ≥6 months), there were no associations between longer duration of breastfeeding and cognitive function.	Age, sex, gestation, birth size, maternal age, parity, socio-economic status, parents' attained schooling and rural/urban residence.	No difference either unadjusted or after adjustment.	B
Whitehouse et al 2011 (98)	Longitudinal cohort, breast feeding data collected prospectively	1195 live born children recruited at approximately 18 weeks gestation, Western Australian Pregnancy Cohort (Raine) Study.	(1) Never breast-fed, (2) Breast-fed predominantly for <4 months, (3) Breast-fed predominantly	Peabody Picture Vocabulary Test – Revised (PPVT-R) at age 10 years. Raw scores are transformed to standard scores, based around a mean of 100 and a SD of 15.	Strong positive association between the duration of predominant breast-feeding and PPVT-R at age 10 years.	Maternal age at conception, maternal education, family income, and the presence of the biological father in the family home, maternal smoking and alcohol consumption during pregnancy, maternal experience	Children who were predominantly breast-fed for >6 months had a mean PPVT-R score that was	A

			for 4–6 months (4) Breast-fed predominantly for >6 months.			of stressful events during pregnancy, parity, gestational age, child’s sex, and proportion of optimal birthweight, a measure of the appropriateness of fetal growth, Family functioning (MMFAD), Parenting Scale, language-learning environment.	4.04 points higher than children who were never breast-fed	
Wigg et al 1998 (99)	Longitudinal cohort, breast feeding data collected prospectively	375 children born in Port Pirie, South Australia.	At 6 month of age: Breast fed vs. bottle-fed. Children who were breast-fed for less than 6 months would have been classified in our analysis as bottle-fed.	Bayley Scales of Infant Development at age 2, the McCarthy Scales of Children's Abilities age 4 and Wechsler Intelligence Scale for Children at age 7,11,13.	Estimated unadjusted advantages for the breast-fed children at ages 2, 4 and 7 years were 5.5 points, 4.6 points and 4.3 points in MDI, GCI and IQ scores, respectively. At age 11 to 13 years, the unadjusted advantage for the breast-fed children was 3.8 points in IQ.	Gender, maternal age at delivery, socioeconomic status, HOME score, maternal IQ, parental smoking habits, birthweight, birth rank, lifetime average blood lead concentration and whether parents were living together.	Adjusting for the covariates diminished the association of the feeding method in infancy with cognitive development. The covariates contributing most to this attenuating effect were the HOME scores, maternal IQ, socioeconomic status and parental smoking habits.	B
Zaini et al 2005 (100)	Retrospective cohort	1397 children from Selangor Malaysia	<6 months >6 months	Raven’s Colored Progressive Matrices at a mean age 9.6 years	Those who were breastfed <u>less</u> than six months performed better (31.01 vs. 30.63 out of max 36)	None.	None.	C
Zhou et al 2007 (101)	Prospective cohort	302 children born between 1998 and 1999 in Adelaide, Australia	Not breast fed, Breast fed at hospital discharge,	Stanford-Binet Intelligence Scale at 4 years of age.	There was no association between duration of breast-feeding and childhood IQ in this relatively well-nourished cohort from an	Birth order and sex of the child, maternal smoking in pregnancy, parental education, parental occupation, and quality of	There was no association between breast-feeding and	B

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

			Less than 6 months, More than 6 months.		industrialized society. Before adjustment, children who were breast-fed for at least 6 m. had higher IQ than those who were breast-fed for less than 6 months	home environment	childhood IQ.	
--	--	--	--	--	---	------------------	---------------	--

For peer review only



PRISMA 2009 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	6
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6-7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	6
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6-7
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	8-9
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2 for each meta-analysis).	



PRISMA 2009 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10+flow diagram
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Table
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Table
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	13
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	13
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	12-14
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	3

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.



Breast milk and cognitive development – the role of confounders: A systematic review

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-003259.R1
Article Type:	Research
Date Submitted by the Author:	11-Jul-2013
Complete List of Authors:	Walfisch, Asnat; Hospital for Sick Children, Motherisk Program Sermer, Corey; Hospital for Sick Children, Motherisk Program Cressman, Alex; Hospital for Sick Children, Motherisk Program Koren, Gideon; Hospital for Sick Children, Motherisk Program
Primary Subject Heading:	Obstetrics and gynaecology
Secondary Subject Heading:	Paediatrics, Reproductive medicine, Public health
Keywords:	Developmental neurology & neurodisability < PAEDIATRICS, PERINATOLOGY, PUBLIC HEALTH, REPRODUCTIVE MEDICINE

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3 Breast milk and cognitive development – the role of
4
5 confounders: A systematic review
6
7
8

9
10 **Contributing Authors:**

11 Asnat Walfisch MD¹, Corey Sermer², Alex Cressman MSc¹, and Gideon Koren MD
12 FACMT FRCPC¹
13

14
15
16
17 **Author Affiliations:**

18
19 ¹ The Motherisk Program, Hospital for Sick Children; Department of Pediatrics and
20
21 Pharmacology, University of Toronto, Toronto, Ontario, Canada
22

23 ²University of Western Ontario, London, Ontario, Canada
24
25
26

27
28 **Corresponding author (and reprint requests):**

29 Gideon Koren, MD, FACMT, FRCPC

30
31 Director of Motherisk Program

32
33 Hospital for Sick Children, 555 University Avenue, M5G 1X8

34
35 Toronto, Ontario, Canada

36
37 Phone: 416-813-6780

38
39 Fax: 416-813-7562

40
41 E-mail: gidiup_2000@yahoo.com
42

43
44
45 **Word count:**

46
47 Abstract: 268 words

48
49 Manuscript: 2721 (excluding references)

50
51 Running title: Breast Milk and Cognitive Development – Walfisch et al
52
53
54
55
56
57
58
59
60

ABSTRACT:

Objectives: The association between breastfeeding and child cognitive development is conflicted by studies reporting positive and null effects. Consequently, debate remains whether such a relationship is confounded by factors associated with breastfeeding, specifically maternal socioeconomic class, IQ and education.

Methods: To try to resolve this debate, we conducted a systematic review of the literature investigating the association between breastfeeding and cognitive outcomes of healthy infants born at term.

Results: Eighty-four studies met our inclusion criteria with 34 rated as high quality, 26 as moderate and 24 as low quality. Critical assessment of accepted studies revealed the following associations: 21 null, 28 positive, 18 null after adjusting for confounders, and 17 positive - diminished after adjusting for confounders.

Directionality of effect did not correlate with study quality, however studies showing a decreased effect after multivariate analysis were of superior quality compared to other study groupings (i.e. 14/17 high quality, 82%). Further, studies that showed null or diminished effect after multivariate analysis corrected for significantly more confounders (7.7 ± 3.4) as compared to those that found no change following adjustment (5.6 ± 4.5) ($p = 0.04$). The majority of included studies were done during childhood ($>1y, <18y, 75%$) and set in high-income countries (85.5%).

Conclusion: The results of this systematic review support a conclusion that much of the reported effect of breastfeeding on child neurodevelopment may be due to confounding. Although it is unlikely that additional work will change the current synthesis, future studies in this field should attempt to rigorously control for all important confounders. Alternatively, study designs using sibling cohorts discordant for breastfeeding may yield more robust conclusions.

1
2
3 **Article summary:**
4

5
6 **Article focus:**
7

- 8
9
10
11
12
13
14
15
16
17
18
19
20
21
- Although most published data supports the association between breast-feeding and IQ of the offspring, debate remains whether this is a causal relationship or an association with favorable parental characteristics.
 - We conducted a systematic review of the literature investigating the association between breastfeeding and cognitive outcomes of healthy term infants.

22 **Key messages:**
23

- 24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
- Over eighty studies addressing this issue were published with conflicting results.
 - Studies, where the initial positive effect of breastfeeding on IQ disappeared or diminished after multivariate analysis, controlled for significantly more confounders than studies showing no such change.
 - Much of the reported effect of breastfeeding on child cognitive abilities is due to maternal cognitive and socioeconomic effects.

40 **Strengths and limitations of this study:**
41

- 42
43
44
45
46
47
- The significant heterogeneity in study design and rigor precluded the conduct of a formal meta-analysis.

48
49 **Funding**
50

51 This research received no specific grant from any funding agency in the public,
52 commercial or not-for-profit sectors.
53
54
55
56
57
58
59
60

INTRODUCTION:

Breastfeeding confers a range of nutritional and immunological advantages upon infants including reduction in childhood illness,¹⁻⁵ diabetes,⁶⁻⁷ and obesity.⁸

The potential of breast milk to enhance cognitive development has been the focus of numerous studies since Hoefler and Hardy's initial observation in 1929.⁹ It is generally agreed that children who breast-feed are more intelligent, however debate remains whether this is a causal relationship or merely an association with favorable parental socioeconomic class and IQ. The beneficial effects of breastfeeding on the child's neurodevelopment are hypothesized, by some, to be mediated by long-chain polyunsaturated fatty acids (LCPUFAs)¹⁰ which are present in human milk, but not in cow's milk or most infant formulas.¹¹ However, a recent systematic review of all randomized trials where mother's diet was supplemented with PUFA's during pregnancy has failed to confirm such an effect.¹²

The pendulum of opinion has swung back and forth with different investigators showing inconsistent results depending upon study design and rigor. The Achilles heel of most of these studies, and the probable explanation for the conflicting results, is the difficulty in controlling for confounders that may affect child development. Furthermore, the ability to clarify this relationship is hindered by ethical considerations, which preclude randomized controlled trials, given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

Well-established confounders in breastfeeding research include demographic and IQ differences between mothers who breastfeed and those who chose not to.¹³ Parents who score high on a range of cognitive abilities have children with above average IQ scores.¹³ In parallel, advantage in mother's IQ more than doubles the odds of

1
2
3 breastfeeding.¹³ Thus, some of the published data demonstrates the disappearance of
4
5 the breast-feeding effect on child's cognition after correction for maternal IQ.
6

7
8 In an attempt to partially overcome these sources of bias, a few randomized trials
9
10 have been published, with randomization to breastfeeding promotion¹⁴⁻¹⁵ or in preterm
11
12 infants.¹⁶ In the breastfeeding promotion intervention trial (PROBIT Trial) by Kramer
13
14 and colleagues, IQ scores and academic performance tests were more favorable in the
15
16 intervention group, however, statistically significant differences were only shown for
17
18 some of the sub-scores.¹⁵
19

20
21 Systematic reviews examining the impact of breastfeeding on cognitive abilities have
22
23 reached conflicting results.^{13, 17-19} The Meta-analysis by Jain et al¹⁸ suggests that less
24
25 than 25% of studies in this area have adjusted for socio-demographic confounders.
26

27
28 There is a paucity of literature critically assessing the current published evidence
29
30 within this field. In trying to address these challenges, the objective of the present
31
32 work was to conduct a systematic review of published studies investigating the
33
34 association between breast-feeding and neurodevelopmental outcome of healthy
35
36 infants born at term.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS:

The study was conducted based on a prospectively prepared protocol, using the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines.²⁰

Literature search:

Searches were conducted in the following databases (all from inception to July 2011): MEDLINE(R) with Daily Update, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, Health Technology Assessment, NHS Economic Evaluation Database, Embase and PsycINFO using the OvidSP interface, and on Science Citation Index Expanded (SCI-EXPANDED), Social Sciences Citation Index (SSCI), Conference Proceedings Citation Index- Science (CPCI-S), and Conference Proceedings Citation Index - Social Science & Humanities (CPCI-SSH) using the Web of Knowledge interface.

A search strategy was developed based on the MEDLINE, EMBASE and PsycINFO database subject headings and the "Used for" synonyms listed in the scope notes for the terms "breast feeding", "human milk", "breast milk", "infant formula", "artificial milk", "cognition", "intelligence" and "intelligent quotient". Reference lists of identified studies, textbooks, previously published systematic reviews, and review articles were also searched. No language restrictions were applied and studies in languages other than English were translated for incorporation into the study.

A second, complimentary, literature search was done on April 2013 for all studies published during the period of August 2011-December 2012.

Study selection:

Both prospective and retrospective studies (RCTs, non-randomized controlled clinical

1
2
3 trials, cohort studies, longitudinal studies, and case-control studies) were included if:

- 4
5 1. One of the study aims was to address the question of breastfeeding and cognitive
6 development;
7
8
9
10 2. The authors used reliable validated methods to evaluate cognitive development
11 [e.g. Bayley scales of infant development, Wechsler Intelligence Scale for
12 Children Revised (WISC-R)];
13
14
15
16 3. Prospective or retrospective documentation of use and duration of breastfeeding
17 was used; and
18
19
20
21 4. The authors focused on healthy term infants and not those at increased biological
22 risk for developmental delays (e.g. prematurity, intrauterine growth restriction).
23
24

25 Studies were excluded if:

- 26
27 1. The study group included preterm or small for gestational age babies.
28
29 2. Evaluation of the cognitive development was done using only a non-reliable or
30 subjective tool (such as school grades, or maternal report).
31
32
33 3. Dietary patterns and breastfeeding were not evaluated since birth.
34
35

36 Titles and abstracts were reviewed for possible exclusion by two reviewers (A.W. and
37 C.S.). If both reviewers excluded a citation, it was eliminated from further review. If
38 at least one reviewer included the citation or if there was insufficient information to
39 make a determination from the title and abstract, the full article was obtained for
40 review. Full text articles were reviewed by three authors (A.W., C.S. and A.C.) for
41 suitability for inclusion. Disagreements regarding study eligibility were resolved by
42 consensus.
43
44
45
46
47
48
49
50

51 Study quality grading

52
53
54 Quality assessment of individual studies was performed by two authors (A.W and
55 C.S.) using the three category summary grading system (A, B, C) suggested by Ip et
56
57
58
59
60

1
2
3 al.¹⁹ Their system defines a generic grading system that is applicable to each type of
4
5 study design including randomized controlled trials, cohort, and case-control studies
6
7 as follows:

8
9
10 **A (good)**

11 A study that adheres mostly to the commonly held concepts of high quality including
12
13 the following: clear description of the population, setting, interventions and
14
15 comparison groups; clear description of the comparison groups; appropriate
16
17 measurement of outcomes; appropriate statistical and analytic methods and reporting;
18
19 no reporting errors; less than 20 percent dropout; clear reporting of dropouts; and
20
21 appropriate consideration and adjustment for potential confounders.
22
23

24
25 **B (fair/moderate)**

26
27 Category B studies do not meet all the criteria in category A because they have some
28
29 deficiencies, but none of which are likely to cause major biases. The study may have
30
31 suboptimal adjustment for potential confounders. The study may also be missing
32
33 information, making it difficult to assess limitations and potential problems.
34
35

36
37 **C (poor)**

38
39 Category C studies either did not consider potential confounders or did not adjust for
40
41 them appropriately. These studies may have serious shortcomings in design, analysis
42
43 or reporting; have large amounts of missing information, or discrepancies in
44
45 reporting.
46

47 **Data extraction:**

48
49 Extracted data were compiled in an evidence table. The table includes a description of
50
51 the studies that addressed the key question according to the inclusion/exclusion
52
53 criteria. The table provides information about study design, target population and
54
55 sample size, description of breast-feeding exposure and method of categorization,
56
57
58
59
60

1
2
3 nature of the comparison group, cognitive development assessment tool used and
4
5 participants' age, summary of the results prior to adjustment for confounders, a list of
6
7 all confounders adjusted for, differences in IQ between the groups after adjustment
8
9 for possible confounders (if available), and study quality grading according to the
10
11 scale described above.¹⁹
12

13
14 Statistical analysis:
15

16 Comparison of studies based on their results or quality was performed by Chi Square
17
18 or ANOVA as appropriate.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS:

The flow of the literature search is displayed in Figure 1. Of the 1,696 potentially relevant citations identified, 84 studies met the *a priori* inclusion criteria for this systematic review (Table 1).^{9,13,15, 21-101} Overall agreement between reviewers on the inclusion of studies was 100%.

Out of these 84 publications, 34 were rated as high quality (grade A), 26 as moderate (grade B) and 24 as low quality (grade C). Overall, based on the primary endpoint of cognitive function, there were 21 studies showing no association between IQ and breastfeeding,^{24-25,30-32,35-36,39,54,56,62-63,73-74,86-87,89,91,97,100-101} 28 positive studies,^{9,15,21,27-29,37-38,42,48,51,57,59,61,64,68,71-72,75,79-82,85,90,94-95,98} 18 initially positive studies that became negative after accounting for confounders,^{13,23,26,33,43-46,49-50,52,55,65-66,69,78,84,96} and 17 studies where the initial positive effect was diminished but remained statistically significant after accounting for confounders (Table 2).^{22,34,40-41,47,53,58,60,67,70,76-77,83,88,92-93,99}

In general, the directionality of the results did not correlate with the studies' quality. However, the studies showing decrease in the effect after multivariate analysis were of superior quality compared to the rest (i.e. 14 of 17 had a quality score of A – Table 2).

Different studies corrected in their analyses for different potential confounders, ranging from zero to 16 total confounders (Table 1). Confounders commonly considered in these studies were socioeconomic status, maternal education, birth weight, gestational age, birth order and gender. Some considered quality and quantity of stimulation of the child to be crucial confounders but did not consider maternal or paternal intelligence and other important factors. Studies that showed null or diminished effect in their multivariate analysis controlled for significantly more

1
2
3 confounders (7.7 ± 3.4) as compared to those that found no change following
4
5 adjustment for confounders (5.6 ± 4.5 , $p = 0.04$). Furthermore, many of the studies
6
7 did not have a clear definition of breastfeeding or cumulative breast milk exposure.
8
9 Table 3 divides the included studies according to their settings: developed vs.
10
11 developing world. The majority of included studies were set in the developed world
12
13 ($71/84$, 85.5% vs. $13/84$, 15.5% in the developing world). The quality of the studies
14
15 set in developing countries were generally poorer given our criteria: 46% graded A +
16
17 B ($6/13$) in studies set in developing countries, compared with 76% ($54/71$) in studies
18
19 set in developed countries. Developing country studies were also more likely to reach
20
21 a null association or null association after adjustment for confounding compared with
22
23 developed country studies ($8/13$, 61% vs. $31/84$, 37% respectively).
24
25
26

27
28 A large variety of cognitive assessment tools were used and study outcomes were
29
30 measured anywhere from 8 days of age into adulthood. Table 4 divides the included
31
32 studies according to age groups of participants: infancy, childhood and adulthood,
33
34 with the corresponding direction of results and study quality. The majority of
35
36 included studies measured intelligence during the childhood period (age 1-18
37
38 years: $70/93$ studies, 75%). Studies performed during childhood and reaching an
39
40 initial positive association, weakened after adjustment, were generally of higher
41
42 quality than other studies ($12/14$ quality grade A, 86% , Table 4). Studies
43
44 performed during infancy or adulthood were more likely to find a null
45
46 association (before or after adjustment) compared with studies performed
47
48 during childhood (Infancy group - 61% , adulthood - 60% , childhood 43% , Table
49
50
51
52
53
54
55
56
57
58
59
60

The significant heterogeneity in study design and rigor precluded the conduct of a formal meta-analysis.

DISCUSSION:

The continuing debate of whether breastfeeding imparts direct advantage on child cognition, or whether this is merely an association with favorable familial socioeconomic status and cognition, is not purely theoretical. From a public health perspective, if breastfeeding has biological effects on a child's IQ - this will be one of very few cost-effective means to significantly improve a child's neurodevelopment. If, on the other hand, there is no such effect, in the case where breastfeeding is either impossible or not sought by the mother, this will allow these women to rest assured that their choice will not have long-term developmental consequences.

In the case of other comparable therapeutic dilemmas, conflicts are typically resolved through randomized controlled trials (RCT), which are not ethically feasible in this case given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

The closest comparison to a formal RCT in reducing selection bias would be sibling-pair analysis, when cognition of breastfed infants is compared to that of their siblings who were formula fed. This design ensures similar socioeconomic and maternal characteristics. Unfortunately, the few studies that have followed this design reached conflicting results.^{13, 40}

The second closest design to RCT was employed in the PROBIT study by Kramer and colleagues,¹⁴⁻¹⁵ who cluster randomized women in Eastern Europe to receive or not receive formal education about the advantages of breastfeeding. This study did show favorable effects, but it has been argued that the mothers randomized for the breastfeeding promotion arm might have been influenced not only in providing higher rates of breastfeeding, but also by improving other positive health behaviors.

1
2
3 Our analysis reveals that there are over eighty studies addressing this issue and that
4 their results divide almost evenly between positive and negative associations. The
5 quality of “positive” or “negative” studies did not differ, except for higher quality on
6 average in studies that showed an apparent decrease in effect after multivariate
7 analysis.
8

9
10
11 We have shown that studies, where the initial positive effect of breastfeeding on IQ
12 disappeared or substantially diminished after multivariate analysis, controlled for
13 significantly more confounders than studies showing no such change. When
14 compared to a meta-analysis conducted 14 years ago,¹⁷ it appears that many more new
15 studies did attempt to control for confounding measures of socioeconomic status and
16 parental education, amongst others.
17
18

19
20
21 Given that more tight control of confounders resulted in greater likelihood of
22 disappearance of breastfeeding effect, it can be argued that the remaining positive
23 effect reflects residual uncontrolled bias, as shown by Der and colleagues in their
24 large study.¹³ In that study, before adjustment, breastfeeding was associated with an
25 increase of around 4 points in mental ability. *Post hoc* analysis revealed that
26 adjustment for maternal intelligence accounted for most of this effect – where full
27 adjustment for a range of relevant confounders yielded a small (0.52) and non-
28 significant effect size (95% confidence interval -0.19 to 1.23).
29
30
31

32
33
34 In our systematic review, a similar effect was recorded by a total of 18 studies, and in
35 addition 17 studies showed substantially diminished effect after adjustment.
36
37

38
39
40 When we examined studies based on setting (Table 3), we found that the majority
41 of the 84 included studies were set in the developed world (85%). Studies completed
42 in mid-low income countries were nearly twice as likely to find a null association
43 (before or after adjustment) compared with studies set in developed countries (61%
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 vs. 43.5% respectively).

4
5 This may be due to the fact that in many low- and middle-income countries high rates
6
7 of some degree of breastfeeding exist¹⁰² and comparisons between breastfed and non-
8
9 breastfed populations may examine more homogenous study groups (i.e. parental
10
11 socioeconomic status, income, and parental IQ).³⁴ In contrast, studies originating in
12
13 the developed world exhibit a greater heterogeneity between breastfed and non-
14
15 breastfed populations¹³ as the choice to breast feed is associated with a family's
16
17 socioeconomic status, maternal education, maternal intelligence, and social
18
19 advantage.^{13,17-19} If a biological effect truly exists between breastfeeding and infant
20
21 IQ, one would expect this relationship to exist in multiple settings, including the
22
23 developing world. The fact that this relationship is less apparent in developing
24
25 countries suggests that much of the observed relationship may be due to parental
26
27 social advantage, confounding the choice to breast-feed.
28
29

30
31 This systematic review includes studies using a large variety of cognitive assessment
32
33 tools and age span. The majority of included studies measured intelligence during
34
35 the childhood period (age 1-18 years, 75%). Studies performed during infancy or
36
37 adulthood were more likely to find a null association (before or after
38
39 adjustment) although number of included studies is small. Possible explanations
40
41 for this finding include reduced accuracy of IQ evaluation in infancy (<1y) on one
42
43 hand, and a variety of additional factors influencing IQ at an older age (>18y), on
44
45 the other hand.
46
47

48
49 Another factor that needs to be seriously considered in our review is the
50
51 existence of bias against the null hypothesis. The likelihood of studies not
52
53 detecting a significant effect in pregnancy to be submitted and published in the
54
55 peer review literature is substantially lower than that of positive studies.¹⁰³⁻¹⁰⁴
56
57
58
59
60

1
2
3 This can create a distorted balance that may seriously affect the conclusions on
4
5 effects of interventions.
6

7
8 In conclusion, this systematic review suggests that much of the reported effect of
9
10 breastfeeding on child cognitive abilities is due to maternal cognitive and
11
12 socioeconomic effects. When considered together with the fact that a recent
13
14 systematic review failed to corroborate a biological effect of milk PUFA on brain
15
16 development, it is likely that breastfeeding does not, by itself, directly affect child
17
18 IQ.
19

20
21 Although it is unlikely that additional studies will change substantially the
22
23 current synthesis, future studies in this field should attempt to rigorously control
24
25 for all important confounders even if they are difficult to obtain (e.g. parental
26
27 IQ). Alternatively, study designs using sibling cohorts discordant for
28
29 breastfeeding may yield more robust conclusions to further clarify this dilemma.
30
31

32
33
34
35 Author's contribution:

36
37 Asnat Walfisch has participated in all phases of this study including literature
38
39 search, data collection and review, quality grading, interpretation of the results,
40
41 and has written a part of the manuscript and the revised manuscript.

42
43 Corey Sermer has participated in all phases of this study including literature
44
45 search, data collection and review, quality grading, and revision of the
46
47 manuscript.

48
49 Alex Cressman has participated in the literature search, data collection and
50
51 review, translation of foreign language manuscripts, and has taken a significant
52
53 part in the manuscript preparation and revision.

54
55 Gideon Koren participated in all phases of this study. He initiated the study and
56
57 supervised actively throughout its conduct. Specifically he was involved in the
58
59 data interpretation, statistical analysis, and has written and revised a substantial
60
61 part of the manuscript.

References

1. Ryan AS, Wenjun Z, Acosta A. Breastfeeding continues to increase into the new millennium. *Pediatrics*. 2002 Dec;110:1103-9.
2. Dell S, To T. Breastfeeding and asthma in young children: Findings from a population based study. *Archives of Pediatric and Adolescent Medicine* 2001;155:1261-1265.
3. Bhandari N, Bahl R, Mazumdar S, Martines J, Black RE, Bhan MK. Effect of community-based promotion of exclusive breastfeeding on diarrhoeal illness and growth: A cluster randomised trial. *Lancet* 2003;361:1418-1423.
4. Galton Bachrach VR, Schwarz E, Bachrach LR. Breastfeeding and the risk of hospitalisation for respiratory disease in infancy. A meta-analysis. *Archives of Pediatric and Adolescent Medicine* 2003;157:237-243.
5. Howie PW, Forsyth JS, Ogston SA, Clark A, Florey CD. Protective effect of breastfeeding against infection. *BMJ* 1990;300:11-16.
6. Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskeine E, Samuelsson U. Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood. *Diabetes Metabolism Research Reviews* 2004;20:150-157.
7. Young TK, Martens PJ, Taback SP, Sellers EAC, Dean HJ, Cheang M, Flett B. Type 2 diabetes mellitus in children prenatal and early infancy risk factors among native Canadians. *Archives of Pediatrics and Adolescent Medicine* 2002;156:651-655.
8. Owen CG, Martin RM, Whincup PH, Davey Smith G, Cook DG. Effect of infant feeding on the risk of obesity across the life course: A quantitative review of the published evidence. *Pediatrics* 2005;115:1367-1377.

- 1
2
3 9. Hoefler C, Hardy MC. Later development of breast fed and artificially fed
4 infants. *J Am Med Assoc.* 1929;92:615–20.
5
6
7 10. Koletzko B, Lien E, Agostoni C, Böhles H, Campoy C, Cetin I, Decsi T,
8 Dudenhausen JW, Dupont C, Forsyth S, Hoesli I, Holzgreve W, Lapillonne A,
9 Putet G, Secher NJ, Symonds M, Szajewska H, Willatts P, Uauy R, World
10 Association of Perinatal Medicine Dietary Guidelines Working Group. The
11 roles of long chain polyunsaturated fatty acids in pregnancy, lactation and
12 infancy: review of current knowledge and consensus recommendations. *J*
13 *Perinat Med* 2008;36:5–14.
14
15 11. Sauerwald TU, Demmelmair H, Koletzko B. Polyunsaturated fatty acid supply
16 with human milk. *Lipids.* 2001;36:991-6.
17
18 12. Lo A, Sienna J, Mamak E, Djokanovic N, Westall C, Koren G. The effects of
19 maternal supplementation of polyunsaturated Fatty acids on visual,
20 neurobehavioural, and developmental outcomes of the child: a systematic
21 review of the randomized trials. *Obstet Gynecol Int* 2012;2012:591531.
22
23 13. Der G, Batty GD, Deary IJ. Effect of breast feeding on intelligence in
24 children: prospective study, sibling pairs analysis, and meta-analysis. *BMJ*
25 2006;333(7575):945.
26
27 14. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzikovich I, Shapiro
28 S, et al for the PROBIT Study Group. Promotion of Breastfeeding Intervention
29 Trial (PROBIT): A randomised controlled trial in the Republic of Belarus.
30 *JAMA* 2001;285:413-420.
31
32 15. Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L,
33 Igumnov S, Fombonne E, Bogdanovich N, Ducruet T, Collet JP, Chalmers B,
34 Hodnett E, Davidovsky S, Skugarevsky O, Trofimovich O, Kozlova L,
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Shapiro S; Promotion of Breastfeeding Intervention Trial (PROBIT) Study
4
5 Group. Breastfeeding and child cognitive development: new evidence from a
6
7 large randomized trial. *Arch Gen Psychiatry* 2008;65(5):578-584.
8
9
10 16. Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and
11
12 subsequent intelligence quotient in children born preterm. *Lancet*
13
14 1992;339:261-4.
15
16 17. Anderson JW, Johnstone BM, Remley DT. Breastfeeding and cognitive
17
18 development: a meta-analysis. *Am J Clin Nutr* 1999;70:525-35.
19
20 18. Jain A, Concato J, Leventhal JM. How good is the evidence linking
21
22 breastfeeding and intelligence? *Pediatrics* 2002;109:1044-53.
23
24 19. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, Trikalinos T, Lau J.
25
26 Breastfeeding and maternal and infant health outcomes in developed
27
28 countries. *Evid Rep Technol Assess* 2007;153:1-186.
29
30 20. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting
31
32 systematic reviews and meta-analyses of studies that evaluate healthcare
33
34 interventions: explanation and elaboration. *BMJ*. 2009; 339:b2700.
35
36 21. Amanda M, Singh RP. A study of intelligence of children in relation to infant-
37
38 feeding practices and nutrition. *Indian Journal of Psychometry & Education*
39
40 1992;23:23-28.
41
42 22. Andres A, Cleves MA, Bellando JB, Pivik RT, Casey PH, Badger TM.
43
44 Developmental status of 1-year-old infants fed breast milk, cow's milk
45
46 formula, or soy formula. *Pediatrics*. 2012;129:1134-40.
47
48 23. Angelsen NK, Vik T, Jacobsen G, Bakketeig LS. Breast feeding and cognitive
49
50 development at age 1 and 5 years. *Arch Dis Child* 2001;85:183-188.
51
52 24. Auestad N, Halter R, Hall RT, Blatter M, Bogle ML, Burks W, Erickson JR,
53
54
55
56
57
58
59
60

- 1
2
3 Fitzgerald KM, Dobson V, Innis SM, Singer LT, Montalto MB, Jacobs JR,
4
5 Qiu W, Bornstein MH. Growth and development in term infants fed long-
6
7 chain polyunsaturated fatty acids: a double-masked, randomized, parallel,
8
9 prospective, multivariate study. *Pediatrics*. 2001;108(2):372-81.
10
11
12 25. Auestad N, Scott DT, Janowsky JS, Jacobsen C, Carroll RE, Montalto MB,
13
14 Halter R, Qiu W, Jacobs JR, Connor WE, Connor SL, Taylor JA, Neuringer
15
16 M, Fitzgerald KM, Hall RT. Visual, cognitive, and language assessments at 39
17
18 months: a follow-up study of children fed formulas containing long-chain
19
20 polyunsaturated fatty acids to 1 year of age. *Pediatrics*. 2003;112(3 Pt 1):e177-
21
22 83.
23
24
25 26. Bartels M, van Beijsterveldt CEM, Boomsma I. Breastfeeding, Maternal
26
27 Education and Cognitive Function: A Prospective Study in Twins. *Behav*
28
29 *Genet* 2009;39:616–622.
30
31
32 27. Bauer G, Ewald ES, Hoffman J, Dubanoski R. Breast feeding and cognitive
33
34 development in three-year-old children. *Arch Dis Child* 1991;64:763–4.
35
36
37 28. Birch EE, Garfield S, Castañeda Y, Hughbanks-Wheaton D, Uauy R, Hoffman
38
39 D. Visual acuity and cognitive outcomes at 4 years of age in a double-blind,
40
41 randomized trial of long-chain polyunsaturated fatty acid-supplemented infant
42
43 formula. *Early Human Development* 2007;83:279–284.
44
45
46 29. Bon N. Relationship between breastfeeding and subsequent IQ of child.
47
48 *Bulletin de Psychologie* 1980;33:935-947.
49
50
51 30. Bouwstra H, Dijck-Brouwer DAJ, Boehm G, Boersma ER, Muskiet FAJ,
52
53 Hadders-Algra M. Long-chain polyunsaturated fatty acids and neurological
54
55 developmental outcome at 18 months in healthy term infants. *Acta Pædiatrica*,
56
57 2005; 94: 26–32.
58
59
60

- 1
2
3 31. Burruchaga MS, Sanz JIR, Zubizarreta JIP, Fernandez JB, Crespo PS.
4
5 Intellectual development in the second year of life in healthy breast-fed
6
7 children vs. formula-fed children. *An Esp Pediatr* 2000; 52: 530-536.
8
9
10 32. Caspi A, Williams B, Kim-Cohen J, Craig IW, Milne BJ, Poulton R,
11
12 Schalkwyk LC, Taylor A, Werts H, Moffitt TE. Moderation of breastfeeding
13
14 effects on the IQ by genetic variation in fatty acid metabolism.
15
16 *Proc.Natl.Acad.Sci.U.S.A.* 2007;104:18860-18865.
17
18
19 33. Clark KM, Castillo M, Calatroni A, Walter T, Cayazzo M, Pino P, Lozoff B.
20
21 Breastfeeding and Mental and Motor Development at 5 1/2 Years. *Ambul*
22
23 *Pediatr.* 2006 ; 6(2): 65–71.
24
25
26 34. Daniels MC, Adair LS. Breast-feeding influences cognitive development in
27
28 Filipino children. *J Nutr* 2005;135:2589–2595.
29
30
31 35. de Andraca,I.; Pino,P.; de la Parra,A.; Rivera,F.; Castillo,M. Risk factors for
32
33 psychomotor development among infants born under optimal biological
34
35 conditions. *Rev Saude Publica* 1998;32:138-147.
36
37
38 36. De Andraca,I.; Salas,MI; Lopez,C.; Cayazzo,MS; Icaza,G. Effect of breast
39
40 feeding an psychosocial variables upon psychomotor development of 12
41
42 months old infants. *Arch Latinoam Nutr* 1999;49:223-231.
43
44
45 37. Di Girolamo D, Braibanti P, Mamone P. Feeding methods and cognitive
46
47 development in the first 3 years of life. *Rassegna di Psicologia* 1993;10:33-49.
48
49
50 38. Eickmann SH, de Lira PI, Lima Mde C, Coutinho SB, Teixeira Mde L,
51
52 Ashworth A. Breast feeding and mental and motor development at 12 months
53
54 in a low-income population in northeast Brazil. *Paediatr Perinat Epidemiol*
55
56 2007;21:129-137.
57
58
59 39. Elwood PC, Pickering J, Gallacher JEJ, Hughes J, Davies D. Long term effect
60

- 1
2
3 of breast feeding: Cognitive function in the Caerphilly cohort.
4
5 J.Epidemiol.Community Health 2005;59:130-133.
6
7
8 40. Evenhouse E, Reilly S. Improved estimates of the benefits of breastfeeding
9
10 using sibling comparisons to reduce selection bias. Health Serv Res
11
12 2005;40:1781-1802.
13
14 41. Fergusson DM, Beautrais AL, Silva PA. Breast-feeding and cognitive
15
16 development in the first seven years of life. Social Science and Medicine
17
18 1982;16:1705-1708.
19
20
21 42. Florey Du CV, Leech AM, Blackhall A. Infant feeding and mental and motor
22
23 development at 18 months of age in first born singletons. Int J Epidemiol
24
25 1995;24:S21-S26.
26
27
28 43. Foroushani AR, Mohammad K, Mahmoodi M, Siassi F. Effect of
29
30 breastfeeding on cognitive performance in a British birth cohort. Eastern
31
32 Mediterranean Health Journal 2010;16:202-208.
33
34
35 44. Gale CR, Martyn CN. Breastfeeding, dummy use, and adult intelligence.
36
37 Lancet 1996;347:1072-1075.
38
39
40 45. Gale CR, Marriott LD, Martyn CN, Limond J, Inskip HM, Godfrey KM, Law
41
42 CM, Cooper C, West C, Robinson SM. Breastfeeding, the use of
43
44 docosahexaenoic acid-fortified formulas in infancy and neuropsychological
45
46 function in childhood. Arch Dis Child 2010;95:174-179.
47
48
49 46. Ghys A, Bakker E, Hornstra G, Van den Hout M. Red blood cell and plasma
50
51 phospholipid arachidonic and docosahexaenoic acid levels at birth and
52
53 cognitive development at 4 years of age. Early Hum Dev 2002;69:83-90.
54
55
56 47. Gibson-Davis CM, Brooks-Gunn J. Breastfeeding and verbal ability of 3-year-
57
58 olds in a multicity sample. Pediatrics 2006;118: e1444-e1445.
59
60

- 1
2
3 48. Gomez-Sanchiz M, Canete R, Rodero I, Baeza JE, Gonzalez JA. Influence of
4 breast-feeding and parental intelligence on cognitive development in the 24-
5 month-old child. *Clin Pediatr* 2004;43:753-761.
6
7
8
9
10 49. GREENE LC, LUCAS A, LIVINGSTONE MBE, HARLAND PSEG,
11 BAKER BA. Relationship between Early Diet and Subsequent Cognitive
12 Performance during Adolescence. *Biochem Soc Trans* 1995;23:S376-s376.
13
14
15 50. Gurka KK, Kellams AL, Locasale-Crouch J, Gurka MJ. Breastfeeding and
16 cognition: The role of confounders. *Breastfeeding Medicine*. Conference: 15th
17 Annual International Meeting of the Academy of Breastfeeding Medicine -
18 Breastfeeding: A Bridge to the Gold Standard San Francisco, CA United
19 States. Conference Start: 20101027 Conference End: 20101030.Conf
20 (TRUNCATED) 2010;5:326.
21
22
23 51. Hart S, Boylan LM, Carroll S, Musick YA, Lampe RM. Brief Report: Breast-
24 fed One-Week-Olds Demonstrate Superior Neurobehavioral Organization. *J*
25 *Pediatr Psychol* 2003;28:529-533.
26
27
28
29 52. Holme A, MacArthur C, Lancashire R. The effects of breastfeeding on
30 cognitive and neurological development of children at 9 years. *Child Care*
31 *Health Dev* 2010;36:583-590.
32
33
34 53. Horwood LJ, Fergusson DM. Breastfeeding and later cognitive and academic
35 outcomes. *Pediatrics*1998;101:E9.
36
37
38 54. Innis SM, Nelson CM, Lwanga D, Rioux FM, Waslen P. Feeding formula
39 without arachidonic acid and docosahexaenoic acid has no effect on
40 preferential looking acuity or recognition memory in healthy full- term infants
41 at 9 mo of age. *Am J Clin Nutr* 1996;64:40-46.
42
43
44 55. Jacobson SW, Chiodo LM, Jacobson JL. Breastfeeding effects on intelligence
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 quotient in 4- and 11-year-old children. *Pediatrics* 1999;103:E71.
- 4
5 56. James PF. Breastfeeding and intelligence. *Lancet* 1992;339:614.
- 6
7 57. Jedrychowski W, Perera F, Jankowski J, Butscher M, Mroz E, Flak E, Kaim I,
8
9 Lisowska-Miszczuk I, Skarupa A, Sowa A. Effect of exclusive breastfeeding
10
11 on the development of children's cognitive function in the Krakow
12
13 prospective birth cohort study. *Eur J Pediatr* 2012;171:151–158.
- 14
15 58. Jiang M, Foster EM, Gibson-Davis CM. Breastfeeding and the Child
16
17 Cognitive Outcomes: A Propensity Score Matching Approach *Matern Child*
18
19 *Health J* 2011;15:1296–1307.
- 20
21 59. Johnson DL, Swank PR, Howie VM, Baldwin CD, Owen M. Breast feeding
22
23 and children's intelligence. *Psychol Rep* 1996;79:1179-1185.
- 24
25 60. Keim SA, Daniels J, Siega-Riz AM, Dole N. Long-Chain Polyunsaturated
26
27 Fatty Acids from Breastmilk and Formula and Infant Cognitive Development.
28
29 *Am J Epidemiol* 2009;169:S11.
- 30
31 61. Lawlor DA, Najman JM, Batty GD, O'Callaghan MJ, Williams GM, Bor W.
32
33 Early life predictors of childhood intelligence: Findings from the Mater-
34
35 University study of pregnancy and its outcomes. *Paediatr Perinat Epidemiol*
36
37 2006;20:148-162.
- 38
39 62. Lucas A, Stafford M, Morley R, Abbott R, Stephenson T, MacFadyen U,
40
41 EliasJones A, Clements H. Efficacy and safety of long-chain polyunsaturated
42
43 fatty acid supplementation of infant-formula milk: a randomised trial. *Lancet*
44
45 1999; 354:1948-1954.
- 46
47 63. Maimaitiming T, Wang YP. Feeding and intellectual development in infants
48
49 and children from Uygur, Kazak, and Han nationality: A comparison of 454
50
51 cases. *Journal of Clinical Rehabilitative Tissue Engineering Research*
52
53
54
55
56
57
58
59
60

- 1
2
3 2007;11:10710-10712.
4
5 64. Makrides M, Neumann MA, Simmer K, Gibson RA . A critical appraisal of
6
7 the role of dietary long-chain polyunsaturated fatty acids on neural indices of
8
9 term infants: A randomized, controlled trial. *Pediatrics* 2000;105:32-38.
10
11 65. Malloy MH, Berendes H. Does breast-feeding influence intelligence quotients
12
13 at 9 and 10 years of age?. *Early Hum Dev* 1998;50:209-217.
14
15 66. Martin NW, Benyamin B, Hansell NK, Montgomery GW, Martin NG, Wright
16
17 MJ, Bates TC. Cognitive Function in Adolescence: Testing for Interactions
18
19 Between Breast-Feeding and FADS2 Polymorphisms. *J Am Acad Child*
20
21 *Adolesc Psychiatry* 2011;50(1):55–62.
22
23 67. McCrory C, Layte R. The effect of breastfeeding on children's educational test
24
25 scores at nine years of age: Results of an Irish cohort study. *Social Science*
26
27 *and Medicine* 2011;72:1515-1521.
28
29 68. Morales E, Bustamante M, Gonzalez JR, Guxens M, Torrent M, et al. (2011)
30
31 Genetic Variants of the FADS Gene Cluster and ELOVL Gene Family,
32
33 Colostrums LC-PUFA Levels, Breastfeeding, and Child Cognition. *PLoS*
34
35 *ONE* 6(2): e17181.
36
37 69. Morris SS, Grantham-McGregor SM, Lira PIC, Assuncao AMS, Ashworth A.
38
39 Effect of breastfeeding and morbidity on the development of low birthweight
40
41 term babies in Brazil. *Acta Paediatr* 1999;88:1101-1106.
42
43 70. Morrow-Tlucak M, Haude RH, Ernhart CB. Breastfeeding and cognitive
44
45 development in the first 2 years of life. *Social Science and Medicine*
46
47 1988;26:635-639.
48
49 71. Mortensen EL, Michaelsen KF, Sanders SA, Reinisch JM. The association
50
51 between duration of breastfeeding and adult intelligence. *J Am Med Assoc*
52
53
54
55
56
57
58
59
60

- 1
2
3 2002;287:2365-2371.
4
5 72. Mukerji M, Sharma K. Mental development as a function of maternal
6 economic status, literacy, occupation and feeding pattern. *Psycho-Lingua*
7 1993;23:77-81.
8
9 73. Nassar MF, Younis NT, El-Arab SE, Fawzi FA. Neuro-developmental
10 outcome and brain-derived neurotrophic factor level in relation to feeding
11 practice in early infancy. *Maternal and Child Nutrition* 2011;7:188-197.
12
13 74. NELSON CM, INNIS SM, WALSEN P, WHITFIELD M. Prospective
14 Measures of Visual and Cognitive-Development in Term Gestation Breast-Fed
15 and Formula-Fed Infants to 18 Months of Age. *Pediatr Res* 1995;37:A315.
16
17 75. Niemela A, Jarvenpaa AL. Is breastfeeding beneficial and maternal smoking
18 harmful to the cognitive development of children?. *Acta Paediatrica,*
19 *International Journal of Paediatrics* 1996;85:1202-1206.
20
21 76. Oddy WH, Kendall GE, Blair E, De Klerk NH, Stanley FJ, Landau LI, Silburn
22 S, Zubrick S. Breast feeding and cognitive development in childhood: A
23 prospective birth cohort study. *Paediatr Perinat Epidemiol* 2003;17:81-90.
24
25 77. Oddy WH, Li J, Whitehouse AJO, Zubrick SR, Malacova E. Breastfeeding
26 duration and academic achievement at 10 years. *Pediatrics* 2011;127:e137-
27 e145.
28
29 78. Paine BJ, Makrides M, Gibson RA. Duration of breast-feeding and Bayley's
30 Mental Developmental Index at 1 year of age. *J Paediatr Child Health*
31 1999;35:82-85.
32
33 79. Pollock JI. Long-term associations with infant feeding in a clinically
34 advantaged population of babies. *Dev Med Child Neurol* 1994;36:429-440.
35
36 80. Quinn PJ, O'Callaghan M, Williams GM, Najman JM, Andersen MJ, Bor W.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Breast Milk and Cognitive Development – Walfisch et al

- 1
2
3 The effect of breastfeeding on child development at 5 years: A cohort study. *J*
4
5 *Paediatr Child Health* 2001;37:465-469.
6
7
8 81. Rao MR, Hediger ML, Levine RJ, Naficy AB, Vik T. Effect of breastfeeding
9
10 on cognitive development of infants born small for gestational age. *Acta*
11
12 *Paediatrica, International Journal of Paediatrics* 2002;91:267-274.
13
14 82. Ribas-Fito N, Julvez J, Torrent M, Grimalt JO, Sunyer J. Beneficial effects of
15
16 breastfeeding on cognition regardless of DDT concentrations at birth. *Am J*
17
18 *Epidemiol* 2007;166:1198-1202.
19
20
21 83. Richards M, Hardy R, Wadsworth MEJ. Long-term effects of breast-feeding
22
23 in a national birth cohort: Educational attainment and midlife cognitive
24
25 function. *Public Health Nutr* 2002;5:631-635.
26
27
28 84. Richards M, Wadsworth M, Rahimi-Foroushani A, Hardy R, Kuh D, Paul A.
29
30 Infant nutrition and cognitive development in the first offspring of a national
31
32 UK birth cohort. *Dev Med Child Neurol* 1998;40:163-167.
33
34
35 85. Rodgers B. Feeding in infancy and later ability and attainment: A longitudinal
36
37 study. *Developmental Medicine & Child Neurology* 1978;20:421-426.
38
39
40 86. Rogan WJ, Gladen BC. Breast-feeding and cognitive development. *Early Hum*
41
42 *Dev* 1993;31:181-193.
43
44
45 87. Santiago Burruchaga M, Ruiz Sanz JI, Pijoan Zubizarreta JI, Benito Fernandez
46
47 J, Sanjurjo Crespo P. Intellectual development in the second year of life in
48
49 healthy breast-fed children vs. Formula-fed children. *An Esp Pediatr*
50
51 2000;52:530-536.
52
53
54 88. Silva AAM, Mehta Z, O'Callaghan FJK. Duration of breast feeding and
55
56 cognitive function: Population based cohort study. *Eur J Epidemiol*
57
58 2006;21:435-441.
59
60

- 1
2
3 89. Silva PA, Buckfield P, Spears GF. Some maternal and child developmental
4 characteristics associated with breast feeding: a report from the Dunedin
5 Multidisciplinary Child Development Study. *Aust Paediatr J* 1978;14:265-268.
6
7
8
9
10 90. Sloan S, Stewart M, Dunne L. The effect of breastfeeding and stimulation in
11 the home on cognitive development in one-year-old infants. *Child Care in*
12 *Practice* 2010;16:101-110.
13
14
15
16 91. Slykerman RF, Thompson JMD, Becroft DMO, Robinson E, Pryor JE, Clark
17 PM, Wild CJ, Mitchell EA. Breastfeeding and intelligence of preschool
18 children. *Acta Paediatrica, International Journal of Paediatrics* 2005;94:832-
19 837.
20
21
22
23
24
25 92. Steer CD, Smith GD, Emmett PM, Hibbeln JR, Golding J. FADS2
26 polymorphisms modify the effect of breastfeeding on child IQ. *PLoS ONE*
27 2010;5(7).
28
29
30
31
32 93. Taylor B, Wadsworth J. Breast feeding and child development at five years.
33 *Dev Med Child Neurol* 1984;26:73-80.
34
35
36 94. Temboury MC, Otero A, Polanco I, Arribas E. Influence of breast-feeding on
37 the infant's intellectual development. *J Pediatr Gastroenterol Nutr* 1994;18:32-
38 36.
39
40
41
42
43 95. Thorsdottir I, Gunnarsdottir I, Kvaran MA, Gretarsson SJ. Maternal body
44 mass index, duration of exclusive breastfeeding and children's developmental
45 status at the age of 6 years. *Eur J Clin Nutr* 2005;59:426-431.
46
47
48
49 96. Tozzi AE, Bisiacchi P, Tarantino V, Chiarotti F, D'Elia L, De Mei B, Romano
50 M, Gesualdo F, Salmaso S. Effect of duration of breastfeeding on
51 neuropsychological development at 10 to 12 years of age in a cohort of
52 healthy children. *Developmental Medicine & Child Neurology* 2012;54:843-
53
54
55
56
57
58
59
60

- 1
2
3 848.
4
5 97. Veena SR, Krishnaveni GV, Srinivasan K, Wills AK, Hill JC, Kurpad AV,
6
7 Muthayya S, Karat SC, Nalinakshi M, Fall CHD. Infant feeding practice and
8
9 childhood cognitive performance in South India. Arch Dis Child 2010;95:347-
10
11 354.
12
13 98. Whitehouse AJO, Robinson M, Li J, Oddy WH. Duration of breast feeding
14
15 and language ability in middle childhood. Paediatr Perinat Epidemiol
16
17 2011;25:44-52.
18
19 99. Wigg NR, Tong S, McMichael AJ, Baghurst PA, Vimpani G, Roberts R. Does
20
21 breastfeeding at six months predict cognitive development?. Aust N Z J Public
22
23 Health 1998;22:232-236.
24
25 100. Zaini MZ, Lim CT, Low WY, Harun F. Effects of nutritional status on
26
27 academic performance of Malaysian primary school children. Asia Pac J
28
29 Public Health 2005;17:81-87.
30
31 101. Zhou SJ, Baghurst P, Gibson RA, and Makrides M. Home
32
33 environment, not duration of breast-feeding, predicts intelligence quotient of
34
35 children at four years Nutrition 2007;23:236–241.
36
37 102. Jolly K, Ingram L, Khan KS, Deeks JJ, Freemantle N, MacArthur C.
38
39 Systematic review of peer support for breastfeeding continuation:
40
41 metaregression analysis of the effect of setting, intensity, and timing. BMJ
42
43 2012; Jan 25;344:d8287. doi: 10.1136/bmj.d8287.
44
45 103. Koren G. Bias against the null hypothesis in maternal-fetal
46
47 pharmacology and toxicology. Clin Pharmacol Ther. 1997 Jul;62(1):1-5.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 104. Koren G, Nickel C. Perpetuating fears: bias against the null hypothesis
4 in fetal safety of drugs as expressed in scientific citations. J Popul Ther Clin
5 Pharmacol. 2011;18(1):e28-32.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only



PRISMA 2009 Checklist

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	5
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	6
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6-7
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	6
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	6-7
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	8-9
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	8
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	9
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2 for each meta-analysis).	



PRISMA 2009 Checklist

Page 1 of 2

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	10+flow diagram
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	10
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Table
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Table
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	13
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	13
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	12-14
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	3

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(6): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Page 2 of 2

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

1

Breast Milk and Cognitive Development – Walfisch et al

Breast milk and cognitive development – the role of
confounders: resolving the nature vs. nurture conflict: A
systematic review

Formatted: Left

Formatted: Font: (Default) Times, 16 pt

Contributing Authors:

Asnat Walfisch MD¹, Corey Sermer², Alex Cressman MSc¹, and Gideon Koren MD
FACMT FRCPC¹

Author Affiliations:

¹ The Motherisk Program, Hospital for Sick Children; Department of Pediatrics and
Pharmacology, University of Toronto, Toronto, Ontario, Canada

²University of Western Ontario, London, Ontario, Canada

Corresponding author (and reprint requests):

Gideon Koren, MD, FACMT, FRCPC

Director of Motherisk Program

Hospital for Sick Children, 555 University Avenue, M5G 1X8

Toronto, Ontario, Canada

Phone: 416-813-6780

Fax: 416-813-7562

E-mail: gidiup_2000@yahoo.com

Word count:

Abstract: ~~268~~ ~~259~~-words

Manuscript: ~~2721~~ (excluding references): ~~2,221~~

Running title: Breast Milk and Cognitive Development – Walfisch et al

ABSTRACT:

Objectives: The association between breastfeeding and child cognitive development is conflicted by studies reporting positive and null effects. Consequently, debate remains whether such a relationship is confounded by factors associated with breastfeeding, specifically maternal socioeconomic class, IQ and education.

Methods: To try to resolve this debate, we conducted a systematic review of the literature investigating the association between breastfeeding and cognitive outcomes of healthy infants born at term.

Results: Eighty-four studies met our inclusion criteria with 34 rated as high quality, 26 as moderate and 24 as low quality. Critical assessment of accepted studies revealed the following associations: ~~14/21~~ (null), ~~28/25~~ (positive), ~~18/15~~ (null after adjusting for confounders), and ~~17/14~~ (positive - diminished after adjusting for confounders), and ~~16 with mixed results with positive associations in sub-group analysis~~. Directionality of effect did not correlate with study quality, however studies showing a decreased effect after multivariate analysis were of superior quality compared to other study groupings (i.e. ~~12/14~~ ~~14/17~~ high quality, ~~82%~~). Further, studies that showed null or diminished effect after multivariate analysis corrected for significantly more confounders (7.79 ± 3.4) as compared to those that found no change following adjustment (5.6 ± 4.5 ~~4.8 ± 4.1~~) ($p = 0.0401$). The majority of included studies were done during childhood (>1y, <18y, 75%) and set in high-income countries (85.5%).

Conclusion: The results of this systematic review support a conclusion that much of the reported effect of breastfeeding on child neurodevelopment may be due to confounding. Although it is unlikely that additional work will change the current synthesis, future studies in this field should attempt to rigorously control for all important confounders. Alternatively, study designs using sibling cohorts discordant

3

Breast Milk and Cognitive Development – Walfisch et al

for breastfeeding may yield more robust conclusions.

|

For peer review only

Article summary:**Article focus:**

- Although most published data supports the association between breast-feeding and IQ of the offspring, debate remains whether this is a causal relationship or an association with favorable parental characteristics.
- We conducted a systematic review of the literature investigating the association between breastfeeding and cognitive outcomes of healthy term infants.

Key messages:

- Over eighty studies addressing this issue were published with conflicting results.
- Studies, where the initial positive effect of breastfeeding on IQ disappeared or diminished after multivariate analysis, controlled for **almost twice as many significantly more** confounders than studies showing no such change.
- Much of the reported effect of breastfeeding on child cognitive abilities is due to maternal cognitive and socioeconomic effects.

Strengths and limitations of this study:

- The significant heterogeneity in study design and rigor precluded the conduct of a formal meta-analysis.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Formatted: Left, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Left

INTRODUCTION:

Breastfeeding confers a range of nutritional and immunological advantages upon infants including reduction in childhood illness,¹⁻⁵ diabetes,⁶⁻⁷ and obesity.⁸

The potential of breast milk to enhance cognitive development has been the focus of numerous studies since Hoefer and Hardy's initial observation in 1929.⁹ It is generally agreed that children who breast-feed are more intelligent, however debate remains whether this is a causal relationship or merely an association with favorable parental socioeconomic class and IQ. The beneficial effects of breastfeeding on the child's neurodevelopment are hypothesized, by some, to be mediated by long-chain polyunsaturated fatty acids (LCPUFAs)¹⁰ which are present in human milk, but not in cow's milk or most infant formulas.¹¹ However, a recent systematic review of all randomized trials where mother's diet was supplemented with PUFA's during pregnancy has failed to confirm such an effect.¹²

The pendulum of opinion has swung back and forth with different investigators showing inconsistent results depending upon study design and rigor. The Achilles heel of most of these studies, and the probable explanation for the conflicting results, is the difficulty in controlling for confounders that may affect child development. Furthermore, the ability to clarify this relationship is hindered by ethical considerations, which preclude randomized controlled trials, given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

Well-established confounders in breastfeeding research include demographic and IQ differences between mothers who breastfeed and those who chose not to.¹³ Parents who score high on a range of cognitive abilities have children with above average IQ scores.¹³ In parallel, advantage in mother's IQ more than doubles the odds of

breastfeeding.¹³ Thus, some of the published data demonstrates the disappearance of the breast-feeding effect on child's cognition after correction for maternal IQ.

In an attempt to partially overcome these sources of bias, a few randomized trials have been published, with randomization to breastfeeding promotion¹⁴⁻¹⁵ or in preterm infants.¹⁶ In the breastfeeding promotion intervention trial (PROBIT Trial) by Kramer and colleagues, IQ scores and academic performance tests were more favorable in the intervention group, however, statistically significant differences were only shown for some of the sub-scores.¹⁵

Systematic reviews examining the impact of breastfeeding on cognitive abilities have reached conflicting results.^{13, 17-19} The Meta-analysis by Jain et al¹⁸ suggests that less than 25% of studies in this area have adjusted for socio-demographic confounders.

There is a paucity of literature critically assessing the current published evidence within this field. In trying to address these challenges, the objective of the present work was to conduct a systematic review of published studies investigating the association between breast-feeding and neurodevelopmental outcome of healthy infants born at term.

METHODS:

The study was conducted based on a prospectively prepared protocol, using the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines.²⁰

Literature search:

Searches were conducted in the following databases (all from inception to ~~December~~ July 2011): MEDLINE(R) with Daily Update, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects, Health Technology Assessment, NHS Economic Evaluation Database, Embase and PsycINFO using the OvidSP interface, and on Science Citation Index Expanded (SCI-EXPANDED), Social Sciences Citation Index (SSCI), Conference Proceedings Citation Index- Science (CPCI-S), and Conference Proceedings Citation Index - Social Science & Humanities (CPCI-SSH) using the Web of Knowledge interface.

A search strategy was developed based on the MEDLINE, EMBASE and PsycINFO database subject headings and the "Used for" synonyms listed in the scope notes for the terms "breast feeding", "human milk", "breast milk", "infant formula", "artificial milk", "cognition", "intelligence" and "intelligent quotient". Reference lists of identified studies, textbooks, previously published systematic reviews, and review articles were also searched. No language restrictions were applied and studies in languages other than English were translated for incorporation into the study.

A second, complimentary, literature search was done on April 2013 for all studies published during the period of August 2011-December 2012.

Study selection:

Both prospective and retrospective studies (RCTs, non-~~randomised~~randomized

Formatted: Left

controlled clinical trials, cohort studies, longitudinal studies, and case-control studies) were included if:

1. One of the study aims was to address the question of breastfeeding and cognitive development;
2. The authors used reliable validated methods to evaluate cognitive development [e.g. Bayley scales of infant development, Wechsler Intelligence Scale for Children Revised (WISC-R)];
3. Prospective or retrospective documentation of use and duration of breastfeeding was used; and
4. The authors focused on healthy term infants and not those at increased biological risk for developmental delays (e.g. prematurity, intrauterine growth restriction).

Studies were excluded if:

1. The study group included preterm or small for gestational age babies.
2. Evaluation of the cognitive development was done using only a non-reliable or subjective tool (such as school grades, or maternal report).
3. Dietary patterns and breastfeeding were not evaluated since birth.

Titles and abstracts were reviewed for possible exclusion by two reviewers (A.W. and C.S.). If both reviewers excluded a citation, it was eliminated from further review. If at least one reviewer included the citation or if there was insufficient information to make a determination from the title and abstract, the full article was obtained for review. Full text articles were reviewed by three authors (A.W., C.S. and A.C.) for suitability for inclusion. Disagreements regarding study eligibility were resolved by consensus.

Study quality grading

Quality assessment of individual studies was performed by two authors (A.W and C.S.) using the three category summary grading system (A, B, C) suggested by Ip et al.¹⁹ Their system defines a generic grading system that is applicable to each type of study design including randomized controlled trials, cohort, and case-control studies as follows:

A (good)

A study that adheres mostly to the commonly held concepts of high quality including the following: clear description of the population, setting, interventions and comparison groups; clear description of the comparison groups; appropriate measurement of outcomes; appropriate statistical and analytic methods and reporting; no reporting errors; less than 20 percent dropout; clear reporting of dropouts; and appropriate consideration and adjustment for potential confounders.

B (fair/moderate)

Category B studies do not meet all the criteria in category A because they have some deficiencies, but none of which are likely to cause major biases. The study may have suboptimal adjustment for potential confounders. The study may also be missing information, making it difficult to assess limitations and potential problems.

C (poor)

Category C studies either did not consider potential confounders or did not adjust for them appropriately. These studies may have serious shortcomings in design, analysis or reporting; have large amounts of missing information, or discrepancies in reporting.

Data extraction:

Extracted data were compiled in an evidence table. The table includes a description of the studies that addressed the key question according to the inclusion/exclusion

criteria. The table provides information about study design, target population and sample size, description of breast-feeding exposure and method of categorization, nature of the comparison group, cognitive development assessment tool used and participants' age, summary of the results prior to adjustment for confounders, a list of all confounders adjusted for, differences in IQ between the groups after adjustment for possible confounders (if available), and study quality grading according to the scale described above.¹⁹

Statistical analysis:

Comparison of studies based on their results or quality was performed by Chi Square or ANOVA as appropriate.

RESULTS:

The flow of the literature search is displayed in Figure 1. Of the 1,696 potentially relevant citations identified, 84 studies met the *a priori* inclusion criteria for this systematic review (Table 1).^{9,13,15, 21-101} Overall agreement between reviewers on the inclusion of studies was 100%.

Out of these 84 publications, 34 were rated as high quality (grade A), 26 as moderate (grade B) and 24 as low quality (grade C). Overall, based on the primary endpoint of cognitive function,

Overall, there were ~~214~~ studies showing no association between IQ and breastfeeding.^{24-25,30-32,35-36,39,54,56,62-63,73-74,86-87,89,91,97,100-101} ~~292~~⁸⁵

positive studies.^{9,15,21,27-29,37-38,42,48,51,57,59,61,64,68,71-72,75,79-82,85,90,94-95,98} ~~171~~⁸⁵ initially

positive studies that became negative after accounting for confounders.^{13,23,26,33,43-}

~~46,49-50,52,55,65-66,69,78,84,96~~ and ~~17~~⁴ studies where the initial positive effect was

diminished but remained statistically significant after accounting for confounders

(Table 2).²² The remaining 16 studies showed mixed results (e.g. association only observed in the female sub-group or

only in carriers of a specific allele^{34,40-41,47,53,58,60,67,70,76-77,83,88,92-93,99}).

In general, the directionality of the results did not correlate with the studies' quality.

However, the studies showing decrease in the effect after multivariate analysis were

of superior quality compared to the rest (i.e. ~~14~~¹² of ~~14~~¹⁷ had a quality score of A =

Table 2).

Different studies corrected in their analyses for different potential confounders,

ranging from zero to 16 total confounders (Table 1). Confounders commonly

considered in these studies were socioeconomic status, maternal education, birth

weight, gestational age, birth order and gender. Some considered quality and quantity

of stimulation of the child to be crucial confounders but did not consider maternal or

paternal intelligence and other important factors. Studies that showed null or

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

diminished effect in their multivariate analysis controlled for significantly more confounders (7.79 ± 3.4) as compared to those that found no change following adjustment for confounders ($5.64.8 \pm 4.51$) ($p = 0.0401$). Furthermore, many of the studies did not have a clear definition of breastfeeding or cumulative breast milk exposure. ~~A large variety of cognitive assessment tools were used and study outcomes were measured anywhere from 8 days of age into adulthood.~~

Table 3 divides the included studies according to their settings: developed vs. developing world. The majority of included studies were set in the developed world (71/84, 85.5% vs. 13/84, 15.5% in the developing world). The quality of the studies set in developing countries were generally poorer given our criteria: 46% graded A + B (6/13) in studies set in developing countries, compared with 76% (54/71) in studies set in developed countries. Developing country studies were also more likely to reach a null association or null association after adjustment for confounding compared with developed country studies (8/13, 61% vs. 31/84, 37% respectively).

A large variety of cognitive assessment tools were used and study outcomes were measured anywhere from 8 days of age into adulthood. Table 4 divides the included studies according to age groups of participants: infancy, childhood and adulthood, with the corresponding direction of results and study quality. The majority of included studies measured intelligence during the childhood period (age 1-18 years: 70/93 studies, 75%). Studies performed during childhood and reaching an initial positive association, weakened after adjustment, were generally of higher quality than other studies (12/14 quality grade A, 86%, Table 4). Studies performed during infancy or adulthood were more likely to find a null association (before or after adjustment) compared with studies performed during childhood (Infancy group - 61%, adulthood - 60%, childhood 43%, Table

Formatted: Font: Times

Formatted: Font: Times

Formatted: Font: (Default) Times

Formatted: Font: Times

Formatted: Font: (Default) Times

Formatted: Font: Times

Formatted: Font: Times

Formatted: Font: Times

Formatted: Font: Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

Formatted: Font: (Default) Times

1
2
3
4
5
6 4).

Formatted: Font: (Default) Cambria

7
8 The significant heterogeneity in study design and rigor precluded the conduct of a
9 formal meta-analysis.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION:

The continuing debate of whether breastfeeding imparts direct advantage on child cognition, or whether this is merely an association with favorable familial socioeconomic status and cognition, is not purely theoretical. From a public health perspective, if breastfeeding has biological effects on a child's IQ - this will be one of very few cost-effective means to significantly improve a child's neurodevelopment. If, on the other hand, there is no such effect, in the case where breastfeeding is either impossible or not sought by the mother, this will allow these women to rest assured that their choice will not have long-term developmental consequences.

In the case of other comparable therapeutic dilemmas, ~~these conflicts~~ is-are typically resolved through randomized controlled trials (RCT), which are not ethically feasible in this case given that breastfeeding has other protective effects and the highly personal nature of the decision to breastfeed.

The closest comparison to a formal RCT in reducing selection bias would be sibling-pair analysis, when cognition of breastfed infants is compared to that of their siblings who were formula fed. This design ensures similar socioeconomic and maternal characteristics. ~~Importantly~~ Unfortunately, the few studies that have followed this design ~~were negative or positive~~ reached conflicting results.^{13, 40}

The second closest design to RCT was employed in the PROBIT study by Kramer and colleagues,¹⁴⁻¹⁵ who cluster randomized women in Eastern Europe to receive or not receive formal education about the advantages of breastfeeding. This study did show favorable effects, but it has been argued that the mothers randomized for the breastfeeding promotion arm might have been influenced not only in providing higher rates of breastfeeding, but also by improving other positive health behaviors.

Our analysis reveals that there are over eighty studies addressing this issue and that their results divide almost evenly between positive and negative associations. The quality of “positive” or “negative” studies did not differ, except for higher quality on average in studies that showed an apparent decrease in effect after multivariate analysis.

We have shown that studies, where the initial positive effect of breastfeeding on IQ disappeared or substantially diminished after multivariate analysis, controlled for ~~almost twice as many~~ significantly more confounders than studies showing no such change. When compared to a meta-analysis conducted 14 years ago,¹⁷ it appears that many more new studies did attempt to control for confounding measures of socioeconomic status and parental education, amongst others.

Given that more tight control of confounders resulted in greater likelihood of disappearance of breastfeeding effect, it can be argued that the remaining positive effect reflects residual uncontrolled bias, as shown by Der and colleagues in their large study.¹³ In that study, before adjustment, breastfeeding was associated with an increase of around 4 points in mental ability. *Post hoc* analysis revealed that adjustment for maternal intelligence accounted for most of this effect – where full adjustment for a range of relevant confounders yielded a small (0.52) and non-significant effect size (95% confidence interval -0.19 to 1.23).

In our systematic review, a similar effect was recorded by a total of ~~1845~~ studies, and in addition ~~14-17~~ studies showed substantially diminished effect after adjustment.

~~When we examined studies based on setting (Table 3), we found that the majority of the 84 included studies were set in the developed world (85%). Studies completed in mid-low income countries were nearly twice as likely to find a null association (before or after adjustment) compared with studies set in developed countries (61%~~

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Line spacing: Double

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

vs. 43.5% respectively).

This may be due to the fact that in many low- and middle-income countries -high rates

of some degree of breastfeeding there exist^{s102} and that comparisons between

breastfed and non-breastfed populations may examine more homogenous study

groups (i.e. parental socioeconomic status, income, and parental IQ).³⁴ In contrast,

studies originating in the developed world exhibit a greater heterogeneity between

breastfed and non-breastfed populations exists¹³ as the choice to breast feed is

associated with a family's socioeconomic status, maternal education, a number of

other measures of maternal intelligence, and social advantage (Der et al, 2006; BMJ

Article: Ref 17-19 in our paper).^{13,17-19} If a biological effect truly exists between

breastfeeding and infant IQ, one would expect this relationship to exist in multiple

settings, including the developing world. The fact that this relationship is less

apparent in developing countries suggests that much of the observed relationship may

be due to parental social advantage, confounding the choice to -breast-feed.

This systematic review includes studies using a large variety of cognitive

assessment tools and age span. The majority of included studies measured

intelligence during the childhood period (age 1-18 years, 75%). Studies

performed during infancy or adulthood were more likely to find a null

association (before or after adjustment) although number of included studies is

small. Possible explanations for this finding include reduced accuracy of IQ

evaluation in infancy (<1y) on one hand, and a variety of additional factors

influencing IQ at an older age (>18y), on the other hand.

Another factor that needs to be seriously considered in our review is the existence of bias against the null hypothesis. The likelihood of studies not

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, Font color: Auto

Formatted: Font: (Default) Times, 12 pt

Formatted: Left

Formatted: Superscript

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Not Highlight

Formatted: Superscript

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Superscript

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Superscript

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Font: (Default) Times, 12 pt

Formatted: Normal, Line spacing: Double, No bullets or numbering, No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font: Cambria

Formatted: Left

detecting a significant effect in pregnancy to be submitted and published in the peer review literature is substantially lower than that of positive studies.¹⁰²⁻¹⁰³⁻

Formatted: Font: Cambria

¹⁰⁴. This can create a distorted balance that may seriously affect the conclusions on effects of interventions.

In conclusion, this systematic review suggests that much of the reported effect of breastfeeding on child cognitive abilities is due to maternal cognitive and socioeconomic effects. When considered together with the fact that a recent systematic review failed to corroborate a biological effect of milk PUFA on brain development, it is likely that breastfeeding does not, by itself, directly affect child IQ.

Although it is unlikely that additional studies will change substantially the current synthesis, future studies in this field should attempt to rigorously control for all important confounders even if they are difficult to obtain (e.g. parental IQ). Alternatively, study designs using sibling cohorts discordant for breastfeeding may yield more robust conclusions to further clarify this dilemma.

Formatted: Font: (Default) Cambria, Font color: Auto

Author's contribution:

Formatted: Font: Cambria, No underline

Asnat Walfisch has participated in all phases of this study including literature search, data collection and review, quality grading, interpretation of the results, and has written a part of the manuscript and the revised manuscript.

Formatted: Font: Cambria

Corey Sermer has participated in all phases of this study including literature search, data collection and review, quality grading, and revision of the manuscript.

Alex Cressman has participated in the literature search, data collection and review, translation of foreign language manuscripts, and has taken a significant part in the manuscript preparation and revision.

Gideon Koren participated in all phases of this study. He initiated the study and supervised actively throughout its conduct. Specifically he was involved in the data interpretation, statistical analysis, and has written and revised a substantial part of the manuscript.

Formatted: Line spacing: single
Formatted: Font: Bold, Underline

For peer review only

References

1. Ryan AS, Wenjun Z, Acosta A. Breastfeeding continues to increase into the new millennium. *Pediatrics*. 2002 Dec;110:1103-9.
2. Dell S, To T. Breastfeeding and asthma in young children: Findings from a population based study. *Archives of Pediatric and Adolescent Medicine* 2001;155:1261-1265.
3. Bhandari N, Bahl R, Mazumdar S, Martines J, Black RE, Bhan MK. Effect of community-based promotion of exclusive breastfeeding on diarrhoeal illness and growth: A cluster randomised trial. *Lancet* 2003;361:1418-1423.
4. Galton Bachrach VR, Schwarz E, Bachrach LR. Breastfeeding and the risk of hospitalisation for respiratory disease in infancy. A meta-analysis. *Archives of Pediatric and Adolescent Medicine* 2003;157:237-243.
5. Howie PW, Forsyth JS, Ogston SA, Clark A, Florey CD. Protective effect of breastfeeding against infection. *BMJ* 1990;300:11-16.
6. Sadauskaite-Kuehne V, Ludvigsson J, Padaiga Z, Jasinskeine E, Samuelsson U. Longer breastfeeding is an independent protective factor against development of type 1 diabetes mellitus in childhood. *Diabetes Metabolism Research Reviews* 2004;20:150-157.
7. Young TK, Martens PJ, Taback SP, Sellers EAC, Dean HJ, Cheang M, Flett B. Type 2 diabetes mellitus in children prenatal and early infancy risk factors among native Canadians. *Archives of Pediatrics and Adolescent Medicine* 2002;156:651-655.

Formatted: Font: Cambria

Formatted: Font: Cambria, No underline

Formatted: Left

Formatted: Left, Line spacing: 1.5 lines

Formatted: Left, Line spacing: single

Formatted: Left

8. Owen CG, Martin RM, Whincup PH, Davey Smith G, Cook DG. Effect of infant feeding on the risk of obesity across the life course: A quantitative review of the published evidence. *Pediatrics* 2005;115:1367-1377.
9. Hoefler C, Hardy MC. Later development of breast fed and artificially fed infants. *J Am Med Assoc.* 1929;92:615–20.
10. Koletzko B, Lien E, Agostoni C, Bo`hles H, Campoy C, Cetin I, Decsi T, Dudenhausen JW, Dupont C, Forsyth S, Hoesli I, Holzgreve W, Lapillonne A, Putet G, Secher NJ, Symonds M, Szajewska H, Willatts P, Uauy R, World Association of Perinatal Medicine Dietary Guidelines Working Group. The roles of long chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *J Perinat Med* 2008;36:5–14.
11. Sauerwald TU, Demmelmair H, Koletzko B. Polyunsaturated fatty acid supply with human milk. *Lipids.* 2001;36:991-6.
12. Lo A, Sienna J, Mamak E, Djokanovic N, Westall C, Koren G. The effects of maternal supplementation of polyunsaturated Fatty acids on visual, neurobehavioural, and developmental outcomes of the child: a systematic review of the randomized trials. *Obstet Gynecol Int* 2012;2012:591531.
13. Der G, Batty GD, Deary IJ. Effect of breast feeding on intelligence in children: prospective study, sibling pairs analysis, and meta-analysis. *BMJ* 2006;333(7575):945.
14. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzikovich I, Shapiro S, et al for the PROBIT Study Group. Promotion of Breastfeeding Intervention Trial (PROBIT): A randomised controlled trial in the Republic of Belarus. *JAMA* 2001;285:413-420.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
15. Kramer MS, Aboud F, Mironova E, Vanilovich I, Platt RW, Matush L, Igumnov S, Fombonne E, Bogdanovich N, Ducruet T, Collet JP, Chalmers B, Hodnett E, Davidovsky S, Skugarevsky O, Trofimovich O, Kozlova L, Shapiro S; Promotion of Breastfeeding Intervention Trial (PROBIT) Study Group. Breastfeeding and child cognitive development: new evidence from a large randomized trial. *Arch Gen Psychiatry* 2008;65(5):578-584.
 16. Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and subsequent intelligence quotient in children born preterm. *Lancet* 1992;339:261-4.
 17. Anderson JW, Johnstone BM, Remley DT. Breastfeeding and cognitive development: a meta-analysis. *Am J Clin Nutr* 1999;70:525-35.
 18. Jain A, Concato J, Leventhal JM. How good is the evidence linking breastfeeding and intelligence? *Pediatrics* 2002;109:1044-53.
 19. Ip S, Chung M, Raman G, Chew P, Magula N, DeVine D, Trikalinos T, Lau J. Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Rep Technol Assess* 2007;153:1-186.
 20. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009; 339:b2700.
 21. Amanda M, Singh RP. A study of intelligence of children in relation to infant-feeding practices and nutrition. *Indian Journal of Psychometry & Education* 1992;23:23-28.
 22. Andres A, Cleves MA, Bellando JB, Pivik RT, Casey PH, Badger TM. Developmental status of 1-year-old infants fed breast milk, cow's milk formula, or soy formula. *Pediatrics*. 2012;129:1134-40.

23. Angelsen NK, Vik T, Jacobsen G, Bakketeig LS. Breast feeding and cognitive development at age 1 and 5 years. *Arch Dis Child* 2001;85:183–188.
24. Auestad N, Halter R, Hall RT, Blatter M, Bogle ML, Burks W, Erickson JR, Fitzgerald KM, Dobson V, Innis SM, Singer LT, Montalto MB, Jacobs JR, Qiu W, Bornstein MH. Growth and development in term infants fed long-chain polyunsaturated fatty acids: a double-masked, randomized, parallel, prospective, multivariate study. *Pediatrics*. 2001;108(2):372-81.
25. Auestad N, Scott DT, Janowsky JS, Jacobsen C, Carroll RE, Montalto MB, Halter R, Qiu W, Jacobs JR, Connor WE, Connor SL, Taylor JA, Neuringer M, Fitzgerald KM, Hall RT. Visual, cognitive, and language assessments at 39 months: a follow-up study of children fed formulas containing long-chain polyunsaturated fatty acids to 1 year of age. *Pediatrics*. 2003;112(3 Pt 1):e177-83.
26. Bartels M, van Beijsterveldt CEM, Boomsma I. Breastfeeding, Maternal Education and Cognitive Function: A Prospective Study in Twins. *Behav Genet* 2009;39:616–622.
27. Bauer G, Ewald ES, Hoffman J, Dubanoski R. Breast feeding and cognitive development in three-year-old children. *Arch Dis Child* 1991;64:763–4.
28. Birch EE, Garfield S, Castañeda Y, Hughbanks-Wheaton D, Uauy R, Hoffman D. Visual acuity and cognitive outcomes at 4 years of age in a double-blind, randomized trial of long-chain polyunsaturated fatty acid-supplemented infant formula. *Early Human Development* 2007;83:279–284.
29. Bon N. Relationship between breastfeeding and subsequent IQ of child. *Bulletin de Psychologie* 1980;33:935-947.
30. Bouwstra H, Dijck-Brouwer DAJ, Boehm G, Boersma ER, Muskiet FAJ,

Breast Milk and Cognitive Development – Walfisch et al

Hadders-Algra M. Long-chain polyunsaturated fatty acids and neurological developmental outcome at 18 months in healthy term infants. *Acta Pædiatrica*, 2005; 94: 26–32.

31. Burruchaga MS, Sanz JIR, Zubizarreta JIP, Fernandez JB, Crespo PS.

Intellectual development in the second year of life in healthy breast-fed children vs. formula-fed children. *An Esp Pediatr* 2000; 52: 530-536.

32. Caspi A, Williams B, Kim-Cohen J, Craig IW, Milne BJ, Poulton R,

Schalkwyk LC, Taylor A, Werts H, Moffitt TE. Moderation of breastfeeding effects on the IQ by genetic variation in fatty acid metabolism.

Proc.Natl.Acad.Sci.U.S.A. 2007;104:18860-18865.

33. Clark KM, Castillo M, Calatroni A, Walter T, Cayazzo M, Pino P, Lozoff B.

Breastfeeding and Mental and Motor Development at 5 1/2 Years. *Ambul Pediatr.* 2006 ; 6(2): 65–71.

34. Daniels MC, Adair LS. Breast-feeding influences cognitive development in

Filipino children. *J Nutr* 2005;135:2589–2595.

35. de Andraca,I.; Pino,P.; de la Parra,A.; Rivera,F.; Castillo,M. Risk factors for

psychomotor development among infants born under optimal biological conditions. *Rev Saude Publica* 1998;32:138-147.

36. De Andraca,I.; Salas,MI; Lopez,C.; Cayazzo,MS; Icaza,G. Effect of breast

feeding an psychosocial variables upon psychomotor development of 12 months old infants. *Arch Latinoam Nutr* 1999;49:223-231.

37. Di Girolamo D, Braibanti P, Mamone P. Feeding methods and cognitive

development in the first 3 years of life. *Rassegna di Psicologia* 1993;10:33-49.

38. Eickmann SH, de Lira PI, Lima Mde C, Coutinho SB, Teixeira Mde L,

Ashworth A. Breast feeding and mental and motor development at 12 months

Breast Milk and Cognitive Development – Walfisch et al

in a low-income population in northeast Brazil. *Paediatr Perinat Epidemiol* 2007;21:129-137.

39. Elwood PC, Pickering J, Gallacher JEJ, Hughes J, Davies D. Long term effect of breast feeding: Cognitive function in the Caerphilly cohort. *J.Epidemiol.Community Health* 2005;59:130-133.

40. Evenhouse E, Reilly S. Improved estimates of the benefits of breastfeeding using sibling comparisons to reduce selection bias. *Health Serv Res* 2005;40:1781-1802.

41. Fergusson DM, Beautrais AL, Silva PA. Breast-feeding and cognitive development in the first seven years of life. *Social Science and Medicine* 1982;16:1705-1708.

42. Florey Du CV, Leech AM, Blackhall A. Infant feeding and mental and motor development at 18 months of age in first born singletons. *Int J Epidemiol* 1995;24:S21-S26.

43. Foroushani AR, Mohammad K, Mahmoodi M, Siassi F. Effect of breastfeeding on cognitive performance in a British birth cohort. *Eastern Mediterranean Health Journal* 2010;16:202-208.

44. Gale CR, Martyn CN. Breastfeeding, dummy use, and adult intelligence. *Lancet* 1996;347:1072-1075.

45. Gale CR, Marriott LD, Martyn CN, Limond J, Inskip HM, Godfrey KM, Law CM, Cooper C, West C, Robinson SM. Breastfeeding, the use of docosahexaenoic acid-fortified formulas in infancy and neuropsychological function in childhood. *Arch Dis Child* 2010;95:174-179.

46. Ghys A, Bakker E, Hornstra G, Van den Hout M. Red blood cell and plasma phospholipid arachidonic and docosahexaenoic acid levels at birth and

Breast Milk and Cognitive Development – Walfisch et al

- cognitive development at 4 years of age. *Early Hum Dev* 2002;69:83-90.
47. Gibson-Davis CM, Brooks-Gunn J. Breastfeeding and verbal ability of 3-year-olds in a multicity sample. *Pediatrics* 2006;118: e1444-e1445.
48. Gomez-Sanchiz M, Canete R, Rodero I, Baeza JE, Gonzalez JA. Influence of breast-feeding and parental intelligence on cognitive development in the 24-month-old child. *Clin Pediatr* 2004;43:753-761.
49. GREENE LC, LUCAS A, LIVINGSTONE MBE, HARLAND PSEG, BAKER BA. Relationship between Early Diet and Subsequent Cognitive Performance during Adolescence. *Biochem Soc Trans* 1995;23:S376-s376.
50. Gurka KK, Kellams AL, Locasale-Crouch J, Gurka MJ. Breastfeeding and cognition: The role of confounders. *Breastfeeding Medicine*. Conference: 15th Annual International Meeting of the Academy of Breastfeeding Medicine - Breastfeeding: A Bridge to the Gold Standard San Francisco, CA United States. Conference Start: 20101027 Conference End: 20101030.Conf (TRUNCATED) 2010;5:326.
51. Hart S, Boylan LM, Carroll S, Musick YA, Lampe RM. Brief Report: Breastfed One-Week-Olds Demonstrate Superior Neurobehavioral Organization. *J Pediatr Psychol* 2003;28:529-533.
52. Holme A, MacArthur C, Lancashire R. The effects of breastfeeding on cognitive and neurological development of children at 9 years. *Child Care Health Dev* 2010;36:583-590.
53. Horwood LJ, Fergusson DM. Breastfeeding and later cognitive and academic outcomes. *Pediatrics* 1998;101:E9.
54. Innis SM, Nelson CM, Lwanga D, Rioux FM, Waslen P. Feeding formula without arachidonic acid and docosahexaenoic acid has no effect on

Breast Milk and Cognitive Development – Walfisch et al

preferential looking acuity or recognition memory in healthy full-term infants at 9 mo of age. *Am J Clin Nutr* 1996;64:40-46.

55. Jacobson SW, Chiodo LM, Jacobson JL. Breastfeeding effects on intelligence quotient in 4- and 11-year-old children. *Pediatrics* 1999;103:E71.

56. James PF. Breastfeeding and intelligence. *Lancet* 1992;339:614.

57. Jedrychowski W, Perera F, Jankowski J, Butscher M, Mroz E, Flak E, Kaim I, Lisowska-Miszczuk I, Skarupa A, Sowa A. Effect of exclusive breastfeeding on the development of children's cognitive function in the Krakow prospective birth cohort study. *Eur J Pediatr* 2012;171:151-158.

58. Jiang M, Foster EM, Gibson-Davis CM. Breastfeeding and the Child Cognitive Outcomes: A Propensity Score Matching Approach *Matern Child Health J* 2011;15:1296-1307.

59. Johnson DL, Swank PR, Howie VM, Baldwin CD, Owen M. Breast feeding and children's intelligence. *Psychol Rep* 1996;79:1179-1185.

60. Keim SA, Daniels J, Siega-Riz AM, Dole N. Long-Chain Polyunsaturated Fatty Acids from Breastmilk and Formula and Infant Cognitive Development. *Am J Epidemiol* 2009;169:S11.

61. Lawlor DA, Najman JM, Batty GD, O'Callaghan MJ, Williams GM, Bor W. Early life predictors of childhood intelligence: Findings from the Mater-University study of pregnancy and its outcomes. *Paediatr Perinat Epidemiol* 2006;20:148-162.

62. Lucas A, Stafford M, Morley R, Abbott R, Stephenson T, MacFadyen U, EliasJones A, Clements H. Efficacy and safety of long-chain polyunsaturated fatty acid supplementation of infant-formula milk: a randomised trial. *Lancet* 1999; 354:1948-1954.

63. Maimaitiming T, Wang YP. Feeding and intellectual development in infants and children from Uygur, Kazak, and Han nationality: A comparison of 454 cases. *Journal of Clinical Rehabilitative Tissue Engineering Research* 2007;11:10710-10712.
64. Makrides M, Neumann MA, Simmer K, Gibson RA . A critical appraisal of the role of dietary long-chain polyunsaturated fatty acids on neural indices of term infants: A randomized, controlled trial. *Pediatrics* 2000;105:32-38.
65. Malloy MH, Berendes H. Does breast-feeding influence intelligence quotients at 9 and 10 years of age?. *Early Hum Dev* 1998;50:209-217.
66. Martin NW, Benyamin B, Hansell NK, Montgomery GW, Martin NG, Wright MJ, Bates TC. Cognitive Function in Adolescence: Testing for Interactions Between Breast-Feeding and FADS2 Polymorphisms. *J Am Acad Child Adolesc Psychiatry* 2011;50(1):55–62.
67. McCrory C, Layte R. The effect of breastfeeding on children's educational test scores at nine years of age: Results of an Irish cohort study. *Social Science and Medicine* 2011;72:1515-1521.
68. Morales E, Bustamante M, Gonzalez JR, Guxens M, Torrent M, et al. (2011) Genetic Variants of the FADS Gene Cluster and ELOVL Gene Family, Colostrums LC-PUFA Levels, Breastfeeding, and Child Cognition. *PLoS ONE* 6(2): e17181.
69. Morris SS, Grantham-McGregor SM, Lira PIC, Assuncao AMS, Ashworth A. Effect of breastfeeding and morbidity on the development of low birthweight term babies in Brazil. *Acta Paediatr* 1999;88:1101-1106.
70. Morrow-Tlucak M, Haude RH, Ernhart CB. Breastfeeding and cognitive development in the first 2 years of life. *Social Science and Medicine*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1988;26:635-639.

71. Mortensen EL, Michaelsen KF, Sanders SA, Reinisch JM. The association between duration of breastfeeding and adult intelligence. *J Am Med Assoc* 2002;287:2365-2371.

72. Mukerji M, Sharma K. Mental development as a function of maternal economic status, literacy, occupation and feeding pattern. *Psycho-Lingua* 1993;23:77-81.

73. Nassar MF, Younis NT, El-Arab SE, Fawzi FA. Neuro-developmental outcome and brain-derived neurotrophic factor level in relation to feeding practice in early infancy. *Maternal and Child Nutrition* 2011;7:188-197.

74. NELSON CM, INNIS SM, WALSEN P, WHITFIELD M. Prospective Measures of Visual and Cognitive-Development in Term Gestation Breast-Fed and Formula-Fed Infants to 18 Months of Age. *Pediatr Res* 1995;37:A315.

75. Niemela A, Jarvenpaa AL. Is breastfeeding beneficial and maternal smoking harmful to the cognitive development of children?. *Acta Paediatrica, International Journal of Paediatrics* 1996;85:1202-1206.

76. Oddy WH, Kendall GE, Blair E, De Klerk NH, Stanley FJ, Landau LI, Silburn S, Zubrick S. Breast feeding and cognitive development in childhood: A prospective birth cohort study. *Paediatr Perinat Epidemiol* 2003;17:81-90.

77. Oddy WH, Li J, Whitehouse AJO, Zubrick SR, Malacova E. Breastfeeding duration and academic achievement at 10 years. *Pediatrics* 2011;127:e137-e145.

78. Paine BJ, Makrides M, Gibson RA. Duration of breast-feeding and Bayley's Mental Developmental Index at 1 year of age. *J Paediatr Child Health* 1999;35:82-85.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
79. Pollock JI. Long-term associations with infant feeding in a clinically advantaged population of babies. *Dev Med Child Neurol* 1994;36:429-440.
80. Quinn PJ, O'Callaghan M, Williams GM, Najman JM, Andersen MJ, Bor W. The effect of breastfeeding on child development at 5 years: A cohort study. *J Paediatr Child Health* 2001;37:465-469.
81. Rao MR, Hediger ML, Levine RJ, Naficy AB, Vik T. Effect of breastfeeding on cognitive development of infants born small for gestational age. *Acta Paediatrica, International Journal of Paediatrics* 2002;91:267-274.
82. Ribas-Fito N, Julvez J, Torrent M, Grimalt JO, Sunyer J. Beneficial effects of breastfeeding on cognition regardless of DDT concentrations at birth. *Am J Epidemiol* 2007;166:1198-1202.
83. Richards M, Hardy R, Wadsworth MEJ. Long-term effects of breast-feeding in a national birth cohort: Educational attainment and midlife cognitive function. *Public Health Nutr* 2002;5:631-635.
84. Richards M, Wadsworth M, Rahimi-Foroushani A, Hardy R, Kuh D, Paul A. Infant nutrition and cognitive development in the first offspring of a national UK birth cohort. *Dev Med Child Neurol* 1998;40:163-167.
85. Rodgers B. Feeding in infancy and later ability and attainment: A longitudinal study. *Developmental Medicine & Child Neurology* 1978;20:421-426.
86. Rogan WJ, Gladen BC. Breast-feeding and cognitive development. *Early Hum Dev* 1993;31:181-193.
87. Santiago Burruchaga M, Ruiz Sanz JI, Pijoan Zubizarreta JI, Benito Fernandez J, Sanjurjo Crespo P. Intellectual development in the second year of life in healthy breast-fed children vs. Formula-fed children. *An Esp Pediatr* 2000;52:530-536.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
88. Silva AAM, Mehta Z, O'Callaghan FJK. Duration of breast feeding and cognitive function: Population based cohort study. *Eur J Epidemiol* 2006;21:435-441.
89. Silva PA, Buckfield P, Spears GF. Some maternal and child developmental characteristics associated with breast feeding: a report from the Dunedin Multidisciplinary Child Development Study. *Aust Paediatr J* 1978;14:265-268.
90. Sloan S, Stewart M, Dunne L. The effect of breastfeeding and stimulation in the home on cognitive development in one-year-old infants. *Child Care in Practice* 2010;16:101-110.
91. Slykerman RF, Thompson JMD, Becroft DMO, Robinson E, Pryor JE, Clark PM, Wild CJ, Mitchell EA. Breastfeeding and intelligence of preschool children. *Acta Paediatrica, International Journal of Paediatrics* 2005;94:832-837.
92. Steer CD, Smith GD, Emmett PM, Hibbeln JR, Golding J. FADS2 polymorphisms modify the effect of breastfeeding on child IQ. *PLoS ONE* 2010;5(7).
93. Taylor B, Wadsworth J. Breast feeding and child development at five years. *Dev Med Child Neurol* 1984;26:73-80.
94. Temboury MC, Otero A, Polanco I, Arribas E. Influence of breast-feeding on the infant's intellectual development. *J Pediatr Gastroenterol Nutr* 1994;18:32-36.
95. Thorsdottir I, Gunnarsdottir I, Kvaran MA, Gretarsson SJ. Maternal body mass index, duration of exclusive breastfeeding and children's developmental status at the age of 6 years. *Eur J Clin Nutr* 2005;59:426-431.
96. Tozzi AE, Bisiacchi P, Tarantino V, Chiarotti F, D'Elia L, De Mei B, Romano

- M, Gesualdo F, Salmaso S. Effect of duration of breastfeeding on neuropsychological development at 10 to 12 years of age in a cohort of healthy children. *Developmental Medicine & Child Neurology* 2012;54:843-848.
97. Veena SR, Krishnaveni GV, Srinivasan K, Wills AK, Hill JC, Kurpad AV, Muthayya S, Karat SC, Nalinakshi M, Fall CHD. Infant feeding practice and childhood cognitive performance in South India. *Arch Dis Child* 2010;95:347-354.
98. Whitehouse AJO, Robinson M, Li J, Oddy WH. Duration of breast feeding and language ability in middle childhood. *Paediatr Perinat Epidemiol* 2011;25:44-52.
99. Wigg NR, Tong S, McMichael AJ, Baghurst PA, Vimpani G, Roberts R. Does breastfeeding at six months predict cognitive development?. *Aust N Z J Public Health* 1998;22:232-236.
100. Zaini MZ, Lim CT, Low WY, Harun F. Effects of nutritional status on academic performance of Malaysian primary school children. *Asia Pac J Public Health* 2005;17:81-87.
101. Zhou SJ, Baghurst P, Gibson RA, and Makrides M. Home environment, not duration of breast-feeding, predicts intelligence quotient of children at four years *Nutrition* 2007;23:236–241.
- 101-102. [Jolly K, Ingram L, Khan KS, Deeks JJ, Freemantle N, MacArthur C. Systematic review of peer support for breastfeeding continuation: metaregression analysis of the effect of setting, intensity, and timing. *BMJ* 2012; Jan 25;344:d8287. doi: 10.1136/bmj.d8287.](#)

1
2
3
4
5
6 | ~~102,103.~~ Koren G. Bias against the null hypothesis in maternal-fetal

7
8 | pharmacology and toxicology. Clin Pharmacol Ther. 1997 Jul;62(1):1-5.

9
10 | ~~103,104.~~ Koren G, Nickel C. Perpetuating fears: bias against the null hypothesis


11 | in fetal safety of drugs as expressed in scientific citations. J Popul Ther Clin

12 | Pharmacol. 2011;18(1):e28-32.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Formatted: Left

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1
Flow diagram:



Author and Reference	Study Design	Target Population	Breast milk exposure	Cognitive development assessment tool and participant age	Results	Confounders adjusted for	Difference in IQ after adjustment	Study Quality Grading
Amanda et al 1992 (21)	Prospective cohort. Feeding method collected retrospectively	84 school aged children from two schools in India	Exclusive breast milk for more than 4 months vs. less than 4 months or formula fed	General mental ability test for children from 7-11 years (Srivastava and Saxena 1988-89)	There were significantly more breast fed (>4 m) children in the higher IQ category (IQ>109)	None.	Not provided	C
Andres et al 2012 (22)	Longitudinal study	391 healthy infants enrolled in the Beginnings Study in Arkansas, USA	Breast fed vs. Soy fed vs. milk-based formula fed	Assessed at ages 3, 6, 9, and 12 months using the Bayley Scales of Infant Development second edition, from which the Mental Developmental Index (MDI) and Psychomotor Development Index (PDI) were derived	BF infants scored slightly higher than formula-fed infants on the Mental Developmental Index (MDI) score at ages 6 and 12 months (P < 0.05). Confounders included in the model.	Socioeconomic status, mother's age and IQ, gestational age, gender, birth weight, head circumference, race, age, and diet history	BF infants scored 1-2 points higher than formula-fed infants on the Mental Developmental Index (MDI) score at ages 6 and 12 months (P<0.05)	A
Angelsen et al 2001(23)	Prospective cohort	345 children in Scandinavia (Norway and Sweden)	< 3 months 3-6 months > 6 months	Bayley Scales of Infant Development (BSID) at 13 months. Wechsler Preschool and Primary Scales of Intelligence (WPPSI-R), and Peabody Developmental Motor Scales (PDMS) at age 5y	Shorter duration of breast-feeding was associated with lower scores on mental developmental tests both at 13 months and at 5 years of age. Unadjusted difference of 8 points	Maternal age, education, smoking, and Raven score (IQ)	Maternal age, education, and intelligence were significant confounder. When analyzing performance IQ and verbal IQ separately, the median IQ value, was not statistically different when adjusting for maternal Raven score	B
Auestad et al 2001 (24)	Prospective randomized longitudinal study comparing different formula types and non randomized breast feeding group.	294 children from four sites in the United states	Breast feeding until age 12m, versus 3 different types of formulas (±AA, DHA)	Bayley scale for infant development at 6 and 12 months Fagan test of infant intelligence at 6 and 9 months MacArthur communicative development inventories at 9 and 14 months	No difference in any of the parameters checked between any study groups.	None.	No difference in any of the parameters checked between any of the study groups.	A

Auestad et al 2003 (25)	Follow up study of Auestad 2001: Prospective randomized longitudinal study comparing different formula types and a non randomized breast feeding group.	157 children from the original 294 children from four sites in the United states	Breast feeding until age 12m, versus 3 different types of formulas (\pm AA,DHA)	At 39 months, standard tests of IQ (Stanford Binet IQ), receptive vocabulary (Peabody Picture Vocabulary Test-Revised), and expressive vocabulary (mean length of utterance).	No difference in any of the parameters checked between any study groups.	None	No difference in any of the parameters checked between any study groups.	A
Bartels et al 2009 (26)	Retrospective cohort. Breast feeding status prospectively assessed	672 monozygotic male twin pairs 637 dizygotic male twin pairs 860 monozygotic female twin pairs 647 dizygotic female twin pairs 679 male-female twin pairs 598 female-male twin pairs in the Netherlands	<Two weeks >Two weeks	Dutch CITO-elementary test at age 12 years.	Breast-fed children of highly educated mothers score on average 7.6 point higher on a standardized test of cognitive abilities than formula-fed children of mothers with a low education.	Maternal education, income.	A significant positive effect of breastfeeding on cognitive abilities above the expected positive effect of maternal education. Exact numbers not provided.	B
Bauer et al 1991 (27)	Prospective cohort	50 children from Honolulu, Hawaii	Breastfeeding as a continuous variable over time.	The McCarthy Scales of Children's Abilities at age 3 years.	The duration of breastfeeding was significantly correlated with scores on the scales, General, cognitive, Quantitative and Memory	Socio-economic status, gender, and pesticide exposure	Remained significant, numbers not provided	C
Birch et al 2007 (28)	Single-center, double-blind, randomized clinical trial comparing different formula types and non randomized breast feeding group.	52 healthy term infants enrolled for DHA and ARA supplementation and 32 breast-fed infants served as controls in Dallas, Texas	Assigned diets were fed exclusively through 17 weeks of age. In the breast feeding group the average duration of breastfeeding was 43 \pm 9 weeks.	The Wechsler Preschool and Primary Scale of Intelligence, Revised (WPPSI-R) was used to assess intelligence at 4 years of age.	The control formula and DHA-supplemented groups had Verbal IQ scores poorer than the breast-fed group. There was no difference in performance or full-scale IQ between all groups.	None.	No adjustment done.	B

Bon 1980 (29)	Retrospective cohort	954 children from France	Exclusive bottle fed Vs. Breastfed for <15 days Vs. Breastfed between 15 days and 2 months; Vs. Breastfed more than two months.	The PM-47 Non-Verbal test, at 6-8.5 years of age.	Higher scores for girls who were breast-fed versus not breast-fed. No difference in boys.	Could not be assessed from the text.	Not performed.	C
Bouwstra et al 2005(30)	A prospective, double-blind, randomized control study comparing different formulas. Non randomized breast fed group as control.	A control formula - n=169, an LC-PUFA supplemented n=146 Breastfed group n=159 in the Netherlands	Supplementation - 2 mo. All formula-fed infants received control formula between 2 and 6 mo.	The Bayley Scales of Infant Development (BSIDII) at 18 months.	Bayley's MDI and PDI result values did not differ significantly between the three groups.	Parity, HOME score, parental Education	No difference between the groups.	B
Burruchaga et al 2000 (31)	Prospective cohort	39 children born at term and from homogeneous socioeconomic status in Spain.	Exclusive breast milk for at least 2 months, vs. exclusive bottle-fed.	The Bayley Scales of Infant Development at 22 months.	Bayley's MDI and PDI result values did not differ significantly between the groups.	Maternal education, head circumference, maternal occupation, birth order, smoking status.	No difference between the groups.	C
Caspi et al 2007 (32)	Retrospective cohort	858 children from the New Zealand (Dunedin) birth cohort 1848 children from the British (E-risk study) twin birth cohort	No breastfeeding Vs. breastfeeding	New Zealand cohort: IQ measured at ages 7, 9, 11, 13 years using Wechsler Intelligence Scale for Children-Revised. IQ scores combined for an overall score. British cohort: revised short form of Wechsler Preschool and Primary Scale of Intelligence-age 5	Difference in IQ test scores between breastfed children and those not breastfed was 5.6 and 6.3 IQ points in the Dunedin and E-risk cohorts, respectively. Benefit mediated by a specific genotype. Only in children carrying the C allele.	Genetic variation in fatty acid metabolism (rs174575) Socioeconomic status, Maternal cognitive ability, Gestational age, Birth weight	Children not carrying the C allele did not benefit from breastfeeding.	A

Clark et al 2006 (33)	Prospective cohort	784 Chilean children	< 2months 2 > 8 months > 8 months	At age 5.5 years. Wechsler Preschool and Primary Scales of Intelligence – Revised (WPPSI-R) Preschool Language Scale – 3rd Edition (PLS-3) the Broad Cognitive Abilities Standard Scale (BCA) of the Bateria Woodcock-Muñoz-Revisada (Bateria-R)	Poorer outcomes on the cognitive and language assessments were found for both the short and long extremes of breastfeeding as the sole milk source. The highest scores were observed in children who received breast milk as the sole milk source for 2–8 months.	Gender, birth weight, child's age at testing, maternal education, IQ, depression, age, father absence, paternal education, HOME score, socioeconomic status, nutritional status, iron deficiency anemia at 12 months, and iron supplementation.	Difference in IQ not significant.	B
Daniels et al 2005 (34)	Prospective cohort	1984 Filipino children	Any breastfeeding during: 0-6m 6-12m 12-18m 18-24m Over 24m	Philippines Nonverbal Intelligence Test of fluid abilities at ages 8.5 and 11.5 y	Poor education and suboptimal living conditions among BF mothers were strong negative confounders, causing inverse associations between BF and cognitive ability. Increased duration of any BF was of small significant benefit for cognitive development at both ages.	Parental education, paternal presence in home, maternal age, parity, alcohol during pregnancy, preterm status of child, mother reads, child's gender, baths (n/wk), dietary variety at 2 y, household income, non-income-producing assets, electricity in home, and environmental hygiene score.	1.6 points among normal birthweight breast-fed for 12 to 18 mo vs. less than 6 mo.	A
De Andraca et al 1998 (35)	Prospective cohort. Sub analysis of an RCT concerning iron fortified formulas	788 infants, 4-6 months of age in Chile	<30 days vs. >30 days. All children in the study were breastfed for average of 75 days.	Bayley Scale of Infant Development – MDI and PDI at 12 months of age.	Breast-feeding for more than 30 days was associated with significantly lower scores (2.5 points less in MDI and 2.3 in PDI). Probably due to low SES.	SES, Education and occupation of parents, alcohol abuse, HOME, maternal intelligence by WAIS, stressful events.	No adjustment was done for the specific association of breastfeeding and Bayley scale.	C
De Andraca et al 1999 (36)	Prospective cohort	138 mother-infant dyad in Chile	Exclusive breast feeding for 6 months or more vs weaning before 45 days of age	Bayley Scale of Infant Development at 12 months of age	No difference between the groups in MDI and PDI	None; study reports similar family characteristics in each group.	No difference.	C
Der et al 2006 (13)	Database analysis of a prospective study, sibling pairs analysis, and metaanalysis	5,475 children 332 pairs of sibling discordant for breastfeeding status 545 discordant for duration of breastfeeding in the	Breastfeeding vs. no breastfeeding. Breastfeeding history obtained mostly within a year of birth	Peabody individual achievement test (PIAT) was administered to children between 5 and 14y	Unadjusted effect of breastfeeding +4.7 compared to non-breastfeeding.	Maternal IQ, education, age, family poverty, HOME stimulation score, and birth order.	After adjustment the difference became non-significant.	A

		USA						
Di girolamo et al 1993 (37)	Prospective cohort. Breastfeeding data collected retrospectively	80 children in Spain	Full breast-feeding for at least 4 months vs. no breast feeding since 2 weeks of age.	Bayley Scales of Infant Development at 8-30 months	Breast-feeding group had a higher average MDI (of 6 points) compared to the bottle fed group, but no difference in PDI.	Parental education and age, gender, birth order.	Remained significant. Data not provided.	C
Eickmann et al 2007 (38)	Prospective cohort	191 Brazilian infants	'Predominantly breast fed' 'partially breast fed' and 'non-breast fed'	Bayley Scales of Infant Development II at 12 months of age.	Full breast-feeding at 1 month was associated with a small significant benefit in mental development compared with partial or no breast-feeding. No additional advantage in mental development was found with longer durations of full breastfeeding.	Adjusted for family income, possession of TV and fridge, flush toilet, maternal work and years of schooling, number of children under 5 years, home stimulation index, smoking during pregnancy, birth weight, infant's sex, hemoglobin, weight-for-age.	Full breast feeding at 1 month was associated with +3.0 points, P = 0.02 compared with partial or no breast feeding.	C
Elwood et al 2005 (39)	Prospective cohort. Breastfeeding data collected retrospectively	779 men from Caerphilly, South Wales, UK	Artificially fed vs. Breastfed, duration unknown vs. Breastfed <3m vs. breastfed >3m	Men aged 60-74 years. 1. The national adult reading test (NART) 2. The AH4 13 verbal and mathematical reasoning 3. The choice reaction time (CRT) for hand-brain reaction speed.	In the normal birth weight group the mean cognitive function was similar in both groups. In the men whose birth weight had been below the median, having been artificially fed was associated with significantly lower results in 2 of the 3 tests.	Age, social class, education, birth order and family size, father's social class, father's unemployment.	In the normal birthweight group, the adjusted mean cognitive function was similar in both groups.	B
Evenhouse et al 2005 (40)	Database analysis of a prospective study, sibling pairs analysis. Data from the National Longitudinal Study of Adolescent Health.	2,734 sibling pairs in the USA	No breastfeeding < 3 months 3-6 months 6-9 months 9-12 months 12-24 months >24 months	Add Health's abbreviated version of the Peabody Picture Vocabulary Test (PVT), normed for age and sex.	Persistent positive correlation between breastfeeding and cognitive ability. Unadjusted Difference in PVT 4.9 percentile points when comparing never breastfed to any breastfeeding in the full sample.	Birth weight, gender, birth order, parental investment, environmental characteristics	1.68 percentile points higher for ever breast fed to never breastfed.	A

Fergusson et al 1982 (41)	Prospective cohort	Birth cohort of children from the Dunedin Multidisciplinary Child Development Study: 1037 children assessed at age 3y 991 at age 5y 954 at age 7 y.	Child bottle- fed; Breast-fed for up to 4 months; Breast- fed for 4 months or longer.	Measures of intelligence at 3, 5 and 7 years. 3y-Peabody Picture Vocabulary Test 5y-Stanford Binet Intelligence Scale 7y-Weschler Child Intelligence Scale	Children who were breast-fed for 4 months or longer had scores which were 3.84 (average) points higher than bottle-fed infants (on scales with a standard deviation of 10).	Maternal intelligence (SRA verbal scale), maternal education, childhood experiences. Maternal training in child rearing, family socioeconomic status. Birth weight and gestational age.	A small but statistically significant benefit in the test scores of breastfed vs. bottle- fed infants (mean = 1.89).	A
Florey et al 1995 (42)	Retrospective cohort	592 first born singletons in Dundee	Breast fed on discharge from the hospital Vs. Bottle-fed.	Age 18 months. Bayley Scales of Infant Mental and Motor Development.	Higher mental development was significantly related to breast-feeding on discharge from hospital.	Partner's social class, mother's education, height, alcohol and cigarette consumption, placental weight and the child's sex, birth weight and gestational age at birth.	After adjustment, the difference in Bayley mental development index between the groups was between 3.7 and 5.7 units.	C
Foroushani et al 2010 (43)	Longitudinal cohort. Breastfeeding data collected retrospectively at age 2y.	5362 singletons born in 1946 in England, Wales and Scotland	No breastfeeding 1-3 months 4+ months	Age 8 – sentence completion, reading and vocabulary. Age 11 – Verbal, reading and vocabulary. Age 15 – Verbal, reading and vocabulary. Age 26 – reading Age 43 – visual and memory.	Children who were breastfed longer scored higher on verbal tests. Tests at older age (26, 43) were not significantly different.	Birth weight, childhood illness, home conditions, parents' age and education, child's behavioral scores, parents interest in the child's development, school type.	Mean score 1.5 points higher for ages 8,11,15. Not significant at older ages.	C
Gale et al 1996 (44)	Prospective cohort	994 men and women, born between 1920 and 1930 in Hertfordshire, UK	Exclusive breast feeding, exclusive bottle feeding, mixed feeding	65-75 years old. AH4 IQ test, taken on a computer.	Participants who had been exclusively breastfed had slightly higher IQ scores compared to the 2 other groups.	The use of a dummy in infancy, number of older siblings, father's occupational class, and mother's age at the participant's birth.	After adjustment, no association was found between adult intelligence, and method of infant feeding.	B
Gale et al 2010 (45)	Prospective cohort	241 children born to the Southampton Women's Survey, UK.	Breast feeding Fortified formula feeding Unfortified formula feeding	Age 4 – Wechsler Pre-School and Primary Scale of Intelligence (3rd ed.), sentence repetition and verbal fluency measured by the NEPSY	In unadjusted analysis children who were breastfed or fed with a fortified formula had significantly higher scores.	Maternal IQ and education, social class, on benefits, age at birth, birthweight.	After adjustment the differences in IQ between groups became non-significant.	A

Ghys et al 2002 (46)	Prospective cohort. Data on breast feeding collected retrospectively	128 full term children in the Netherlands	Breast-feeding was expressed as the number of months of breast-feeding regardless of possible additional formula feeding.	At 4 years: Dutch adaptation of the Kaufman Assessment Battery for Children (K-ABC), the Groningen Developmental Scale (GOS). And items of the motor scale of the McCarthy Scales of Children's Mental Abilities.	Duration of breast-feeding showed significant correlations with cognitive development (Pearson correlation coefficient 0.26)	Plasma and RBC DHA and AA, maternal intelligence, birth weight, duration of breast-feeding and paternal educational attainment, smoking during pregnancy.	In the regression analysis the correlation disappeared.	B
Gibson-Davis et al 2006 (47)	Longitudinal birth cohort study. Breast-feeding information collected retrospectively at 1 year.	1645 American-born mothers and their babies.	Breastfeeding for at least 1 month vs none.	At three years of age: Peabody Picture Vocabulary Test-Third Edition	In unadjusted mean comparisons, breastfed children had Peabody Picture Vocabulary Test scores that were 6.6 points higher than children who were not breastfed.	An extensive set of demographic characteristics, including mother's Peabody Picture Vocabulary Test and the Home Observation for Measurement of the Environment score. Mothers were categorized into 1 of 3 educational-status groups.	After adjusting for demographic characteristics and maternal verbal ability, the coefficient dropped to 1.72.	A
Gómez-Sanchiz et al 2004 (48)	Prospective cohort, information on feeding collected retrospectively.	238 healthy babies born at term, non-IUGR in Spain	Formula fed, Breast-fed up to 4 months, Breast-fed for more than 4 months.	Bayley Infant Development Scale at 24 months of age	Infants breast-fed for longer than 4 months scored higher on the mental development scale than those breast-fed for less time.	Socio-demographic and neonatal variables including parental IQ score.	The results of multiple linear regression analysis showed that infants breast-fed for longer than 4 months scored 4.3 points more than those breast-fed for less time.	A

Greene et al 1995 (49)	Retrospective cohort	432 subjects. 208 males, 224 females in Ireland	Breast-fed vs. non-breast fed. And Breast fed for up to 12 weeks vs. more than 12 weeks.	Age: 11-16 years. Raven's Standard progressive matrices test and subsets of the Primary Mental Ability Test namely verbal meaning, reasoning and number facility.	The breastfed children showed a highly significant advantage over the non-breastfed children for all measures of IQ assessed, ranging from a 4.3 point advantage in Raven's IQ to a 6.0 point advantage in Primary mental abilities IQ.	Birth weight, gestational age, birth rank, child's sex, social class, mother's age and mother's educational level.	Following adjustment, the beneficial effect of breastfeeding (yes vs no) was statistically non-significant. A 6 point advantage in verbal IQ and 5.4 point advantage in reasoning IQ were observed for subjects breastfed for > 12 weeks, compared with less.	C
Gurka et al 2010 abstract (50)	Prospective cohort	1,050 children, from the National Institute of Child Health and Development Study of Early Child Care and Youth Development in the USA	Never; 0-6 months; longer than 6 months	Age: 4 years old. Standardized (mean.100; SD.15) cognitive outcomes.	Significant positive associations were observed between breastfeeding and cognitive outcomes before adjusting for other factors.	Maternal age, education, observed quality of the home environment, mother's attitude regarding modernity of parenting, and maternal verbal IQ	No difference between the groups.	A
Hart et al 2003 (51)	Prospective cohort	83 healthy full term infants in Texas, USA	Exclusively breastfed vs exclusively non-breastfed	Brazelton Neonatal Behavioral Assessment Scale (BNBAS) at mean age of 8.95 days	Breast-fed infants surpassed formula-fed infants on the items of the orientation, motor, range of state, and state regulation dimensions of the BNBAS. Breast-fed infants also exhibited fewer abnormal reflexes, signs of depression, and withdrawal.	Socioeconomic status	After adjustment for SES only, the differences remained significant.	B

Hofer et al 1929 (9)	Retrospective cohort	383 children in Illinois, USA	Artificially fed, breast-fed for three months or less, from four to nine months, and from ten to twenty months.	Age – 7-13 years. Stanford Revision of the Binet-Simon intelligence test and the Pintner-Patterson performance scale (a nonverbal intelligence test), and by a group educational test, the Stanford achievements test.	Artificially fed were inferior in all standardized measurements to those breast fed from four to nine months, and, with one exception, to those breast fed three months or less. Artificially fed equaled or excelled those breast fed from ten to twenty months.	None; similar paternal IQ	Adjustment was not done although they mention paternal IQ to be similar between the groups.	C
Holme et al 2010 (52)	A secondary analysis of data from follow-up study of a RCT of an intervention to reduce smoking in pregnancy	1218 children in Birmingham, UK	Not breastfed, Breastfed up to 2 months, 2-4 months, over 4 months. Also, any breastfeeding vs none	British Ability Scales (Total IQ, Visual IQ, and verbal IQ), and Quick Neurological Screening Test (QNST) at age 9 years.	Before adjustment, breastfeeding was significantly associated with higher total, verbal and visual IQ scores, and increasing duration was significantly correlated with IQ scores. Breastfeeding was associated with a crude total IQ increase of 5.49 points, which was reduced to 1.78 points on analysis with breastfeeding as a binary variable (yes/no – still significant)	Maternal demographics (including education, race, and age), smoking history, work patterns, depression, social support, neonatal details and ill health in the child	Total IQ became non-significant after adjustment	A
Horwood et al 1998 (53)	Longitudinal study – children studied at birth, 4 months, 1 year, at annual intervals to age 16, and 18 yrs.	1064 children born in New Zealand in 1977	Not breastfed, breastfed for <4 months, breastfed for 4-7 months, breastfed for >=8 months	Revised Wechsler Intelligence Scale for Children (total IQ) – age 8 and 9 Teachers rating of school performance (reading and mathematics) – age 8 and 12 Progressive Achievement Test of Reading Comprehension – ages 10 and 12 Progressive Achievement Test of Mathematics – age 11 Tests of scholastic abilities – age 13 High School Outcomes	Increasing duration of breastfeeding was associated with consistent and statistically significant increases in cognitive abilities, and children who were breastfed for >=8 months had mean test scores that were between 0.35 and 0.59 SD units higher (more than 5 points) than children who were bottle fed.	Maternal age, maternal education, family socioeconomic status, averaged standard of living, averaged family income, maternal smoking during pregnancy, gender, birth order, birth weight	Upon adjustment associations were reduced and children who were breastfed for >=8 months had scores that were 0.11-0.3 SD units (less than 3 points) higher than children who were bottle fed.	A

Innis et al 1996 (54)	Retrospective cohort	433 full term infants born in 1994 in Vancouver	Never breastfed, breastfed less than 1 month, 1-3 months, 3-6 months, 6-8 months, more than 8 months, and mixed feeding (breast and formula milk)	Visual acuity measured using acuity card procedure with Teller Acuity Cards. Cognitive development measured using the Fagan Test of Infant Intelligence, (version 4.1) at 39 +/- 1 weeks of age	There were no differences in visual acuity or novelty preference among the infants when they were stratified by incidence or duration of breastfeeding.	None.	No difference.	B
Jacobson et al 1999 (55)	Prospective Longitudinal study	323 predominantly white, middle-class children born from 1980-81, at age 4, and 280 children at age 11, from 2 cohorts of similar demographic information in Michigan, USA *most exposed to PCB during pregnancy	Breastfed vs. not breastfed	McCarthy Scales of Children's Abilities and the Peabody Picture Vocabulary Test-Revised at the age of 4. Wechsler Intelligence Scale for Children-III, Wide Range Achievement Test-Revised, and the Woodcock Word, Passage, and Reading Comprehension test at the age of 11.	At both the ages of 4, and 11, breastfeeding was significantly related to higher IQ scores.	Social class, education, maternal IQ, parenting skills (Home observation for measurement of the environment (HOME))	The relationship was reduced to non-significance after adjustment for maternal IQ and parenting skills (HOME)	B
James et al, letter 1992 (56)	Prospective cohort	38 full-term children (taken as a sample from an extended Pembrokeshire farming family, UK)	Bottle-fed babies vs. breast-fed babies	IQ test At age 16 (no details)	No difference in IQ	No adjustment	No difference	C
Jedrychowski et al 2012 (57)	Longitudinal study	468 term babies in Krakow, Poland.	Complementary breastfeeding (including none), Vs. exclusive breastfeeding up to 3 months, 4-6 months, or longer than 6	In the first 3 years of the follow-up the Bayley Mental Scales of Infant Development-second edition (BSID-II) were used. At the age of 6 and 7, the Wechsler intelligence test for children (WISC-R) was administered.	Children on mixed breastfeeding achieved lower total IQ scores at each of IQ check-ups compared with those who were exclusively breastfed.	Maternal education, baby's gender, parity, and weight gain in pregnancy.	Children breastfed exclusively for >6 months increased by 3.8 points (95% CI, 2.11-5.45).	B

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

			months.					
Jiang et al 2011 (58)	Longitudinal study	3,271 children and their mothers from the US participating in the Child Development Supplement of the Panel Study of Income Dynamics	Yes or no ever breast feeding and, never breastfed; less than 6 months, 7–12 months, and more than 12 months.	Woodcock Johnson Psycho-Educational Battery-Revised (WJ-R) test score and Wechsler Intelligence Scale for Children-Revised (WISC-R) test score at 3 and 6 years of age.	Breastfed children had higher scores on WJ-R and WISC-R tests.	Child’s age, race and ethnicity, sex, number of siblings, whether the child was first born to the mother, whether the child was born preterm, whether the child was born SGA, HOME scale, maternal IQ, age, education, health status, insurance, marital status, working, income.	Three out of the five effects remain significant; the effect sizes are smaller, with only one effect size being larger than one-fifth of a standard deviation. Longer spells of breastfeeding are uncorrelated with increases in the measures of achievement.	A
Johnson et al 1996 (59)	Longitudinal study	204 Euro-American full-term infants were followed up to the age of 3 from Galveston (Texas) area	Breastfed vs. not breastfed AND Duration of breastfeeding	At age 3 years; Stanford-Binet Fourth Edition and Peabody Picture Vocabulary Test-Revised.	Breastfeeding added significantly to the prediction of the Composite IQ Comprehension factor, Vocabulary, Absurdities, Memory for Sentences, and Peabody Picture Vocabulary Test-Revised. Duration of breastfeeding only added to the prediction of Vocabulary scores.	Socioeconomic status, HOME scores (parenting skills), mother’s intelligence, mother’s smoking behavior, gender, and birth order of the child	Breast feeding was associated with 4.6-point higher mean in children's Intelligence.	A
Keim et al 2009-abstract (60)	Prospective cohort	347 children in the USA	Exclusively breastfed vs formula fed	Mullen Scales of Early Learning at 1 yr of age	Infants exclusively breastfed demonstrated better visual reception, fine motor and overall cognitive development at 12 months than formula fed infants (4-6 points).	Preterm birth, smoking, race/ethnicity, education	Differences were weakened after adjustment. No numbers provided in abstract.	C

Kramer et al 2008 (15)	Cluster-randomized trial.	13,889 Belarussian children born between June 1996 and December 1997	Experimental group vs control group: Experimental group was encouraged to breastfeed and had greater levels of breastfeeding at 3, 6, 9 and 12 months (exclusive breastfeeding also 7 fold higher at 3 months)	Wechsler Abbreviated Scales of Intelligence and teacher evaluations of academic performance in reading, writing, mathematics, and other subjects at 6.5 yrs of age.	The experimental group had higher means on all of the Wechsler Abbreviated Scales of Intelligence measures, with cluster-adjusted mean differences of 7.5 points for verbal IQ, 2.9 points for performance IQ, and 5.9 points for full-scale IQ. Teachers' academic ratings were significantly higher in the experimental group for both reading and writing.	Maternal (and paternal) IQ, as well as all other demographic and confounding variables should be distributed randomly between the treatment groups and should not confound the treatment effect.	Cluster-adjusted mean differences of 7.5 points for verbal IQ, 2.9 points for performance IQ, and 5.9 points for full-scale IQ.	A
Lawlor et al 2006 (61)	Prospective cohort	3794 woman who delivered a singleton baby between 1981-84 in Brisbane, Australia	Never, <4 months, >= 4 months	Peabody Picture Vocabulary Test at age of 5 Raven's standard progressive matrices (Raven's SPM) and the Wide Range Achievements Test version 3 (WRAT3) at 14 years	Univariate analysis of breastfeeding vs IQ showed a significant difference between the breastfeeding groups (higher scores associated with longer breastfeeding up to 8.6 points difference).	Gender, maternal age, maternal ethnicity, maternal education, paternal education, family income, gravidity, maternal smoking, fetal distress, duration of the first and second stage of labour, mod of delivery, apgar scores at 1 and 5 min, birthweight for sex and gestational age (z score), height for age and sex (z score), BMI for age and sex (z score)	Significance remained with mean difference in IQ of 6.8 between never and over 4 months of breast-feeding.	A
Lucas et al 1999 (62)	Prospective double-blind, randomized controlled study (different formulas, breastfed control group not randomized)	447 healthy full-term children born in the UK between 1993-1995.	Breastfed for at least 6 weeks vs formula fed.	Bayley Scales of Infant Development II at 18 months.	No differences in overall developmental scores at 9 months or 18 months or in any subscale quotient at 9 months were found.	Sex, centre, maternal age, maternal education, maternal marital status, and social class	No difference with adjustment for potential confounding factors.	A
Maimaitimi	Retrospective	442 infants and	Breast fed vs	Denver Developmental Screening	There were no differences in	none	No difference.	C

ng et al 2007 (63)	cohort	children inhabited by Uygur, Han, or Kazak nationality in West China	mixed feeding	Test at age up to 3.	scores between breastfeeding vs mixed feeding groups.			
Makrides et al 2000 (64)	Prospective randomized double blind controlled. Different formulas. Breastfed group was used as control, but not randomized.	68 formula fed infants and 46 breastfed infants born at term of appropriate weight in Australia	Formula fed vs breastfed	Infant VEP acuity at 16 and 34 weeks Bayley's Scales of Development at 1 year and 2 years.	At 1 yr of age, MDI scores of breastfed and formula-fed infants were not different, but at 2 years of age, MDI scores of breastfed infants were higher than formula fed infants. PDI scores were similar.	Home screening questionnaire scores, occupational prestige, parental education, gender, birth weight, maternal smoking, birth order, feeding mode.	MDI score was higher at age 2 in the breastfed group. 95% CI 3-16.8.	A
Malloy et al 1997 (65)	Retrospective cohort	518 children born in 1978-79 in Washington DC who were exposed to chloride deficient formulas.	No breastfeeding (176) vs any time length of breastfeeding (342)	Weschler Intelligence Scale-Revised at 9 or 10 years old	Breastfed scored significantly higher on full scale IQ; Further analyses limited to those exclusively breastfed for the first 60 days failed to demonstrate any significant relationship.	Maternal education, paternal education and annual income	Following adjustment, the difference was not significant.	B
Martin et al 2011 (66)	Retrospective cohort	1431 children (twin siblings) from the greater Brisbane area	Exclusively formula fed, any breastfeeding between birth and 3 months, exclusively breastfed for 3 to 6 months, exclusively breastfed for 6 months or more	FSIQ (full scale IQ) assessed using the Multi-dimensional Aptitude Battery (MAB) at 16 years of age	Breastfeeding was significantly associated with FSIQ scores. No effect of duration of breastfeeding on FSIQ was found.	Socioeconomic status, paternal education, maternal education, gestational age, and birth weight.	The effect was no longer significant after adjustment.	A
McCrory et al 2011 (67)	Retrospective cross-sectional study	8568 school children in Ireland born between 1997 and 1998.	Breastfed vs not breastfed AND Never breastfed, <=5 weeks, 6-15 weeks, 16-25 weeks, 26+ weeks	Age 9 years. Vocabulary component of the Drumcondra Primary Reading Test-Revised and part 1 of the Drumcondra Primary Mathematics Test-Revised.	In unadjusted analysis, children who were breastfed scored 8.67 percentage points higher on reading and 7.42 percentage points higher on math. Evidence of dose-response relationship was weak.	Gender, birth weight, gestation period, NICU, primary and secondary carer's social class, primary carer's education level, household income, mothers age at birth, Irish/Non-Irish, number of children's books in the home.	After adjustment, remained significant but weakened: 3.24 and 2.23 percentage points in reading and math respectively	A

Morales et al 2011 (68)	Retrospective cohort analyzing polymorphisms in genes encoding enzymes involved in LC-PUFA synthesis	Two population-based birth cohorts n = 400 mother-child pairs from INMA-Sabadell; and n = 340 children from INMA-Menorca in Spain	Different types of formula fed vs breastfed	Mental development was assessed at age 14 months using the Bayley Scales of Infant Development, first Edition – MDI only, and at age 4 years old by the Spanish version of the McCarthy Scales of Children’s Abilities (MCSA)	Children with variants associated with lower synthesis of LC-PUFA had higher scores when breastfed, while those with greater capacity to synthesize these fatty acids had higher scores regardless of breastfeeding practices.	Sex, child age, psychologist, quality of neuropsychological test, maternal education, breastfeeding, and use of gas stove at home.	Not being breastfed conferred an 8- to 9-point disadvantage in cognition among children with low FADS1 activity and in a 5-8 point disadvantage in cognition among children with low ELOVL5 activity.	A
Morris et al 1999 (69)	Prospective cohort	102 normal birthweight children born in Brazil at term.	Number of breastfeeds per day (recorded daily).	Bayley Scales of Infant Development at 6 months and 12 months (MDI and PDI).	Breastfeeding frequency over the first 4 weeks of life, but not later, was significantly associated with mental development (MDI) at 6 months of age.	Socioeconomic data: family income, a household resources index, a housing quality index, a water and sanitation index, and maternal and paternal literacy	An average of one extra breastfeed per day resulted in an increase of approximately one-quarter of a point. The effect was no longer apparent at 12 months of age.	B
Morrow-Tlucak et al 1988 (70)	Prospective cohort	229 children born in Ohio between 1981-1982	No breastfeeding, breastfeeding =<4 months, breastfeeding >4 months	Bayley Scales of Infant Development – Mental Development Index (MDI) at 6 months, 1 year, and 2 years of age	Significant differences in MDI scores between the three groups (longer breastfeeding = higher scores) at both 1 year and 2 years of age (no significant difference at 6 months).	Parent education (mean of both parents), maternal attitude (Authoritarian Family Ideology), maternal intelligence (PPVT-R), cigarette use, maternal age, race, marital status, Home Observation for Measurement of the Environment (HOME) at age 1, HOME at age 2, exact age at time of testing	With covariate control, a small but significant relationship between duration of breastfeeding and Bayley MDI at 1 and 2 years was detected. Infants breastfed for 4 months or more scored on	A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

							average 9 points higher compared to the bottle-fed infants.	
Mortensen et al 2002 (71)	Prospective longitudinal birth cohort	Mixed sample: 973 men and women All-male sample: 2280 men in Denmark	Divided to five groups: < 1 month 2-3 months 4-6 months 7-9 months > 9 months	Wechsler Adult Intelligence Scale (WAIS) at a mean age of 27.2 years in the mixed-sex sample. Borge Priens Prøve (BPP) test at a mean age of 18.7 years in the all-male sample.	Duration of breast-feeding was associated with significantly higher scores on the Verbal, Performance, Full Scale WAIS IQs and BPP test.	Parental social status Parental education Single mother status Mother's height, age, and weight gain during pregnancy cigarette consumption during the third trimester number of pregnancies Gestational age birth weight Birth length Indexes of pregnancy Delivery complications	The results remained significant with 4.6 points higher IQ for those breast-fed over 9 months compared to less than 1 month.	B
Mukerji et al 1993 (72)	Cross-sectional study	100 children aged 0-3 years in India	No breast-feeding or mixed. Exclusive breast-feeding for 4-9 months. Exclusive breastfeeding for more than 9 months	Developmental Screening Test (DST)	In the 4-9 months exclusive breast fed group 100% had average or above average IQ. In the two other groups this number was significantly lower.	None.	No adjustments were made.	C
Nassar et al 2011 (73)	Cross-sectional study	42 healthy infants in Cairo, Egypt	Breastfed, artificially fed, mixed feeding	Bayley scale of infant development-second edition (BSID-II) between 4-6 months of age.	No difference in MDI or PDI. There was significant difference only in total behavior rating scale (TBRS) and motor quality percentile rank.	No significant difference between groups in terms of age, sex, and socioeconomic standard (indirectly adjusted for with multiple regression)	Significant increase in mean adjusted TBRS and motor quality percentile rank.	B
Nelson et al 1995 (74)	Prospective cohort	Term gestation infants in USA	Breast fed or formula fed for 3 months	Teller Acuity Cards at 14d, 3, 4, 8, and 18 months Fagan Test of Infant Intelligence at 8, 10 and 12 months Bayley Scales of Infant Development at 4, 8, and 18 months	There were no significant differences between breast-fed and formula-fed infants in visual acuity at 14d, 3, 8, or 18 months, or recognition memory or the Bayley PDI or MDI at any age		Meeting abstract only available	C

Niemela et al 1996 (75)	Prospective follow-up	726 children born between 1985-86 in Finland	Breastfed <5 months, breastfed >5 months (matched pairwise based on maternal education and sex)	Non-verbal Columbian Mental Maturity Scale (CMM), visual integration using the Beery test and active vocabulary by naming of pictures at 56 months	Children breastfed for 5 months or more attained higher scores in developmental tests (sig difference found in relation to the general cognitive capacity and visual motor integration). No evidence of any interaction between verbal development and breastfeeding to 5 months or more.	Groups matched pairwise based on sex and maternal education.	Maternal education and parental status correlated with all cognitive scores. Sex and breastfeeding correlated with scores of the general cognitive capacity and visual motor integration tests in multiple linear regressions.	B
Oddy et al 2003 (76)	Prospective cohort	1401 children at first follow up and 1283 children at second follow up from the Western Australian Pregnancy Cohort Study following 2860 children in Perth, Australia	Never breastfed, fully breastfed 0-4 months, 4-6 months, more than 6 months	Peabody Picture Vocabulary Test-Revised (PPVT-R) for receptive English vocabulary – verbal intelligence at 6 yrs Performance subtest by the Wechsler Intelligence Scale for Children – Third Edition (WISC-III) – Block Design Test at age 8yrs	On average, children breast fed for more than 6 months had mean verbal IQ scores that were 6.44 points higher and Block Design scores that were 1.13 points higher than children never breastfed (small but significant differences).	Gestational age, maternal age, maternal education, parental smoking and older siblings (all covariates that were significantly correlated with verbal IQ and the performance subtest or breastfeeding)	breast feeding for >6 months was associated with an increase in verbal IQ of 3.56 points. The Performance subtest was weakened and was no longer a significant difference after adjustment.	A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Oddy et al 2011 (77)	Prospective cohort	980 children from the Western Australian Pregnancy Cohort Study following 2860 children in Perth, Australia	Breastfed less than 4 vs more than 4 months And Breastfed less than 6 vs more than 6 months	Western Australian Monitoring Standards in Education (WAMSE) scores in: math, reading, writing, and spelling at age 10yrs	Continuous breastfeeding was significantly associated with an increase in scores with each additional month of breastfeeding for math, reading, writing, and spelling.	Gender, maternal age, maternal education, family income, marital status, parent looks at book with child at age 5, maternal country of birth	Results were attenuated when adjusted for confounders. Significant interactions were found in math and spelling, revealing that boys were more likely to have improved academic scores if breastfed for a longer period.	A
Paine et al 1999 (78)	Retrospective cohort	96 healthy full term Caucasian children from the Adelaide area in Australia	Duration of exclusive breastfeeding	Bayley Scales of Infant Developmental at age 10-14 months	Duration of exclusive breast-feeding significantly predicted mental development scores for boys, but not for girls. Duration of breast-feeding did not predict psychomotor development scores.	Duration of breast-feeding, parents occupational prestige, parents' education level and smoking habits, number of siblings, birth order, HSQ score, gestational age, birthweight, age of testing, maternal age and gender were considered as possible independent variables – independent variables with p < 0.02 included in the model – gender, maternal age, birthweight, and duration of breastfeeding.	None.	C

<p>Pollock et al 1994 (79)</p>	<p>Prospective cohort</p>	<p>3838 children from the 1970 British Births Survey at full term and healthy birth weight</p>	<p>Wholly breast fed for more than 3 months vs. wholly bottle fed</p>	<p>Human figure drawing score, copying design score and English Picture Vocabulary Test score at age 5. Dichotomised outcomes: Pictorial Language Test, Friendly Maths test, Edinburgh Reading test, Spelling Test, British Ability Scales (Word definitions, similarities, matrices, total) and Continuous outcomes: Pictorial Language test, Friendly Maths test, Edinburgh Reading test, Spelling test, and British Ability Scales (Word definitions, similarities, matrices, total) at age 10</p>	<p>Significant difference found between groups at age 5 for the English Picture Vocabulary test. Significant difference found for both dichotomized outcomes and continuous outcomes for the British Ability Scales and dichotomized outcomes for the Pictorial Language test at age 10.</p>	<p>Age father left full time education, age mother left full time education, highest educational qualifications of mother, mother's smoking behavior during pregnancy, antenatal labour preparation class attendance, infant's place of birth</p>	<p>All results adjusted for all other independent predictors of breast-feeding: English Picture Vocabulary test: aOR 1.50. Dichotomized outcomes: Pictorial Language test aOR 1.49 Word definitions aOR 1.55 Similarities aOR 1.64 Total aOR 1.64 Continuous outcomes: Word definitions mean diff 3.5 Matrices mean diff 2.6 Similarities mean difference 3.0 Total mean difference 3.1</p>	<p>A</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1									
2									
3									
4									
5									
6									
7									
8	Quinn et al 2001 (80)	Prospective cohort	3880 healthy children from the Mater Hospital-University of Queensland Study of Pregnancy project	Never breastfed, < 3 weeks, 3-7 weeks, 7 weeks – 4 months, 4-6 months, still breast feeding at 6 months	Peabody Picture Vocabulary Test Revised (PPVT-R) at 5 yrs. of age.	Significantly increasing scores were found between duration of breastfeeding and the PPVT-R scores.	Birthweight, poverty, maternal education, maternal age, time in daycare/preschool, number of children in household at 5 years, English speaking background for mother & father, and degree of infant stimulation	After adjusting for a wide range of biological and social factors, the adjusted mean for those breastfed for 6 months or more was 8.2 points higher for females and 5.8 higher for males than compared to those never breastfed (this was significant).	A
9									
10									
11									
12									
13									
14									
15									
16									
17									
18									
19									
20									
21	Rao et al 2002 (81)	Prospective cohort	299 children born in Norway/Sweden at appropriate size for gestational age (comparison group)	<12 weeks Vs. >12 weeks of breastfeeding AND Duration of breastfeeding as a continuum	Bayley Scales of Infant Development at 13 months of age Norwegian version of the Wechsler Preschool and Primary Scales of Intelligence – Revised at 5 yrs. of age Peabody Development Motor Scale measured at 5 yrs. of age	There were statistically significant differences in IQ between the 2-breastfeeding groups.	Site of enrolment, maternal education, maternal IQ, maternal smoking, admission to a neonatal intensive care unit, kindergarten attendance, gender and asymmetric intrauterine growth retardation	3.7 points for total IQ and 4.1 points for performance IQ. Results remained unaltered when adjusted for confounding variables.	B
22									
23									
24									
25									
26									
27									
28	Ribas-Fito et al 2007 (82)	Prospective cohort	391 children born in Spain between 1997-1999	0-2 weeks breastfeeding 2-20 weeks breastfeeding 20+ weeks breastfeeding	Spanish version of the McCarthy Scales of Children's Abilities (general cognitive scale, verbal scale, perceptual-performance scale, memory scale, quantitative scale, motor scale subsets) at 4 yrs. of age.	Children with longer periods of breastfeeding performed significantly better on the McCarthy cognitive scale (except motor).	Gender, academic trimester at examination, psychologist, maternal social class, maternal education, and maternal use of alcohol and tobacco during pregnancy	After adjustment for confounders, significance remained for the general cognitive scale, and a trend remained in other subsets.	A
29									
30									
31									
32									
33									
34									
35									
36									
37									
38									
39									
40									
41									
42									
43									
44									
45									
46									
47									
48									
49									

<p>Richards et al 2002 (83)</p>	<p>Retrospective cohort</p>	<p>1741 people from the MRC National Survey of Health and Development (NSHD), also known as the British 1946 birth cohort</p>	<p>Never Up to 2 months 2-7 months 7+ months</p>	<p>National Adult Reading Test (NART), word-list learning task (verbal memory) and timed letter search task (mental speed and concentration) at age 53.</p>	<p>Educational Attainment: Odds of obtaining higher qualifications by the age of 26 were statistically significant, with greater odds for longer breastfeeding. Cognitive Function: Regression coefficients for the unadjusted effect show that breastfeeding was significantly associated with the NART and verbal memory tests.</p>	<p>Early background variables: Sex, father's social class, mother's education, birth order, parental interest in education and material home conditions.</p>	<p>Effect of BF on education was strengthened after adjusting for the early background variables and remained significant. No longer significant with further adjustment for cognitive ability at age 15. Cognitive function: Coefficients were reduced after adjusting for the early background variables and were further reduced after adjusting for educational attainment and adult social class Only NART remained sig.</p>	<p>A</p>
--	-----------------------------	---	--	---	---	--	---	-----------------

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Richards et al 1998 (84)	Retrospective cohort	511 first born offspring of the British 1946 cohort and their parent	Duration of breastfeeding.	Sentence completion test, reading test and vocabulary test at age 8 yr.	Breastfeeding was positively associated with cognitive function at age 8 in the first offspring of a national birth cohort. This association was not evident in the subsample of mothers of these offspring. Association in the offspring cohort became non-significant after adjusting for social class, maternal education, or maternal cognitive performance.	Social class, parental educational attainment, material home conditions, maternal age at birth, birth order, family size, maternal cigarette smoking, parental interest in education, attendance at nursery school, whether offspring cohort members had been taught cognitive skills at age 4, and cognitive test scores of the mothers.	Non-significant; after adjusting for social class, maternal education, or maternal cognitive performance.	C
Rodgers et al 1978 (85)	Longitudinal study	1464 children at age 8 and 1398 children at age 15 from the British 1946 cohort	Never bottle-fed vs. never breastfed	Picture intelligence test and word reading test at 8 years of age Non-verbal ability, mathematics, and sentence completion tests at age 15 years of age	A preliminary analysis indicated that low scores were more likely for those who had been bottle-fed than breast-fed. The mean sentence completion scores between the two groups is statistically significant. Every test at both ages was significant except for word reading scores after correction for background factors.	Sex, social group, parental interest in education, material home conditions, sample stratification, father's education, mother's education, family size, birth rank, age at weaning	After correction for confounders, every test at both ages was significant except for word reading scores.	B
Rogan et al	Prospective	855 newborns being	Bottle fed	Bayley Mental and Psychomotor	Bayley Mental and Psychomotor:	Age, race, occupation,	Confounders	B

<p>1993 (86)</p>	<p>cohort</p>	<p>followed in North Carolina were enrolled between 1978-1982 and followed up to 5 yrs old</p>	<p>‘short breastfeeding’ ‘medium breastfeeding’ ‘long breastfeeding’ ‘very long breastfeeding’</p>	<p>Development Index at 6, 12, 18, and 24 months of age. McCarthy General Cognitive, Verbal, Quantitative, Memory, Perceptual Performance, and Motor scales at 36, 48, and 60 months of age Report Card Grades from 3rd, 4th and 5th grade (averaged)</p>	<p>After adjusting for co-variables, the results at all four time points were similar, differences among the groups were only statistically sig. at 24 months. McCarthy: All scales showed trends towards higher scores with increasing length of breastfeeding, but the relationship was weakest for the motor scale. Differences after adjustment were only significant at 3 and 4 years (marginally at 5 years) between length of breastfeeding groups Report Cards: Showed slight increase with breastfeeding. Marginally significant after adjustment for English scores and not significant for math.</p>	<p>education, smoking, drinking, child’s sex, birthweight, number of older siblings, prenatal PCB exposure and dichlorodiphenyl dichloroethene exposure , Identity of the examiner.</p>	<p>integrated into model; unadjusted results not shown.</p>	
<p>Santiago Burruchaga et al 2000 (87)</p>	<p>Prospective cohort</p>	<p>39 children born at term and from homogenous sociocultural status in Spain</p>	<p>Breastfed for at least 2 months vs. formula fed</p>	<p>Bayleys scale at 2 years of age</p>	<p>No statistically significant differences between groups were found in cognitive function.</p>	<p>Maternal age, level of education, occupation, number of children in the family, smoking.</p>	<p>None; confounders integrated into model.</p>	<p>C</p>
<p>Silva et al 2006 (88)</p>	<p>Longitudinal population based</p>	<p>9367 children from the 1970 British</p>	<p>Never breastfed Less than 1</p>	<p>British Ability Scale, the Shortened Edinburgh Reading Test (word</p>	<p>Breast-feeding showed a positive association with cognition at 10</p>	<p>Socioeconomic class, birth weight, parity, gestational age,</p>	<p>Breastfeeding was weakly</p>	<p>A</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	cohort	Cohort Study comprising individuals born during April 5-11 in the UK	month 1-3 months More than 3 months	recognition), the Friendly Math Tests, and the Pictorial Language Comprehension Test at age of 10 yrs.	years before adjustment.	maternal age and maternal smoking	associated with cognitive function after adjustment (standardized coefficient 0.07). However, this effect was much smaller in the structural equation model adjusting for the same variables and did not reach significance level, suggesting that it is of little clinical importance.	
Silva et al 1978 (89)	Retrospective cohort	1037 children from the Dunedin Multidisciplinary Child Development Study	< 1 week 1-4 weeks 5-12 weeks 13-24 weeks 25-36 weeks 37-51 weeks 51+ weeks	Gross motor co-ordination, fine motor co-ordination, verbal comprehension and verbal expression, intelligence, child behavior problems at age 3 yrs.	Comparison of the groups resulted in only one significant difference among 96 comparisons made. No significant differences in age of attainment of milestones, gross or fine motor ability scores, verbal comprehension or expression, ability, intelligence, the incidence of separation problems, hyperactivity, height, weight and head circumference.	Socioeconomic class, general mental ability (IQ), level of education. Pairwise comparison with group matching to account for confounders.	None; pairwise matching said to account for confounders.	B
Sloan et al 2010 (90)	Cross-sectional observational. Breastfeeding data were obtained retrospectively.	137 infants and mothers in Ireland	Breast fed (defined as more than 1 month) vs. not breast fed (defined as less than a month a not at all)	Bayley Scales of Infant Development (second edition; Bayley, 1993), between 10 and 18 months old.	Mean cognitive scores were significantly higher in breast-fed infants (110) compared with formula-fed infants (105). Breastfeeding duration had a positive linear association with cognitive scores at one year.	Home environment (HOME score) and infant gender	The adjusted standardized beta for breastfeeding was 0.285, which remained significant after adjustment.	C
Slykerman 2005 (91)	Cross-sectional observational.	550 infants, 50% of which were SGA at	Not at all, less than 6 m,	Stanford Binet Intelligence Scale, 4th Edition	Breastfeeding was not significantly related to	Examiner administering the intelligence test, gestation,	No difference in IQ.	B

	Breastfeeding data were obtained retrospectively.	birth and 50% AGA in New Zealand.	6 to 12 m, >12 m.	at age 3.5y-4y	intelligence scores in the AGA group.	gender, maternal education, marital status, parental occupation, maternal age, parity, and smoking.		
Steer et al 2010 (92)	Prospective observational. Breastfeeding data were obtained prospectively.	9656 children from the Avon Longitudinal Study of Parents and Children cohort, in United Kingdom	Breastfeeding within the first month of life vs. never breast feeding	Wechsler Intelligence Scale for Children, 8 years old.	Breastfeeding showed a strong association with full-scale IQ with breastfed children scoring 8 points higher IQ on average in unadjusted analyses.	Maternal education, paternal social class, low birth weight, pre-term gestation, home environment, parenting and gender	The breastfeeding effect attenuated to a 3- point advantage after adjustment.	B
Taylor et al 1984 (93)	Longitudinal population based cohort. Breastfeeding data were obtained retrospectively.	13,135 children from The Child Health and Education Study in the United Kingdom	Never breastfed Less than 1 month 1-3 months More than 3 months	English picture vocabulary test (EPVT) adapted from the American Peabody Picture Vocabulary Test, at 5 years of age.	Children breast-fed for three or more months scoring over one-quarter of a standard deviation above the norm.	The age of the child at testing; the child's sex and birthweight; whether there were older or younger siblings in the home when the study child was five years old; home furnishings and equipment; maternal age at the child's birth; maternal smoking and the social index	Breast feeding remained a significant influence on EPVT scores, but the difference between the groups was small: +0.12 in the standardized EPVT score	A
Temboury et al 1994 (94)	Prospective cohort	229 infants in Spain	Breast fed – at least three months Bottle-fed – none or less than 1 month	Baylay's scale at age 18-29 months.	Bottle fed infants had lower IMD scores (index of mental development)	Maternal age, education and social class, job, psychosocial risk, number of children, infants shyness, tantrum, hyperactivity, gender, birthweight, height, place of birth	The result remained significant.	C
Thorsdottir et al 2005 (95)	Longitudinal cohort	85 children in Iceland	Duration of exclusive breastfeeding	The Icelandic developmental inventory at age 6 years. Information collected from the mothers.	Duration of exclusive breastfeeding, in months, was positively related to children's total developmental index.	Maternal and paternal education, and family income, birthweight, maternal BMI.	Total developmental index (B=1.07±0.5, P=0.04)	C
Tozzi et al 2012 (96)	Prospective cohort	1403 children originally included in	Duration of exclusive	An estimated IQ was obtained from scores of the vocabulary,	Scores of only a few neuropsychological tests were	Sex, birthweight, gestational age, mother's age at birth, type	In the regression analysis, the	A

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

		an Italian clinical trial on acellular Pertussis vaccines	breastfeeding both as a continuous variable and as a categorical variable (<6mo, >6mo).	similarities, block design, and coding tests at 10-12 years of age. An estimation of IQ was obtained from the scores of the four WISC-R subtests.	affected by exclusive breastfeeding duration: Mean scores on vocabulary, similarities, the Boston naming test, and estimated IQ improved with the duration of breastfeeding, whereas performance in one of the subtests for writing praxis decreased with breastfeeding duration.	of delivery, family composition, parents' education, presence of chronic diseases, current prescription of antihistamines or antiepileptic drugs, and the amount of thimerosal to which children were exposed through vaccines.	score on one subcategory of the California verbal learning test was negatively associated with breastfeeding for longer than 6 months. No difference was detected in any of the other test scores included in the analysis	
Veena et al 2010 (97)	Longitudinal cohort	514 children from the Mysore Parthenon birth cohort in south India	Six categories from <3 to ≥18 months	Kaufman Assessment Battery at 9- to 10-year-old children	Within this cohort, in which prolonged breastfeeding was the norm (90% breastfed ≥6 months), there were no associations between longer duration of breastfeeding and cognitive function.	Age, sex, gestation, birth size, maternal age, parity, socio-economic status, parents' attained schooling and rural/urban residence.	No difference either unadjusted or after adjustment.	B
Whitehouse et al 2011 (98)	Longitudinal cohort, breast feeding data	1195 live born children recruited at approximately 18	(1) Never breast-fed, (2) Breast-fed	Peabody Picture Vocabulary Test – Revised (PPVT-R) at age 10 years. Raw scores are transformed to	Strong positive association between the duration of predominant breast-feeding and	Maternal age at conception, maternal education, family income, and the presence of the	Children who were predominantly	A

	collected prospectively	weeks gestation, Western Australian Pregnancy Cohort (Raine) Study.	predominantly for <4 months, (3) Breast-fed predominantly for 4–6 months (4) Breast-fed predominantly for >6 months.	standard scores, based around a mean of 100 and a SD of 15.	PPVT-R at age 10 years.	biological father in the family home, maternal smoking and alcohol consumption during pregnancy, maternal experience of stressful events during pregnancy, parity, gestational age, child's sex, and proportion of optimal birthweight, a measure of the appropriateness of fetal growth, Family functioning (MMFAD), Parenting Scale, language-learning environment.	breast-fed for >6 months had a mean PPVT-R score that was 4.04 points higher than children who were never breast-fed	
Wigg et al 1998 (99)	Longitudinal cohort, breast feeding data collected prospectively	375 children born in Port Pirie, South Australia.	At 6 month of age: Breast fed vs. bottle-fed. Children who were breast-fed for less than 6 months would have been classified in our analysis as bottle-fed.	Bayley Scales of Infant Development at age 2, the McCarthy Scales of Children's Abilities age 4 and Wechsler Intelligence Scale for Children at age 7,11,13.	Estimated unadjusted advantages for the breast-fed children at ages 2, 4 and 7 years were 5.5 points, 4.6 points and 4.3 points in MDI, GCI and IQ scores, respectively. At age 11 to 13 years, the unadjusted advantage for the breast-fed children was 3.8 points in IQ.	Gender, maternal age at delivery, socioeconomic status, HOME score, maternal IQ, parental smoking habits, birthweight, birth rank, lifetime average blood lead concentration and whether parents were living together.	Adjusting for the covariates diminished the association of the feeding method in infancy with cognitive development. The covariates contributing most to this attenuating effect were the HOME scores, maternal IQ, socioeconomic status and parental smoking habits.	B
Zaini et al 2005 (100)	Retrospective cohort	1397 children from Selangor Malaysia	<6 months >6 months	Raven's Colored Progressive Matrices at a mean age 9.6 years	Those who were breastfed <u>less</u> than six months performed better (31.01 vs. 30.63 out of max 36)	None.	None.	C

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Zhou et al 2007 (101)	Prospective cohort	302 children born between 1998 and 1999 in Adelaide, Australia	Not breast fed, Breast fed at hospital discharge, Less than 6 months, More than 6 months.	Stanford-Binet Intelligence Scale at 4 years of age.	There was no association between duration of breast-feeding and childhood IQ in this relatively well-nourished cohort from an industrialized society. Before adjustment, children who were breast-fed for at least 6 m. had higher IQ than those who were breast-fed for less than 6 months	Birth order and sex of the child, maternal smoking in pregnancy, parental education, parental occupation, and quality of home environment	There was no association between breast-feeding and childhood IQ.	B
------------------------------	--------------------	--	---	--	---	---	---	----------

Table 2 – Studies by directionality and quality

Direction of findings	Number of studies	Quality grading (%)
No association between IQ and BF	21*	A - 4 (19) B - 9 (43) C - 8 (38)
Positive association between IQ and BF	28**	A - 10 (36) B - 7 (25) C - 11 (39)
Initial positive association which became negative after adjustment for confounders	18***	A - 6 (33.3) B - 8 (44.4) C - 4 (22.3)
Initial positive association, weakened after adjustment for confounders but remained statistically significant	17****	A - 14 (82) B - 2 (12) C - 1 (6)

IQ - Intelligence quotient, BF -breastfeeding

*References: 24-25,30-32,35-36,39,54,56,62-63,73-74,86-87,89,91,97,100-101

** References: 9,15,21,27-29,37-38,42,48,51,57,59,61,64,68,71-72,75,79-82,85,90,94-95,98

*** References: 13,23,26,33,43-46,49-50,52,55,65-66,69,78,84,96

**** References: 22,34,40-41,47,53,58,60,67,70,76-77,83,88,92-93,99

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 3 – Studies by setting, directionality and quality grading

Setting	Direction of findings	Number of studies (%)	Quality grading (%)
Developing Countries* (n = 13)	No association between IQ and BF	6 (46)	A – 0 (0) B – 2 (33.3) C – 4 (66.6)
	Positive association between IQ and BF	4 (30)	A – 1 (25) B – 0 (0) C – 3 (75)
	Initial positive association which became negative after adjustment for confounders	2 (15)	A – 0 (0) B – 2 (100) C – 0 (0)
	Initial positive association, weakened after adjustment for confounders but remained statistically significant	1 (8)	A – 1 (100) B – 0 (0) C – 0 (0)
Developed Countries** (n = 71)	No association between IQ and BF	15 (21)	A – 4 (26.6) B – 7 (46.6) C – 4 (26.6)
	Positive association between IQ and BF	24 (34)	A – 9 (37.5) B – 7 (29.2) C – 8 (33.3)
	Initial positive association which became negative after adjustment for confounders	16 (22.5)	A – 6 (37.5) B – 6 (37.5) C – 4 (25)
	Initial positive association, weakened after adjustment for confounders but remained statistically significant	16 (22.5)	A – 13 (81.25) B – 2 (12.5) C – 1 (6.25)

IQ – Intelligence Quotient; BF – breastfeeding

Developing: *15, 21, 33, 34, 35, 36, 38, 63, 69, 72, 73, 97, 100

Developed: **9, 13, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 70, 71, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 98, 99, 101

Table 4 – Studies by age group, directionality and quality grading

Age Group	Direction of findings	Number of studies (%)	Quality grading (%)
Infancy * ≤1 year of age n=18	No association between IQ and BF	9 (50)	A – 3 (33.3) B – 3 (33.3) C – 3 (33.3)
	Positive association between IQ and BF	5 (28)	A – 0 (0) B – 1 (20) C – 4 (80)
	Initial positive association which became negative after adjustment for confounders	2 (11)	A – 0 (0) B – 1 (50) C – 1 (50)
	Initial positive association, weakened after adjustment for confounders but remained statistically significant	2 (11)	A – 1 (50) B – 0 (0) C – 1 (50)
Childhood ** 1-18 years of age n=70	No association between IQ and BF	14 (20)	A – 3 (21) B – 5 (36) C – 6 (43)
	Positive association between IQ and BF	26 (37)	A – 10 (38.5) B – 6 (23) C – 10 (38.5)
	Initial positive association which became negative after adjustment for confounders	16 (23)	A – 6 (37.5) B – 6 (37.5) C – 4 (25)
	Initial positive association, weakened after adjustment for confounders but remained statistically significant	14 (20)	A – 12 (86) B – 2 (14) C – 0 (0)
Adulthood *** ≥18 years of age n=5	No association between IQ and BF	2 (40)	A – 0 (0) B – 1 (50) C – 1 (50)
	Positive association between IQ and BF	1 (20)	A – 0 (0) B – 1 (100)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

			C - 0 (0)
	Initial positive association which became negative after adjustment for confounders	1 (20)	A - 0 (0) B - 1 (100) C - 0 (0)
	Initial positive association, weakened after adjustment for confounders but remained statistically significant	1 (20)	A - 1 (100) B - 0 (0) C - 0 (0)

IQ – Intelligence Quotient; BF – breastfeeding

Infancy*: 22,24,35,36,37^,38,51,54,60,64^,69,70^,72^,73,74^,78^,86^,90^

Childhood**: 9,13,15,21,23,25,26,27,28,29,30,31,32,33,34,37^,40,41,42,43^,45,46,47,48,49,50,52,53,55,56,57,58,59,61,62,63,64^,65,66,67,68,70^,72^,74^,75,76,77,78^,79,80,81,82,84,85,86^,87,88,89,90^,91,92,93,94,95,96,97,98,99,100,101

Adulthood***: 39,43^,44,71,83

^Study examined two different age groups and therefore included in several categories.