

Online Resource 1- Prevalence of Cardiovascular Disease Risk Factors for Long-term Cancer Survivors by Race/Ethnicity and Cancer Type

	Race/Ethnicity						χ^2 , p-value
	Total		Non-Hispanic, White (n=580)	Hispanic, White (n=207)	African-American (n= 368)	Asian/Pacific Islander (n=348)	
	N	Wt% (95% CI)	Wt% (95% CI)	Wt% (95% CI)	Wt% (95% CI)	Wt% (95% CI)	
BMI							
<25	599	38.0 (34.3, 41.8)	40.8 (35.1, 46.8)	26.8 (19.7, 35.5)	23.5 (16.6, 32.1)	60.7 (53.5, 67.5)	9.99, p<.0001
25-<30	520	38.0 (34.3, 41.9)	36.9 (31.3, 42.8)	44.0 (35.2, 53.2)	44.9 (36.1, 54.0)	25.7 (20.4, 31.8)	
30+	384	24.0 (20.8, 27.5)	22.3 (17.8, 27.7)	29.2 (21.4, 38.4)	31.6 (24.4, 39.8)	13.6 (9.1, 19.9)	
Physical Inactivity							
Meets Guideline	606	44.0 (40.3, 47.8)	43.7 (38.1, 49.6)	47.4 (38.4, 56.5)	45.8 (37.8, 54.1)	38.5 (32.2, 45.2)	2.62, p=.02
Some Activity	589	37.9 (34.2, 41.6)	42.0 (36.4, 47.9)	31.4 (23.6, 40.6)	30.0 (23.1, 38.0)	40.6 (33.5, 48.1)	
No Activity	324	18.1 (15.6, 20.9)	14.2 (10.7, 18.7)	21.2 (14.7, 29.5)	24.2 (18.9, 30.3)	20.9 (15.4, 27.8)	
Smoking Status							
Never smoked	811	54.7 (50.8, 58.5)	55.6 (49.8, 61.3)	54.8 (45.5, 63.8)	40.8 (32.1, 50.0)	69.1 (62.5, 75.1)	4.75, p<.0001
Former smoker	573	40.2 (36.5, 44.0)	40.0 (34.5, 45.7)	41.8 (33.1, 51.1)	49.8 (40.6, 59.0)	26.8 (21.1, 33.3)	
Current smoker	91	5.1 (3.8, 6.9)	4.4 (2.8, 6.8)	3.4 (1.3, 8.4)	9.5 (5.3, 16.3)	4.1 (1.9, 8.5)	
Hypertension							
No	637	45.0 (41.1, 48.9)	52.1 (46.2, 57.9)	43.6 (34.6, 53.1)	30.4 (22.7, 39.4)	39.5 (33.2, 46.2)	6.14, P<0.0004
Yes	877	55.0 (51.1, 58.9)	47.9 (42.1, 53.8)	56.4 (46.9, 65.4)	69.6 (60.6, 77.3)	60.5 (53.8, 66.8)	
Diabetes							
No	1161	79.3 (76.0, 82.2)	86.0 (81.4, 89.6)	70.4 (60.9, 78.4)	72.9 (64.4, 80.0)	73.1 (66.4, 78.9)	6.80, p=0.0001
Yes	364	20.7 (17.8, 24.0)	14.0 (10.4, 18.6)	29.6 (21.6, 39.1)	27.1 (20.0, 35.6)	26.9 (21.1, 33.6)	

Physical activity was classified as meeting American College of Sports Medicine/ American Heart Association’s physical activity guidelines for adults[18] (\geq 150 minutes of moderate-intensity or 60 minutes of vigorous-intensity physical activity per week), some moderate or vigorous physical activity, but below guideline level, or those reporting no moderate and no vigorous activity in the past 4 weeks. Self-report of ever diagnosis with hypertension or diabetes defined these risk factors. The number of observations listed for each race/ethnicity group is for BMI; the number of observations for other risk factors was very similar.

	Cancer Type				X², p-value
	Breast (n= 376) Wt% (95% CI)	Prostate (n= 382) Wt% (95% CI)	Colorectal (n= 382) Wt% (95% CI)	Gynecologic (n= 391) Wt% (95% CI)	
BMI					
<25	42.7 (35.8, 49.9)	30.4 (24.1, 37.6)	42.6 (35.7, 49.8)	40.9 (33.1, 49.2)	6.15, p<.001
25-<30	32.8 (26.5, 39.9)	51.1 (43.7, 58.5)	33.3 (26.8, 40.5)	22.4 (17.1, 28.6)	
30+	24.5 (18.9, 31.1)	18.5 (13.4, 24.9)	24.1 (18.0, 31.4)	36.7 (28.9, 45.2)	
Physical Inactivity					
Meets Guidelines	37.1 (30.5, 44.3)	54.2 (47.1, 61.1)	41.9 (34.7, 49.4)	31.3 (25.0, 38.4)	4.92, p<.001
Some Activity	44.1 (37.1, 51.3)	32.4 (26.2, 39.4)	41.7 (34.3, 49.5)	37.4 (30.2, 45.2)	
No Activity	18.8 (14.0, 24.8)	13.4 (9.6, 18.4)	16.5 (12.2, 21.8)	31.3 (24.4, 39.2)	
Smoking Status					
Never smoked	65.8 (58.6, 72.3)	49.2 (41.7, 56.6)	41.9 (34.9, 49.3)	67.0 (59.9, 73.4)	7.66, p<.001
Former smoker	27.2 (21.3, 34.1)	46.6 (39.2, 54.2)	54.2 (46.9, 61.3)	27.1 (21.0, 34.2)	
Current smoker	7.0 (4.0, 11.9)	4.3 (2.3, 7.9)	3.9 (2.3, 6.3)	5.9 (3.2, 10.5)	
Hypertension					
No	54.8 (48.0, 61.4)	43.2 (35.9, 50.7)	38.1 (31.2, 45.5)	46.6 (38.7, 54.6)	3.96, p=0.008
Yes	45.2 (38.6, 52.0)	56.8 (49.3, 64.1)	61.9 (54.5, 68.8)	53.4 (45.4, 61.3)	
Diabetes					
No	81.1 (75.1, 86.0)	77.2 (70.7, 82.6)	79.7 (73.7, 84.7)	79.4 (72.5, 85.0)	0.32, p=0.81
Yes	18.9 (14.0, 24.9)	22.8 (17.4, 29.3)	20.3 (15.3, 26.3)	20.6 (15.0, 27.5)	

Physical activity was classified as meeting American College of Sports Medicine/ American Heart Association's physical activity guidelines for adults[18] (\geq 150 minutes of moderate-intensity or 60 minutes of vigorous-intensity physical activity per week), some moderate or vigorous physical activity, but below guideline level, or those reporting no moderate and no vigorous activity in the past 4 weeks. Self-report of ever diagnosis with hypertension or diabetes defined these risk factors. The number of observations listed for each cancer type is for BMI; the number of observations for other risk factors was very similar.