

ONLINE SUPPLEMENT

HO-1 Induction Improves Type-1 Cardiorenal Syndrome in Mice with Impaired

Ang II-induced Lymphocyte Activation (SCID Mice)

Sumit R. Monu^{1,4}, Paola Pesce^{1,2}, Komal Sodhi^{1,4}, Massimo Boldrin^{1,3},

Nitin Puri⁴, Larisa Fedorova⁴, David Sacerdoti², Stephen J. Peterson⁵,

Nader G. Abraham¹ and Attallah Kappas⁶

¹Department of Medicine, Joan C. Edwards School of Medicine, Marshall University, Huntington,
WV, 25755, USA

²Department of Medicine, ³CORIT-Consortium for Research in Organ Transplantation,
University of Padova, 35100 Padova, Italy

⁴Department of Physiology & Pharmacology,

University of Toledo College of Medicine, Toledo, OH, 43614, USA

⁵Department of Medicine, New York Medical College, Valhalla, NY, 10595, USA

⁶The Rockefeller University, New York, NY, 10065, USA

Running Title: T-lymphocyte and HO-1 in cardiorenal syndrome

Corresponding author: Nader G. Abraham, PhD., Dr. H.C., FAHA, Professor,

Marshall University Joan Edwards School of Medicine, Huntington, West Virginia, 25701-3655,

FAX: 304 691-1726, Tel 304-691-1791, E-mail: abrahamn@marshall.edu

RESULTS

Plasma Renin and Angiotensin II levels:

Figure S




Figure S1. Effect of CoPP in C57 and SCID mice on plasma Renin and Angiotensin II levels 30 days post MI. **(A, B)** Plasma Renin level (ng/ml) * $p < 0.05$ vs. Control, # $p < 0.05$ vs. MI, + $p < 0.05$ vs. MI+CoPP; **(C, D)** Plasma Angiotensin II level (pg/ml) of C57 and SCID mice respectively; * $p < 0.05$ vs. Control, # $p < 0.05$ vs. MI, + $p < 0.05$ vs. MI+CoPP.