

Appendix 1 (as supplied by the authors): Descriptive variables for patients included in the 2-week follow-up survey

	Study group; no. (%) children		Statistical test used for comparison and p-value
	LET n = 72 (50.35)	Placebo n = 71 (49.65)	
Median (IQR) age in years	4.33 (2.75, 8.08)	3.33 (2.17, 5.17)	Wilcoxon Rank Sum test p = 0.04
Sex, female	26 (37.68)	31 (44.29)	Chi-square test for association between sex and allocation p = 0.4
Location of laceration - Face - Torso - Extremities - Other	65 (94.20) - 4 (5.80) -	65 (92.86) - 5 (7.14) -	Fisher's exact test p = 1
Median (IQR) length of laceration, cm	10.50 (8.00, 15.00)	12.00 (10.00, 17.00)	Wilcoxon Rank Sum test p = 0.1
Median (IQR) age of laceration, hours [time of injury to time at triage]	1.22 (0.67, 1.88)	1.08 (0.82, 1.72)	Wilcoxon Rank Sum test p = 0.6
Median (IQR) amount of time LET applied, hours [time LET applied to time at completion of glue application]	1.08 (0.95, 1.38)	1.05 (0.92, 1.33)	Wilcoxon Rank Sum test p = 0.5