

Appendix 2 (as supplied by the authors): Descriptive variables for per-protocol placebo group versus those surveyed

	Study group; no. (%) children		Statistical test used for comparison and p-value
	Placebo group n = 98 (100)	In survey n = 70 (71.43)	
Median (IQR) age in years	3.42 (2.17, 5.25)	3.33 (2.17, 5.17)	Wilcoxon Rank Sum test p = 0.8
Sex, female	42 (42.86)	31 (44.29)	Chi-square test for association between sex and allocation p = 0.9
Location of laceration			Fisher's exact test p = 0.9
- Face	92 (93.88)	65 (92.86)	
- Torso	-	-	
- Extremities	5 (5.10)	5 (7.14)	
- Other	1 (1.02)	-	
Median (IQR) length of laceration, cm	12.00 (10.00, 15.00)	12.00 (10.00, 17.00)	Wilcoxon Rank Sum test p = 0.7
Median (IQR) age of laceration, hours [time of injury to time at triage]	1.12 (0.82, 1.73)	1.08 (0.82, 1.72)	Wilcoxon Rank Sum test p = 0.7
Median (IQR) amount of time LET applied, hours [time LET applied to time at completion of glue application]	1.07 (0.92, 1.33)	1.05 (0.92, 1.33)	Wilcoxon Rank Sum test p = 0.9