

Infant cortex responds to other humans from shortly after birth

Teresa Farroni^{1,3}, Antonio Chiarelli², Sarah Lloyd-Fox³, Stefano Massaccesi⁴, Arcangelo Merla²,
Valentina Di Gangi¹, Tania Mattarello¹, Dino Faraguna^{5,6}, Mark H. Johnson³.

¹*Dipartimento di Psicologia dello Sviluppo e della Socializzazione, Università di Padova, Padova, Italy,*

²*Infrared Imaging Lab. ITAB -Institute of Advanced Biomedical Technologies and Dept. of Neuroscience and Imaging, University of Chieti-Pescara, Italy,*

³*Centre for Brain and Cognitive Development, Birkbeck, University of London, United Kingdom,*

⁴*Dipartimento di Psicologia, Università di Padova, Padova, Italy,*

⁵*Dipartimento di Pediatria, Ospedale di Monfalcone, Monfalcone, Italy*

⁶*IRCCS Burlo Garofolo, Childrens Hospital Trieste, Italy*

Figure S1




Figure S2




Figure S3





Figure S4



Supplementary Figure Legends

Figure S1: Average time course changes in HbO₂ and HHb concentration (μM), $\pm\text{SEM}$ (Standard Error of the Mean), for each source–detector couple, in response to H-F stimulus.

Figure S2: Average time course changes in HbO₂ and HHb concentration (μM) for each source–detector couple, in response to M stimulus in Experiment 1.

Figure S3: Average time course changes in HbO₂ and HHb concentration (μM), $\pm\text{SEM}$ (Standard Error of the Mean), for each source–detector couple, in response to H-A stimulus.

Figure S4: Average time course changes in HbO₂ and HHb concentration (μM) for each source–detector couple, in response to M stimulus in Experiment 2.