

Supporting Information

Oligodeoxynucleotides containing multiple thiophene-modified isomorphous fluorescent nucleosides

Mary S. Noé, Renatus W. Sinkeldam, and Yitzhak Tor*

University of California, San Diego, Department of Chemistry and Biochemistry, La Jolla, California 92093 (USA).

*to whom correspondence should be addressed: ytor@ucsd.edu

Contents:

- **S.1** Photophysical studies.....S2
 - **Figure S1.1** Stern-Volmer plot of **1** in water.....S2
 - **Figure S1.2** Temperature dependent fluorescence spectra for ds **4•6** in triplicateS2
 - **Figure S1.3** Temperature dependent fluorescence spectra for ss **4** in triplicateS2
 - **Figure S1.4** Temperature dependent fluorescence spectra for ds **5•7** in triplicateS2
 - **Figure S1.5** Temperature dependent fluorescence spectra for ss **5** in triplicateS3
- **S.3** NMR spectra of compound **2**.....S3

S1 Spectra and graphs




Figure S1.1. Stern-Volmer plot of the fluorescence intensity of **1** in the presence of increasing concentrations of each native nucleotide (AMP, dTMP, GMP, and CMP) in water.




Figure S1.2. Temperature dependent fluorescence spectra for ds **4•6** in triplicate, and a plot of the change in fluorescence intensity at 433 nm as a function of temperature.




Figure S1.3. Temperature dependent fluorescence spectra for ss **4** in triplicate, and a plot of the change in fluorescence intensity at 433 nm as a function of temperature.




Figure S1.4. Temperature dependent fluorescence spectra for ds **5•7** in triplicate, and a plot of the change in fluorescence intensity at 433 nm as a function of temperature.




Figure S1.5. Temperature dependent fluorescence spectra for ss **5** in triplicate, and a plot of the change in fluorescence intensity at 433 nm as a function of temperature.

S.2 NMR Spectra

