



Cardiovascular health knowledge, attitude, and practice/behavior in an urbanizing community of Nepal: a population-based cross-sectional study from Jhaukhel-Duwakot Health Demographic Surveillance Site

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002976
Article Type:	Research
Date Submitted by the Author:	30-Mar-2013
Complete List of Authors:	Vaidya, Abhinav; Kathmandu Medical College, Community Medicine Aryal, Umesh Raj; Kathmandu Medical College, Krettek, Alexandra; Nordic School of Public Health NHV,, ; Institute of Medicine, Sahlgrenska Academy at University of Gothenburg, Department of Internal Medicine and Clinical Nutrition
Primary Subject Heading:	Public health
Secondary Subject Heading:	Cardiovascular medicine
Keywords:	PUBLIC HEALTH, PREVENTIVE MEDICINE, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3 **Cardiovascular health knowledge, attitude, and practice/behavior in an urbanizing**
4 **community of Nepal: a population-based cross-sectional study from Jhaukhel-Duwakot**
5 **Health Demographic Surveillance Site**
6
7
8
9
10

11 Abhinav Vaidya^{1,2}, and Alexandra Krettek^{2,3}
12
13

14
15
16 ¹Department of Community Medicine, Kathmandu Medical College, Kathmandu, Nepal

17 ²Nordic School of Public Health NHV, Gothenburg, Sweden, and

18 ³Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska
19 Academy at University of Gothenburg, Gothenburg, Sweden
20
21
22
23

24
25 **Correspondence to**

26 Abhinav Vaidya; Abhinav.Vaidya@nhv.se
27
28
29
30

31 **Keywords:** knowledge, attitude, and practice/behavior; Nepal; cardiovascular health; health
32 literacy
33
34

35
36 **Word count:** 4, 337 (excludes title page, abstract, references, tables, and figures, per author
37 guidelines)
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ARTICLE SUMMARY

Article focus

- We aimed to gain a deeper understanding of knowledge, attitude, and behavior/practice regarding cardiovascular health in a semi-urban community of Nepal, a low-income South Asian country.

Key messages

- The study site has a high burden of cardiovascular risk factors. In particular, low physical activity and overweight/obesity exceed the national average.
- Unprompted responses to cardiovascular health questions revealed a poor level of knowledge, and respondents' attitude toward cardiovascular health showed low risk perception.
- Poor correlation between knowledge, attitude, and behavior scores imply that commonly applied health promotion strategies that merely increase knowledge of cardiovascular health may be insufficient, thus emphasizing the need for a more holistic approach to disease prevention.

Strengths and limitations of the study

- This study is the first to extensively explore knowledge, attitude, and behavior/practice regarding cardiovascular health in Nepal.
- Random sampling derived from a sampling frame generated from a health demographic surveillance site provides the logistic advantage of further follow-up and intervention.
- Using both prompted and unprompted questions in the same study illustrates the variation that may be obtained by these approaches.
- Except for the WHO STEPS questionnaire on behavioral risk factors, we used questions from many unvalidated sources.
- Unintentional oversampling of women respondents yielded unsegregated gender-wise results that are more likely to represent knowledge, attitude, and practice/behavior in women.

Abstract:

Objectives: This study aimed to determine the knowledge, attitude, and practice/behavior of cardiovascular health in residents of a semi-urban community of Nepal.

Design: To increase understanding of knowledge, attitude, and practice/behavior toward cardiovascular health, we conducted in-home interviews using a questionnaire based on WHO STEPS and other resources, scoring all responses. We also recorded blood pressure and took anthropometric measurements.

Setting: Our study was conducted as part of the HARDIC project at the Jhaukhel-Duwakot Health Demographic Surveillance Site in two urbanizing villages near Kathmandu. Our results present baseline data from the project that aims to improve cardiovascular health literacy through health promotion.

Participants: The study population included 777 respondents from six randomly selected clusters in both villages.

Primary outcome measure: Proportion of the study population with correct knowledge about the causes, manifestation, immediate management, and preventability of cardiovascular diseases. Our data reveals participants' attitude toward cardiovascular health and its preventability as well as cardiovascular health behavior, as reflected in the risk factor status and positive actions taken toward achieving them.

Secondary outcome measure: Our data calculated and correlated participants' scores regarding knowledge, attitude, and practice/behavior.

Results: Participants showed only poor knowledge of heart disease causes, heart attack symptoms, and appropriate action. In general, the participants' attitude toward heart health was inappropriate. The burden of cardiovascular risk factors among participants was high. Median percent scores for knowledge, attitude, and practice/behavior were 79.25, 74.29, and 48.00, respectively. We observed poor correlations between the Pearson coefficient for knowledge and

1
2
3 attitude; attitude and behavior/practice; and knowledge and behavior/practice (0.025, 0.126, and
4
5 0.121, respectively).
6
7

8
9 Conclusions: Our study demonstrates poor level of cardiovascular health knowledge, attitude,
10 and practice/behavior in a semi-urban community in a low-income nation. Tailoring health
11 promotion activities to our findings might improve public health literacy for cardiovascular
12 disease.
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 INTRODUCTION

28
29
30 Noncommunicable diseases (NCDs) pose increasingly important public health problems in low-
31 and middle- income countries (LMICs). In 2008, 80% of global NCD deaths occurred in
32 LMICs,¹ and NCDs recently accounted for 7.9 million of 14.5 million deaths (54%) in Southeast
33 Asia.² Poverty, illiteracy, poor health infrastructure, and demographic transition, including
34 increasing life expectancy, are major contributors to rising burden of NCDs in the region. Other
35 postulated causes include poor fetal and childhood nutrition.^{3,4} A similar pattern is evident in
36 South Asian countries like Nepal, where NCDs currently occur more frequently than
37 communicable diseases and behavioral risk factors such as tobacco smoking are high.⁵
38
39
40
41
42
43
44

45 Globally, cardiovascular diseases (CVDs) comprise 39% of all NCD deaths.⁶ Eighty percent of
46 CVDs result from tobacco smoking, unhealthy diet, physical inactivity, and harmful use of
47 alcohol.⁷ The risk of all-cause mortality among individuals who score poorly in behavioral
48 factors is four fold higher than those who score best.⁸ In high-income countries, declining CVD
49 mortality is largely attributed to the reduction of behavioral risk factors.⁹ Underlying
50 socioeconomic parameters and societal influences affect exposure and vulnerability to CVD risk
51 factors. Therefore, surveillance of NCD risk factors should include social determinants of
52
53
54
55
56
57
58
59
60

1
2
3 cardiovascular health.^{1, 10} Currently, understanding the individual and social determinants of
4 cardiovascular health behaviors is among the top 20 priority areas for NCD research in LMICs.¹¹
5
6

7
8 As a precursor of favorable health behavior, experts have advocated health literacy since
9 community-based interventions (e.g., Finland's North Karelia Project) began exploring the
10 potential impact of behavioral changes on CVD.^{1, 12} The World Health Organization (WHO)
11 promotes public awareness of diet and physical activity as a "best buy" to improve
12 cardiovascular behavior.⁹ Similarly, the American Heart Association emphasizes the importance
13 of improving community awareness of cardiovascular health issues.^{13, 14} However, achieving
14 desired health behavior requires deeper understanding of the underlying social context of
15 disease.¹⁵ Hence, studies on knowledge, attitude, and practice (KAP) increase our understanding
16 of cardiovascular health literacy and community behavior.¹⁶ Currently, such studies have been
17 limited in South Asia.
18
19
20
21
22
23
24

25
26 In Pakistan, inability to recognize the symptoms of heart attack associates independently with
27 delayed hospitalization.¹⁷ An earlier study reported a low level of knowledge about heart attack¹⁸
28 and diabetes symptoms in Nepal.¹⁹ Nepal's high prevalence of underlying behavioral risk factors,
29 particularly tobacco and alcohol consumption, and poor cardiovascular health knowledge is a
30 matter of concern.²⁰
31
32
33
34
35

36
37 Currently, attempts to improve cardiovascular health knowledge and behavior in Nepal focus
38 mainly on campaigns that transmit information but pay little attention to the population's actual
39 health literacy. To gain greater insight into this important issue, we undertook a community-based
40 KAP study in the Jhaukhel-Duwakot Health-Demographic Surveillance Site (JD-HDSS) outside
41 Kathmandu.²¹ The baseline census in these villages suggested that NCDs, particularly CVDs, are
42 the predominant cause of morbidity and mortality.²¹ Additionally, our pilot study in the Duwakot
43 population demonstrated poor cardiovascular health knowledge.²²
44
45
46
47
48

49
50 Heart-Health Associated Research and Dissemination in the Community (HARDIC) is the
51 cardiovascular health component of JD-HDSS. In Nepalese, hardic means "heartily" or "from the
52 heart". Using community-level interventions, HARDIC aims to improve cardiovascular health
53 literacy and practice/behavior and this paper reports the baseline results from HARDIC. Our
54
55
56
57
58
59
60

1
2
3 study is the first to conduct a detailed analysis of KAP/health literacy in relation to
4 cardiovascular health.
5
6
7
8
9

10 **METHODS**

11 **Study site and population**

12 We conducted our descriptive, community-based, cross-sectional study between September and
13 November 2011 in the JD-HDSS of Bhaktapur district in the Kathmandu Valley, 13 kilometers
14 from Kathmandu. An earlier report described the establishment of the surveillance site.²¹
15 Duwakot and Jhaukhel each contain nine administrative wards. We randomly selected three
16 wards from each village. Based on the JD-HDSS 2010 baseline census,²¹ we compiled a
17 complete list of households containing adults aged 25-59 years in the selected wards. Next, we
18 selected one adult, of either sex, from all interview households. If a household contained more
19 than one adult, we used the Kish technique to select the respondent.²³
20
21
22
23
24
25
26
27
28

29 **Study tools**

30 Interviews based on the WHO-NCD STEPS questionnaire

31 We structured the basic questionnaire using the WHO-NCD STEPS²⁴ and other sources.^{17, 25-39}
32 Questions were related to respondents' knowledge of cardiovascular health (e.g., causes and risk
33 factors, warning signs, and action for suspected heart attack); heart-healthy diet; and disease
34 prevention. The first two areas included unprompted (spontaneous, open-ended) and prompted
35 (clues provided, closed-ended) responses, and the latter two areas included only prompted
36 responses. We asked respondents about their attitude toward different aspects of cardiovascular
37 health: did they perceive themselves at risk for CVD? Did they think behavior change would
38 reduce their risk of developing CVD?
39
40
41
42
43
44
45
46
47

48 Behavior-related questions were based on Step 1 of the WHO STEPwise approach to
49 surveillance (STEPS), which uses a questionnaire to collect data on risk factors.²⁴ Our questions
50 probed current and past smoking behavior; smokeless tobacco; alcohol consumption in the past
51 30 days (current drinker), and drinking frequency among current drinkers. The physical activity
52 section asked respondents to report the number of days and amount of time they participated in
53 vigorous and moderate activities at work, leisure, and travel. After calculating the Metabolic
54
55
56
57
58
59
60

1
2
3 Equivalent of Task (MET) minutes/week, we grouped participants as highly active, moderately
4 active, or inactive (low physical activity) in accord with the STEPS manual. Additional practice
5 and behavioral questions explored whether participants had engaged in any heart-friendly
6 behavior in the previous year and for what reason.
7
8
9

10
11 Cronbach's alpha, which measures question reliability, for the overall knowledge and attitude
12 was 0.941 (range: 0.775–0.913) and 0.824 (range: 0.791–0.822), respectively. Cronbach's alpha
13 for practice and behavior ranged from low (0.257) for physical activity questions to high (0.929)
14 for blood pressure measurement.
15
16
17

18
19 Twelve local female enumerators (two per ward) conducted the interviews. All enumerators had
20 completed at least grade 10 in school. Supervisors included two public health graduates; a field
21 coordinator who holds both master's and PhD degrees in public health; two PhD students
22 including one author (AV).
23
24
25
26

27 Scoring knowledge, attitude, and practice/behavior questions

28
29 To compare and correlate the three components, we scored responses regarding KAP/behavior
30 for cardiovascular health. The maximum possible score for KAP/behavior was 53, 70, and 25
31 respectively. See Supplement for further detail.
32
33
34

35 Physical measurements

36
37 Enumerators recorded anthropometric measurements (i.e., weight, height, and waist and hip
38 circumferences) according to STEPs guidelines.²⁴ Weight was measured to one decimal place
39 using a Microlife BR-9201 Weighing Scale (Microlife AG Swiss Corporation, Widnau,
40 Switzerland). Height and waist and hip circumferences were measured using non-stretchable
41 tapes (Jonson Tapes Ltd, New Delhi, India). Blood pressure was determined digitally by
42 Microlife BP 3AP1-3E (Microlife AG Swiss Corporation, Widnau, Switzerland). Enumerators
43 recorded three readings for each respondent, taken at 5-minute intervals. We averaged the
44 readings to attain the final blood pressure measurements. Respondents with known history of
45 hypertension (diagnosed cases) and those diagnosed during the study according to Joint National
46 Committee -VII⁴¹ were termed as having "hypertension."
47
48
49
50
51
52
53
54

55 **Data management and analysis**
56
57
58
59
60

Supervisors checked all completed questionnaires and, if necessary, asked enumerators to revisit households. We outsourced data entry to a team of data entry operators, public health graduates by qualification. Data was coded and entered in Epidata version 2.1 and analyzed with SPSS version 17.0 (IBM, Armonk, New York, USA) and STATA version 10.0 (StataCorp, Texas, USA). Demographic variables were taken as independent variables; other variables, including KAP components, were analyzed as dependent variables. Categorical data is presented as percentages, and continuous data is presented as mean (standard deviation). KAP scores are presented as median score percents. We used the Chi-square test to compare proportions and the Kruskal–Wallis test to compare medians. $P < 0.05$ was considered statistically significant.

Ethical considerations

The Nepal Health Research Council and the Institutional Review Board of Kathmandu Medical College approved this study. Following a pattern established at the outset of JD-HDSS, we consulted local leaders and sought their informal permission for HDSS activities. In addition, we periodically consulted with health and administrative authorities in both villages. All participants gave informed consent, and privacy was maintained during the interview process. To maintain confidentiality, all data were secured in the HDSS office. Computer data were accessible only to the research team. Although respondents received no monetary incentives, we provided a referral that included discounted consultation fees at Kathmandu Medical College or Nepal Medical College for self-reported illness and newly diagnosed hypertension.

RESULTS

We surveyed 789 of 840 households in randomly selected six wards of Duwakot and Jhaukhel (non-response rate = 6.07%). After omitting incomplete questionnaires, we included 777 individuals in the analysis. Except for missing recordings for height, weight, or blood pressure, the data reported here cover all 777 respondents.

Demographic variables

Table 1 describes the respondents. Seventy percent of all participants were female, and 25% lacked formal education.

Table 1: Demographic characteristics and risk factor status of the study population (n=777)

Demographic variables (n=777)	Male (%)	Female (%)	Total (%)
Age (years)			
25–34	30.1	34.3	33.1
35–44	34.1	36.9	36.0
45–59	35.8	28.8	30.9
Ethnic group			
Brahmin	37.1	38.1	37.8
Chhetri	24.0	25.4	25.0
Newar	29.3	23.2	25.0
Minorities	9.6	13.3	12.2
Education*			
No formal schooling	11.8	33.2	26.9
Primary school	32.3	31.4	31.7
Secondary school	32.3	22.6	25.5
High school or more	23.6	12.8	16.0
Occupation[†]			
Employee	30.6	8.4	14.9
Self-employed	32.4	6.6	13.9
Housewife		67.3	47.5
Agriculture	21.0	14.2	16.3
Other	17.0	3.5	17.5
Cardiovascular risk factors[§]			
Smoking (current) (n=777)	33.5	14.7	20.1
Smokeless tobacco (current) (n=777)	20.2	3.3	8.3
Alcohol drinkers (ever) (n=777)	36.7	14.1	20.7
Alcohol drinkers (current) (n=777)	34.5	12.6	19.0
Low physical activity (n=777)	38.3	45.2	43.3
Hypertension (known and new) (n=670)	20.6	22.1	21.6

Self-reported high blood sugar (n=777)	4.8	3.5	3.9
Overweight/obese (n=762)	25.0/5.4	31.6/11.2	29.7/9.4
Increased waist circumference (cm) (n=773)	21.6	56.6	46.3
Increased waist/hip ration (n=750)	57.9	55.6	56.3

Note: Classification of ethnic groups is based on the National Central Bureau of Statistics.⁴¹

*Primary and secondary schooling indicate education up to grade 4 and 10, respectively.

†Employee describe individuals working for governmental or nongovernmental organizations. “Other” occupations include students, unemployed individuals (able or unable to work), retired, and non-paid workers.

§Risk factor categories are based on the WHO-NCD Risk Factor STEPS Survey manual.²⁴ Current smokers included those who responded “yes” to “Do you smoke?” Past smokers included those who replied “yes” to “Did you ever smoke in the past?” “Ever drinkers” included respondents who had “consumed a drink that contained alcohol ever in lifetime,” and current drinkers included respondents who had “consumed alcohol within the previous month.” Percentage of drinking frequency represents current drinkers who consumed at least one standard drink per occasion. Blood pressure data exclude respondents who did not submit to all three readings. Likewise, data for body mass index, waist circumference, and waist-hip ratio exclude respondents whose weight, height, and waist and/or hip measurements were not taken. Increased waist circumference includes waist measurements of ≥ 80 cm (females) and ≥ 90 cm (males); increased waist-hip ratio is ≥ 0.85 (females) and ≥ 0.90 (males).

High burden of risk factors related to cardiovascular health

Table 1 also shows the prevalence of cardiovascular risk factors in the study population. One in four respondents was a past or current smoker. One in five respondents had consumed at least one alcoholic beverage in the previous month. Both males and females consumed fruit 2.6 ± 1.98 and 3.3 ± 2.17 days/week, respectively. Likewise, males and females consumed vegetables 5.33 ± 1.69 and 5.64 ± 1.55 days/week, respectively. More than 40% of the respondents reported low physical activity. Prevalence of known and newly diagnosed hypertension was 21.6%. As measured by BMI, the prevalence of overweight and obesity was 29.7% and 9.4 %, respectively. Abdominal obesity (i.e., increased waist circumference and waist-hip ratio) was 46.3% and 56.3%, respectively (Table 1).

Poor knowledge of cardiovascular health

Cause of heart disease

When seeking unprompted responses, enumerators asked participants to spontaneously name the causes of CVD (Table 2). Respondents showed low overall knowledge of the factors that cause CVDs, ranging from 1.0% (high blood sugar) to 29.2% (smoking). Generally, males knew more

than females; younger respondents knew more than older ones; ethnic minorities knew less; and better-educated individuals and governmental or nongovernmental employees were better informed than those with less education and in other employment situations.

Table 2: Percentage of respondents citing various causes of heart disease (n=777)

	Hypertension	High blood sugar	High cholesterol	Physical inactivity	Overweight	Smoking	Smokeless tobacco	Excessive alcohol	Excess stress
Sex									
Male	18.8*	1.3	20.5*	7.0	5.7	34.9*	8.3	31.4*	16.2
Female	10.9*	0.9	13.5*	4.0	5.3	26.8*	4.7	23.9*	13.5
Age (years)									
25–34	14.8	1.6	17.5	5.4	6.2	31.1	6.2	28.4	16.0*
35–44	13.2	0.4	17.1	5.7	6.1	26.8	5.0	25.0	17.1*
45–59	11.7	1.3	11.7	3.3	3.8	30.0	6.3	25.0	9.2*
Ethnicity									
Brahmin	14.3*	1.0	21.1 [#]	7.5*	5.8	29.3	8.2	25.2	12.9
Chhetri	19.1*	1.0	16.5 [#]	4.6*	6.7	32.5	3.6	25.8	20.1
Newar	10.8*	.5	9.8 [#]	3.6*	5.7	27.8	6.2	28.4	10.8
Minorities	3.2*	2.1	8.4 [#]	0.0*	1.1	25.3	2.1	25.3	13.7
Education									
Primary school	13.0*	0.8	14.2 [#]	5.3 [#]	4.5*	30.5	8.1	27.6	15.0*
Secondary school	15.7*	1.5	18.7 [#]	8.6 [#]	8.6*	29.8	3.0	27.8	16.7*
High school or more	21.0*	1.6	27.4 [#]	4.8 [#]	8.9*	33.9	8.1	26.6	18.5*
No formal schooling	6.7*	0.5	7.2 [#]	1.0 [#]	1.4*	24.4	4.3	22.5	8.6*
Occupation									
Employee	26.7 [#]	0.9	32.8 [#]	10.3	8.6	30.2	8.6	31.9	23.3*
Self-employed	17.6 [#]	2.8	14.8 [#]	1.9	7.4	29.6	2.8	25.9	15.7*
Housewife	10.6 [#]	1.1	11.7 [#]	3.3	5.4	25.7	3.5	23.8	13.0*
Agriculture	7.1 [#]	0.0	14.3 [#]	7.1	1.6	37.3	8.7	26.2	11.1*

Others	8.6 [#]	0.0	10.3 [#]	5.2	3.4	31.0	13.8	29.3	8.6*
Overall	13.3	1.0	15.6	4.9	5.4	29.2	5.8	26.1	14.3

Note: Figures are based on spontaneous participant responses. The table includes only established CVD risk factors and excludes responses that cited other non-established causes (e.g., food hygiene, air pollution, etc.). Calculated with Chi-square test, p-values compare all categories in the variables.

*P-value <0.05; # p-value <0.01]

Knowledge of heart attack symptoms

Around 60% of respondents did not know any heart attack symptoms compared with 20% who knew 2–4 symptoms. This ratio was uniform across all the demographic variables (Table 3).

Importantly, less than 25% of respondents identified chest pain—the most prominent and important indicator of heart attack—as a symptom (data not shown). This knowledge decreased with age (p=0.013) and level of education (p=0.055). Brahmin respondents (p<0.001) and governmental or nongovernmental employees (p=0.021) had slightly better knowledge about chest pain.

Table 3: Correctly reported warning signs of heart attack (n=777)

	Signs reported (number)			P-value
	0 (%)	1 (%)	2–4 (%)	
Sex				
Male	59.4	18.3	22.3	0.605
Female	59.5	20.8	19.7	
Age (years)				
25–34	63.4	16.7	19.8	0.494
35–44	57.1	21.4	21.4	
45–59	57.9	22.1	20.0	
Ethnicity				
Brahmin	56.8	22.1	21.1	0.395

Chhetri	62.9	18.0	19.1	
Newar	59.3	17.0	23.7	
Others	61.1	24.2	14.7	
Highest education				
No formal schooling	60.3	22.0	17.7	
Up to primary school	58.1	18.7	23.2	0.493
Up to secondary school	60.1	22.2	17.7	
High school or more	59.7	16.1	24.2	
Main job				
Employee	55.2	19.0	25.9	
Self-employed (self)	57.4	20.4	22.2	
Housewife	59.1	21.4	19.5	0.687
Agriculture	65.1	15.9	19.0	
Others	62.1	22.4	15.5	
Overall	59.5	20.1	20.5	

Note: Warning signs were based on a list provided by American Heart Association.⁴²

P-values were calculated using the Chi-square test and compare all categories in the variables.

Action in case of a heart attack

When asked what action they would take for a suspected heart attack, 75% of respondents said they would immediately take the person to a hospital. About 3% said they would try home therapy, consult a traditional healer, or stay home and wait for the symptoms to go away. More than 20% gave no response.

Heart-healthy foods

When asked to identify a particular heart-healthy food, most respondents identified green leafy vegetables (94.6%) and fruit (92.1%). Around 7% thought fried food or saturated fat (e.g., ghee) are healthy foods. Nearly 22% thought salty food (e.g., chips) is heart-healthy.

Heart disease prevention

After reading aloud a list of specific actions that help prevent CVD, enumerators asked respondents to comment if they thought a particular action would help. Actions included quitting smoking; getting physical exercise; losing weight; reducing intake of dietary fat, animal products, and salt; reducing stress; and maintaining healthy blood pressure. A majority of respondents thought these actions are helpful, ranging from 86.1% for weight reduction to 92.5% for smoking cessation.

Attitude toward heart health in an urbanizing community

Enumerators asked respondents about their attitude toward different issues of heart health (Table 4). Regarding lifestyle and preventive measures, 72.9% of males and 68.0% females did not perceive themselves at risk of CVD and more than half did not want to change their current lifestyle (males = 63.3%; females = 53.1%). Seventy-five percent said they did not eat more than they need; 80% did consider themselves overweight; and three fifths did not think that changing their behavior would lower their cardiovascular risk. Further, 52.4% of males and 60.8% of females thought that God or a higher power determines their health.

Table 4: Response to attitude-related questions on heart health

Attitude	Strongly disagree		Somewhat disagree		Somewhat agree		Strongly agree		Don't know/ Can't say	
	M	F	M	F	M	F	M	F	M	F
I don't perceive myself to be at risk for heart disease	31.4	30.8	41.5	37.2	15.7	21.9	9.6	8.6	1.7	1.5
I don't want to change my lifestyle (activity, smoking, alcohol)	17.5	20.3	17.9	25.4	30.1	20.8	33.2	32.3	1.3	1.3
I feel the amount of food I eat is too much, and I would like to cut down	33.6	28.5	49.3	45.1	10.0	15.3	5.2	10.2	1.7	0.9

I feel my present weight is too high for my health	36.2	34.9	47.6	41.4	7.9	11.9	7.0	10.4	1.3	1.5
I don't think changing my behavior will reduce my risk of developing heart disease	13.1	15.0	21.8	23.4	38.4	32.5	24.9	24.5	1.7	4.7
I don't have time to take care of myself	44.1	39.4	22.3	29.9	13.5	14.4	18.3	14.6	1.7	1.6
God or some higher power ultimately determines my health	25.8	23.7	20.1	13.5	25.3	22.3	27.1	38.5	1.7	2.0
Opinions on various preventive aspect of heart health	Very helpful	Somewhat helpful	Not very helpful	Not helpful	Don't know					
Access to more fruits, vegetables, and other healthy foods	93.4	93.2	5.2	3.8	0.4	0.2	0.4	0.7	0.4	2.0
Greater access to indoor and outdoor public recreational facilities	56.3	67.3	31.0	22.4	7.0	3.6	0.9	0.5	4.8	6.0
Smoking bans	92.1	92.0	3.9	4.2	0.9	0.9	1.7	0.7	1.3	2.2
More awareness programs for healthy heart	85.2	87.0	9.6	8.6	3.1	0.9	0.9	0.4	13	31
Availability of health facilities to detect and treat heart diseases	89.1	87.6	7.0	7.7	1.7	0.7	0.9	0.7	1.3	3.3

Local volunteers that shall try to change any adverse health behavior	65.1	64.6	17.0	20.6	3.1	2.9	1.7	0.4	13.1	11.5
---	------	------	------	------	-----	-----	-----	-----	------	------

Note: Figures indicate percentage of respondents among males (M) and females (F).

Next, enumerators asked for respondent opinions on various preventive aspects of heart health (e.g., improved access to fruits, vegetables, and other healthy foods) (Table 4). Most thought that smoking bans or increased access to fruits, vegetables, and other healthy foods would be very helpful. About two thirds of all respondents thought that greater access to indoor and outdoor public recreational facilities would be very helpful and one quarter thought it would be somewhat helpful. Around 10% of females seemed unsure whether local health volunteers trying to change the community's health practice/behavior would be useful.

Practice/behavior in relation to heart disease

Enumerators elicited practice- and behavior-related responses in two ways. First, they asked WHO STEPs-based questions about four major behavior-related risk factors (i.e., tobacco smoking, alcohol consumption, physical activity, and fruit and vegetables) (Table 1). Second, they asked respondents if they had participated in any heart-friendly practice/behavior in the previous year (e.g., blood pressure, blood sugar, or other diagnostic tests; tobacco cessation; increased physical activity; weight loss). Half of all respondents had their blood pressure measured and 10% reported getting a diagnostic test for CVD. About 60% attempted to reduce their consumption of unhealthy foods and 25% tried to lose weight or increase physical activity. Half of the smokers said they had attempted to quit smoking.

When enumerators asked why responders who reported one or more positive health behaviors tried to change their lifestyle or practice/behavior, participants responded that they wanted to feel better (70%); had seen, read, or heard information (65.4%); did it for their family (45.9%); or were encouraged to do so by a healthcare professional (41.3%), family member (39.0%), or friend (31.8%). Other reasons for changing lifestyle or practice/behavior included wanting avoid taking medications (20.6%); a family member, relative, or friend who developed CVD, became ill, or died (16.5%); and developing possible cardiovascular symptoms (14.8%).

KAP scores and demographic variables

Median percent scores were similar across the various demographic variables (Table 5). Males, younger people, or better-educated respondents scored 2%–8% higher than females, older respondents, and the less educated. Similarly, median percent scores for attitude tended to decrease about 2% in older respondents, Newars, and those without formal schooling. Behavioral median percent scores were 4%–8% higher among younger and more-educated respondents, Brahmins, Chhetris, and government employees or housewives (Table 5).

Table 5: Median percent score of knowledge, attitude, and practice/behavior according to demographic variables

	Knowledge median percent score	P-value	Attitude median percent score	P-value	Behavior median percent score	P-value
Sex						
Male	79.2	<0.001	74.29	0.139	48.00	0.660
Female	77.4		74.29		48.00	
Age (years)						
25–34	79.79	0.005	74.29	0.074	48.00	0.018
35–44	79.24		74.29		44.00	
45–59	75.47		72.86		44.00	

Ethnic group						
Brahmin	75.47		74.29		48.00	
Chhetri	79.24	<0.001	74.29	<0.001	48.00	0.030
Newar	79.24		72.14		44.00	
Other	79.24		74.29		44.00	
Education						
No formal schooling	75.47		72.86		44.00	
Primary school	77.35		74.29		44.00	
Secondary school	79.24	<0.001	74.29	0.001	48.00	<0.001
High school or more	83.02		75.71		52.00	
Occupation						
Employed	79.24		74.29		48.00	
Self-employed	79.24		74.29		44.00	
Housewife	79.24	<0.001	74.29	0.940	48.00	0.156
Agriculture	75.47		74.29		44.00	
Other	79.24		74.29		48.00	
Overall	79.24		74.29		48.00	

Note: Knowledge scores were based on prompted responses.

Calculated with the Kruskal-Wallis test, p-values compared all categories in the variables.

Correlation between knowledge, attitude, and practice/behavior scores

Scatter-plots (Figure 1a–1c) show the correlations between the three components. We observed a poor positive correlation between knowledge and attitude median scores (Pearson correlation: 0.025, $p=0.480$) [Figure 1a]) and a negative correlation between attitude and behavior (Pearson correlation: -0.126 , $p<0.001$) [Figure 1b]). However, we observed a positive correlation between knowledge and behaviour (Pearson correlation: 0.121, ($p=0.001$) [Figure 1c]).

DISCUSSION

The present study is the first to conduct a detailed analysis of the relationship between KAP/health literacy and cardiovascular health in a semi-urban setting near Kathmandu.

Cardiovascular risk factors are abundant in the study population

1
2
3 Our respondents showed a high burden of behavioral and biological risk factors. The prevalence
4 of tobacco smoking and vegetable and fruit consumption were similar to national averages and
5 alcohol consumption was lower.⁴³ In accord with our earlier review on obesity in Nepal, the
6 effect of urbanization is evident in low physical activity and overweight/obesity that exceeds the
7 national average.^{22, 43} The prevalence of hypertension in our study population is similar to the
8 national average (22%) but less by about one third than a study in Bhadrabas, a similar semi-
9 urban Kathmandu community,⁴⁴ most probably due to different age groups. Self-reported
10 prevalence of diabetes mellitus (3.9%) confirms the HDSS baseline figure reported earlier.²¹
11
12

13 **Poor knowledge of cardiovascular health**

14 Our study population showed low overall knowledge of the causes of CVD. Unprompted
15 responses suggested that respondents identified behavioral factors (e.g., smoking) more
16 frequently than physiological causes (e.g., hypertension). Other studies have reported similar
17 findings for unprompted³⁸ and prompted responses.³⁴ Conversely, unhealthy diet, obesity, and
18 family history were common responses in Taiwan,³² and American women frequently identified
19 obesity.³⁵ Many of our respondents also attributed CVD to environmental factors such as food
20 hygiene and air pollution. Indeed, acute inflammatory reactions to air pollutants accelerate
21 coronary atherosclerosis and trigger arrhythmia or myocardial infarction.⁴⁵ Unlike the Strong
22 Heart Study³⁹ in American populations, our results did not exhibit a higher probability of
23 knowledge of a risk factor in persons who have that risk factor, demonstrating poor
24 cardiovascular health literacy even among those already affected.
25
26

27 Similar to Pakistan,¹⁷ a large majority (60%) of our respondents could not identify even one heart
28 attack symptom. However, similar to an earlier study in Nepal,¹⁸ 6%–22% of individuals in our
29 demographic subsets recognized chest pain as a symptom, which is less compared to 60% of
30 Vietnamese Americans who spontaneously mentioned chest pain.⁴⁷ Importantly, two thirds of
31 heart attack patients delayed their arrival at a Karachi hospital because they did not recognize
32 heart attack symptoms.¹⁷ Similar to a previous study,¹⁸ three quarters of our respondents believed
33 they would immediately take a suspected heart attack patient to a hospital but one fifth did not
34 answer the question or opted instead for home therapy or traditional healing. Combined with our
35 earlier finding of not knowing heart attack symptoms, this lack of knowledge about appropriate
36 action underscores the need to educate the Nepalese population about this issue. In contrast, 85%
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 of Vietnamese Americans knew that they should immediately call 911 if they suspect a heart
4 attack.⁴⁶ Despite inadequate knowledge about the causes of CVD, our respondents were much
5 better informed about prevention measures. This finding concurs with the perception level of
6 American women,³⁵ probably because enumerators asked prompted questions in this section of
7 the questionnaire. In Taiwan, only 25% of respondents thought that CVD is preventable and 50%
8 were “not sure.”³²

14 **Inappropriate attitude toward cardiovascular health**

17 In contrast to their opinion that behavioral modifications can prevent CVD, most of our
18 respondents did not consider themselves at risk and did not want to alter their lifestyle. This may
19 be explained by the Health Belief Model⁴⁷ (i.e., a person may be knowledgeable about disease
20 prevention but is more likely to undertake prevention when s/he perceives herself/himself to be
21 at risk of developing the disease). Others have described a similarly optimistic bias.⁴⁸ Risk
22 indicators (e.g., number of cigarettes smoked; death of a parent due to CVD; and self-assessment
23 of blood pressure, weight, and cholesterol levels) account for a 24% variance in perceived risk.⁴⁸
24 The likelihood of perceiving oneself at higher risk increases when the presence of a risk factor is
25 known.³² However, respondents often underestimate their cardiovascular risk.⁴⁸

34 **Attempts toward better cardiovascular health practice/behavior**

36 Compared to Nepal’s national average (69%), only half of our respondents underwent blood
37 pressure measurement in the previous year.⁴⁴ Ninety percent of the known hypertensive
38 population had their blood pressure under control, exceeding the 6% and 9.5% control rate
39 reported in two other suburban Kathmandu studies^{44,49} and an urban study in Eastern Nepal
40 (58.5%).⁵⁰ Half of the smokers reported here that they had tried to quit smoking. Our
41 questionnaire did not explore whether they had actually quit and did not identify the barriers
42 against quitting, information that might be important for public health. The theory of reasoned
43 action⁴⁷ argues that intention is a critical component of healthy behavior. Although confirming
44 the genuineness of intention is difficult, such intention may provide an opportunity for public
45 health intervention. Importantly, we identified our respondents’ reasons for attempting positive
46 health behavior. Many constructs of the Health Belief Model,⁴⁷ such as perceived benefit
47 (“wanted to feel better”), cues to action (“encouraged by a health professional, etc.”), fear (“death
48 of a family member, etc.”), appear to affect behavioral decisions.

Gender

Seventy percent of our respondents were women. We reported earlier a male: female ratio (1.15) in the population.²¹ Enumerators used the Kish technique to select respondents at the household level, expecting a proportionate sampling.²³ However, enumerators might have interviewed more women than men because women were more likely to be available in the households when enumerators visited. Nevertheless, our analysis includes a gender-wise stratification, and we present data separately whenever we observed marked gender differences. In our study, more men than women smoked tobacco or drank alcohol. The prevalence of hypertension and self-reported diabetes was similar in both groups. On the other hand, low physical activity and obesity were higher in the women. Although men had better knowledge of heart disease, their attitude was similar to those reported by women. However, more women than men strongly agreed that their health was ultimately determined by God or a higher power. A Canadian study³⁶ observed a similar knowledge level among women and men. One major factor for the gender discrepancy in our study is that more women (33.2%) than men (11.8%) lacked formal education.

Effect of other socio-demographic variables on KAP

Age-wise, KAP/behavior scores decreased around 4% from the youngest to the oldest age group. An earlier study reported a higher level of knowledge among young respondents.³⁶ Interestingly, a study in adolescents reported a very poor level of cardiovascular knowledge.⁵¹ Otherwise, knowledge levels rise linearly from childhood until middle age, when the level starts to plateau.⁵² In our study, better-educated respondents ate more fruit and consumed less tobacco and alcohol, but they also reported less physical activity and more hypertension. The highest education category (high school or more) scored higher than the lowest category (no formal schooling) but only by 3%–8%. Other studies observed a consistent link between better KAP scores and higher education.^{32, 36, 53} Occupation-wise, employed respondents knew more about biological causes (e.g., high blood pressure, diabetes, and high cholesterol), possibly due to more education, wider exposure to information, and better access to screening and treatment. A Canadian study reported similar findings.³⁶

Prompted vs. unprompted responses

1
2
3 Although we used unprompted responses to evaluate general knowledge, we used prompted
4 responses to calculate scores, thus providing a denominator (“full marks”) to calculate median
5 score percent. In other settings, KAP studies have variably used prompted^{34, 39, 46, 54} or
6 unprompted questions,^{18, 32, 38} some studies used both.³⁵ Other studies did not identify their
7 chosen method.³³ Although the accuracy of prompted vs. unprompted questions in knowledge
8 assessment has been a topic of debate, prompted questions confer a higher level of knowledge.⁵⁵
9 When we analyzed knowledge responses based on prompted responses, the percent of correct
10 answers increased drastically. The psychological explanation is that respondents must simply
11 “recognize” answers to prompted questions, but they must engage in the more difficult mental
12 process of “recalling” answers to unprompted questions.⁵⁶ We believe that unprompted responses
13 more accurately reflect knowledge level because they emerge from a thought process that
14 involves respondents’ knowledge, experience, and perception regarding a specific issue.
15
16
17
18
19
20
21
22
23
24

25 **Poor correlation between KAP scores**

26
27 Poor correlation between KAP scores was unsurprising because a linear relationship between
28 variables may not exist in cardiovascular health. Valente argued that each of six possible
29 behavior change models contains a varying order of KAP/practice and that a population can be
30 described in terms of the percentage that fits into each model.⁵⁷ This implies that no universal
31 intervention can improve the cardiovascular health KAP of every individual in a community. The
32 challenge lies in identifying individuals who fit into different linear models. Furthermore, health
33 behavior theories (e.g., social cognitive theory and the theory of reasoned action⁴⁷) suggest that
34 knowledge and attitude are only two components of the pathways that lead to behavior.
35 Constructs (e.g., self-efficacy) and other personal and environmental factors (in social cognitive
36 theory) and perceived behavioral control, subjective norms, and ultimately intention (in the
37 theory of reasoned action) play crucial roles in shaping one’s health behavior.⁴⁷
38
39
40
41
42
43
44
45
46

47 **CONCLUSION**

48
49 Our community-based cross-sectional HARDIC study demonstrated poor knowledge,
50 inappropriate attitude, and highly adverse behavior regarding cardiovascular health in a semi-
51 urban community of Nepal. We aimed to determine a baseline KAP in the community to aid the
52 planning of health promotion activities. Our results show the potential and possibility that
53 community health promotion can improve cardiovascular health literacy.
54
55
56
57
58
59
60

Acknowledgments

The authors are grateful to the study participants. We acknowledge all enumerators; field supervisors (Rachana Shrestha, Vishal Bhandari, Chandra Shova Khaitu, and Shova Poudel); and our field coordinator, Prof. Dr. Muni Raj Chhetri. We thank Umesh Raj Aryal for field supervision and advice regarding statistical analysis. We thank scientific editor Karen Williams (Kwills Editing Services, Weymouth, MA, USA) for providing professional English-language editing of this article.

Contributors

AV designed the study, performed statistical analysis, and drafted the manuscript. AK helped design the study and provided critical revision of the manuscript. Both authors have read and approved the final manuscript.

Funding Statement

This study was supported by grants from the Wilhelm & Martina Lundgren's Foundation (vet1-367/2011 and vet1-379/2012) and the University of Gothenburg, Sweden through a "Global University" grant (A11 0524/09).

Competing interests

None declared.

Data sharing

No additional data available.

References

1. Alwan A. *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organization; 2011.
2. World Health Organization. *Noncommunicable diseases in the South-East Asia Region: Situation and response 2011*. New Delhi: World Health Organization, Regional Office for South-East Asia; 2011.
3. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998;97:596-601.
4. Reddy KS. Cardiovascular diseases in the developing countries: dimensions, determinants, dynamics and directions for public health action. *Public Health Nutr* 2002;5:231-37.
5. Engelgau MM, El-Saharty S, Kudesia P, Rajan V, Rosenhouse S, Okamoto K. *Capitalizing on the demographic transition: tackling noncommunicable diseases in South Asia*. Washington, DC: World Bank Publications; 2011.
6. Mathers C, Fat DM, Boerma J. *The global burden of disease: 2004 update*. Geneva: World Health Organization; 2008.
7. Ezzati M, Lopez AD, Rodgers A, et al. *Comparative quantification of health risks: global and Regional Burden of Disease Attributable to Selected Major Risk Factors*. Geneva: World Health Organization; 2004.
8. Khaw KT, Wareham N, Bingham S, et al. Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS Medicine* 2008;5:e12.
9. Mendis S, Puska P, Norrving B. *Global atlas on cardiovascular disease prevention and control*. Geneva: World Health Organization; 2011.
10. Alwan A, Maclean DR, Riley LM, et al. Monitoring and surveillance of chronic non-communicable diseases: progress and capacity in high-burden countries. *Lancet* 2010; 376:1861-68.
11. Mendis S, Alwan A. *Prioritized research agenda for prevention and control of noncommunicable diseases*. Geneva: World Health Organization; 2011.
12. Puska P. From Framingham to North Karelia: from descriptive epidemiology to public health action. *Prog Cardiovasc Dis* 2010;53:15-20.
13. Pearson TA, Bazzarre TL, Daniels SR, et al. American Heart Association guide for improving cardiovascular health at the community level: a statement for public health

- 1
2
3 practitioners, healthcare providers, and health policy makers from the American Heart
4 Association Expert Panel on Population and Prevention Science. *Circulation* 2003;107:645-51.
5
6
7 14. Lloyd-Jones DM, Hong Y, Labarthe D, *et al.* Defining and setting national goals for
8 cardiovascular health promotion and disease reduction: the American Heart Association's
9 strategic impact goal through 2020 and beyond. *Circulation* 2010;121:586-613.
10
11 15. Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health
12 education and communication strategies into the 21st century. *Health Promot Internation*
13 2000;15:259-67.
14
15 16. Gazmararian JA, Williams MV, Peel J, *et al.* Health literacy and knowledge of chronic
16 disease. *Patient Educ Couns* 2003;51(3):267-75.
17
18 17. Khan MS, Jafary FH, Faruqui AM, *et al.* High prevalence of lack of knowledge of symptoms
19 of acute myocardial infarction in Pakistan and its contribution to delayed presentation to the
20 hospital. *BMC Public Health* 2007;7:284.
21
22 18. Limbu YR, Malla R, Regmi SR, *et al.* Public knowledge of heart attack in a Nepalese
23 population survey. *Heart & Lung: The Journal of Acute and Critical Care* 2006;35:164-69.
24
25 19. Upadhyay DK, Palaian S, Shankar PR, *et al.* Knowledge, attitude and practice about diabetes
26 among diabetes patients in Western Nepal. *Rawal Med J* 2008;33:8-11.
27
28 20. Vaidya A. Tackling cardiovascular health and disease in Nepal: epidemiology, strategies and
29 implementation. *Heart Asia* 2011;3:87-91.
30
31 21. Aryal UR, Vaidya A, Shakya-Vaidya S, *et al.* Establishing a health demographic surveillance
32 site in Bhaktapur district, Nepal: initial experiences and findings. *BMC Research Notes*
33 2012;5:489.
34
35 22. Vaidya A, Shakya S, Krettek A. Obesity prevalence in nepal: public health challenges in a
36 low-income nation during an alarming worldwide trend. *International Journal of Environmental*
37 *Research and Public Health* 2010;7:2726-44.
38
39 23. Kish L. Sampling organizations and groups of unequal sizes. *Am Sociol Rev* 1965; 30:564-
40 72.
41
42 24. WHO STEPS Instrument (Core and Expanded): The WHO STEPwise approach to chronic
43 disease risk factor surveillance (STEPS). Available at:
44 http://www.who.int/chp/steps/STEPS_Instrument_v2.1.pdf.
45
46 25. Lin W, Yang H, Hang C, *et al.* Nutrition knowledge, attitude, and behavior of Taiwanese
47 elementary school children. *Asia Pac J Clin Nutr* 2007;16:534-46.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
26. Torabi MR, Yang J, Li J. Comparison of tobacco use knowledge, attitude and practice among college students in China and the United States. *Health Promot Internation* 2002;17:247-53.
27. Kosaryan M, Vahidshahi K, Siami R, *et al.* Knowledge, attitude, and practice of reproductive behavior in Iranian minor thalassemia couples. *Saudi Med J* 2009;30:835-39.
28. Neamatollahi H, Ebrahimi M, Talebi M, *et al.* Major differences in oral health knowledge and behavior in a group of Iranian pre-university students: a cross-sectional study. *J Oral Sci* 2011;53:177-84.
29. Sharda AJ, Shetty S. A comparative study of oral health knowledge, attitude and behaviour of first and final year dental students of Udaipur city, Rajasthan, India. *International Journal of Dental Hygiene* 2008;6:347-53.
30. Yap J, Lee V, Yau T, *et al.* Knowledge, attitudes and practices towards pandemic influenza among cases, close contacts, and healthcare workers in tropical Singapore: a cross-sectional survey. *BMC Public Health* 2010;10(1):442.
31. Maimaiti N, Shamsuddin K, Abdurahim A, *et al.* Knowledge, Attitude and Practice regarding HIV/AIDS among University students in Xinjiang. *Global Journal of Health Science* 2010;2:51.
32. Chen W, Yu Y, Glaser K. The Knowledge and Attitudes of Coronary Heart Disease Prevention among Middle and Older Aged People in a Community in Taipei. *Taiwan Geriatrics & Gerontology* 2009;4:251-62.
33. Dodani S, Mistry R, Khwaja A, *et al.* Prevalence and awareness of risk factors and behaviours of coronary heart disease in an urban population of Karachi, the largest city of Pakistan: a community survey. *Journal of Public Health* 2004;26:245-49.
34. Saeed O, Gupta V, Dhawan N, *et al.* Knowledge of modifiable risk factors of Coronary Atherosclerotic Heart Disease (CASHD) among a sample in India. *BMC International Health and Human Rights* 2009;9:2.
35. Mosca L, Jones WK, King KB, *et al.* Awareness, perception, and knowledge of heart disease risk and prevention among women in the United States. *Arch Fam Med* 2000;9(6):506.
36. Potvin L, Richard L, Edwards AC. Knowledge of cardiovascular disease risk factors among the Canadian population: relationships with indicators of socioeconomic status. *Can Med Assoc J* 2000;162:S5-S11.
37. Lim K, Sumarni M, Amal N, *et al.* Tobacco use, knowledge and attitude among Malaysians age 18 and above. *Trop Biomed* 2009;26:92-99.
38. Wong BM, Garcia Y, Barr A, *et al.* Cardiovascular risk factor awareness in a disadvantaged inner-city population—implications for preventive strategies. *Can J Cardiol* 2008;24:677-82.

- 1
2
3 39. Schweigman K, Eichner J, Welty TK, *et al.* Cardiovascular disease risk factor awareness in
4 American Indian communities: The Strong Heart Study. *Ethnicity and disease* 2006;16:647.
5
6
7 40. Lenfant C, Chobanian AV, Jones DW, *et al.* Seventh Report of the Joint National Committee
8 on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7).
9 *Circulation* 2003;107:2993-94.
10
11 41. Central Bureau of Statistics. *National Population and Housing Census 2011*. Kathmandu,
12 Nepal: Government of Nepal, National Planning Commission Secretariat; 2012.
13
14
15 42. Warning Signs of a Heart Attack. Available at:
16 <http://www.heart.org/HEARTORG/Conditions/HeartAttack/WarningSignsofaHeartAttack/Warni>
17 [ng-Signs-of-a-Heart-Attack_UCM_002039_Article.jsp](http://www.heart.org/HEARTORG/Conditions/HeartAttack/WarningSignsofaHeartAttack/Warni).
18
19
20 43. Ministry of Health and Population, Nepal. *Nepal non-communicable diseases risk factors*
21 *survey 2007*. Kathmandu, Nepal: Ministry of Health and Population, Nepal; 2008.
22
23
24 44. Vaidya A, Pathak RP, Pandey MR. Prevalence of hypertension in Nepalese community
25 triples in 25 years: a repeat cross-sectional study in rural Kathmandu. *Indian Heart J*
26 2012;64:128-31.
27
28
29 45. Brook RD, Franklin B, Cascio W, *et al.* Air pollution and cardiovascular disease A statement
30 for healthcare professionals from the expert panel on population and prevention science of the
31 American Heart Association. *Circulation* 2004;109:2655-71.
32
33
34 46. Nguyen TT, Liao Y, Gildengorin G, *et al.* Cardiovascular risk factors and knowledge of
35 symptoms among Vietnamese Americans. *Journal of General Internal Medicine* 2009;24:238-
36 43.
37
38
39 47. Redding CA, Rossi S, Rossi R, *et al.* Health behavior models. *The International Electronic*
40 *Journal of Health Education* 2000;3:180-93.
41
42
43 48. Avis NE, Smith KW, McKinlay JB. Accuracy of perceptions of heart attack risk: what
44 influences perceptions and can they be changed?. *Am J Public Health* 1989; 79:1608-12.
45
46 49. Sharma D, Bkc M, Rajbhandari S, *et al.* Study of prevalence, awareness, and control of
47 hypertension in a suburban area of Kathmandu, Nepal. *Indian Heart J* 2006;58:34-37.
48
49
50 50. Vaidya A, Pokharel P, Karki P, *et al.* Exploring the iceberg of hypertension: a community
51 based study in an eastern Nepal town. *Kathmandu Univ Med J* 2007;5:349-59.
52
53
54 51. Vanhecke TE, Miller WM, Franklin BA, *et al.* Awareness, knowledge, and perception of
55 heart disease among adolescents. *European Journal of Cardiovascular Prevention &*
56 *Rehabilitation* 2006;13:718-23.
57
58
59
60


- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
52. White CW, Albanese MA. Changes in cardiovascular health knowledge occurring from childhood to adulthood. A cross-sectional study. *Circulation* 1981;63:1110-15.
53. Yahya R, Muhamad R, Yusoff HM. Association between Knowledge, Attitude and Practice on Cardiovascular Disease among Women in Kelantan, Malaysia. *International Journal of Collaborative Research on Internal Medicine & Public Health* 2012;4:1507-23.
54. Muhamad R, Yahya R, Yusoff HM, *et al.* Knowledge, Attitude and Practice on Cardiovascular Disease among Women in North-Eastcoast Malaysia. *International Journal of Collaborative Research on Internal Medicine & Public Health* 2012;4:85-98.
55. Waller J, McCaffery K, Wardle J. Measuring cancer knowledge: comparing prompted and unprompted recall. *Br J Psychol* 2004;95:219-34.
56. Glass AL, Holyoak KJ. *Cognition* . Second ed. New York: McGraw-Hill; 1986.
57. Valente TW, Paredes P, Poppe PR. Matching the message to the process the relative ordering of knowledge, attitudes, and practices in behavior change research. *Human Communication Research* 1998;24:366-85.

1
2
3 **Figure legends:**
4

5
6 **Figure 1: Scattergrams showing correlation between median score percents for (a)**
7 **knowledge and attitude, (b) attitude and behavior, and (c) knowledge and behavior**
8 **(n=777).**
9


10
11 Scores were obtained from the prompted knowledge questions, attitudinal questions, and
12 practice/behavior responses. Median score percents were calculated and plotted. See Supplement
13 for greater detail about score derivation.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1 (a)




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Figure 1 (b)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Figure 1 (c)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Vaidya et al: Cardiovascular health knowledge, attitude, and practice behavior . . .

Supplement

Methods

Scoring system

Knowledge score

To calculate the knowledge score, we considered prompted responses to the five sections that explored that topic. The total score on knowledge combines the scores for cause of heart diseases, warning signs of heart attack, action in case of a heart attack, knowledge regarding heart-healthy foods, and knowledge regarding prevention of heart diseases. Maximum possible score was $11+15+7+10+10=53$.

Cause of heart diseases

Enumerators read aloud a list of conventional causes of heart disease. Score 1 was given if the respondent answered “yes.” No negative marking was given for “no” or “don’t know” responses.

Warning signs of heart attack

Nine common symptoms that are considered warning signs of a heart attack by the American Heart Association⁴³ and two incorrect signs (abdominal pain and leg pain) were mingled together. Respondents were asked if the symptoms were correct. If they answered “yes”, we allocated the scores as follows: 3 points for chest pain; 2 points each for breathlessness, sweating, and vomiting; 1 point each for arm, tooth, or jaw pain; 1 point for light-headedness and unconsciousness; 0 points for abdominal or leg pain. Additionally, we scored 1 point if respondents answered “no” for abdominal or leg pain.

Action in case of a heart attack

We asked respondents what actions they would take or not take in case of a heart attack. If the responses were “yes”, we allocated scores as follows: treat at home (0), immediately take the person to a hospital (2), take to a pharmacy/medicine shop (0), give home-made therapy (0), wait for appointment with a heart specialist (0), take the person to a traditional healer (0).

Heart-healthy foods

Vaidya et al: Cardiovascular health knowledge, attitude, and practice behavior . . .

We asked each respondent if they considered a particular food item from a pre-selected list to be heart-healthy. Responses were recorded as “yes,” “no,” or “don’t know.” We scored 1 point for each correct answer, depending on whether or not the particular food is actually heart-healthy (e.g., fruits) or not (e.g., fat).

Prevention of heart diseases

We asked each respondent about the various preventive actions that are considered good for improving heart health. Possible answers were “yes,” “no,” or “don’t know.” We scored 1 point for each “yes” answer.

Attitude

Each respondent completed a 14-item questionnaire related to her/his attitude towards heart health. Responses were recorded as strongly agree/disagree, somewhat agree/disagree, or don’t know/can’t say. Depending on whether the attitude was proper or not, we allotted scores ranging between 1 and 5 points. Similarly, participants were also asked for their opinions on various preventive aspects of heart. We recorded those responses on a scale from “very helpful” to “not at all helpful” as the extremes. Maximum possible score for the attitude section was $14 \times 5 = 70$.

Practice and behavior score

We asked each respondent if s/he undertook any particular action in the previous year that would increase her/his heart health and scored the answers accordingly. We also scored participants according to their risk factor status. Maximum score possible was 25.

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	Comment [AV1]: Yes Comment [AV2]: Yes
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Comment [AV3]: Yes
Objectives	3	State specific objectives, including any prespecified hypotheses	Comment [AV4]: Yes
Methods			
Study design	4	Present key elements of study design early in the paper	Comment [AV5]: Yes
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Comment [AV6]: Yes
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Comment [AV7]: Yes
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Comment [AV8]: Yes
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Comment [AV9]: Yes
Bias	9	Describe any efforts to address potential sources of bias	Comment [AV10]: Yes
Study size	10	Explain how the study size was arrived at	Comment [AV11]: Purposive
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Comment [AV12]: Yes
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses	Comment [AV13]: Yes Comment [AV14]: Yes Comment [AV15]: Not applicable
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	Comment [AV16]: Done, mentioned in the tables, etc as well Comment [AV17]: Yes Comment [AV18]: Not relevant
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest	Comment [AV19]: Yes Comment [AV20]: Yes
Outcome data	15*	Report numbers of outcome events or summary measures	Comment [AV21]: Yes
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Comment [AV22]: Yes Comment [AV23]: Yes
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion		
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

- Comment [AV24]: Yes
- Comment [AV25]: Yes
- Comment [AV26]: Yes
- Comment [AV27]: Yes
- Comment [AV28]: Yes

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.



Cardiovascular health knowledge, attitude, and practice/behavior in an urbanizing community of Nepal: a population-based cross-sectional study from Jhaukhel-Duwakot Health Demographic Surveillance Site

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2013-002976.R1
Article Type:	Research
Date Submitted by the Author:	07-Aug-2013
Complete List of Authors:	Vaidya, Abhinav; Kathmandu Medical College, Community Medicine Aryal, Umesh Raj; Kathmandu Medical College, Community Medicine Krettek, Alexandra; Nordic School of Public Health NHV,, ; Institute of Medicine, Sahlgrenska Academy at University of Gothenburg, Department of Internal Medicine and Clinical Nutrition
Primary Subject Heading:	Public health
Secondary Subject Heading:	Cardiovascular medicine
Keywords:	PUBLIC HEALTH, PREVENTIVE MEDICINE, EPIDEMIOLOGY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cardiovascular health knowledge, attitude, and practice/behavior in an urbanizing community of Nepal: a population-based cross-sectional study from Jhaukhel-Duwakot Health Demographic Surveillance Site

Abhinav Vaidya^{1,2}, Umesh Raj Aryal^{1,2}, and Alexandra Krettek^{2,3}

¹Department of Community Medicine, Kathmandu Medical College, Kathmandu, Nepal

²Nordic School of Public Health NHV, Gothenburg, Sweden, and

³Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden

Correspondence to

Abhinav Vaidya; Abhinav.Vaidya@nhv.se

Keywords: knowledge, attitude, and practice/behavior; Nepal; cardiovascular health; health literacy

Word count: 4,894 (excludes title page, abstract, references, tables, and figures, per author guidelines)

ABSTRACT

Objectives: This study determined the knowledge, attitude, and practice/behavior of cardiovascular health in residents of a semi-urban community of Nepal.

Design: To increase understanding of knowledge, attitude, and practice/behavior toward cardiovascular health, we conducted in-home interviews using a questionnaire based on WHO STEPS and other resources, scoring all responses. We also recorded blood pressure and took anthropometric measurements.

Setting: Our study was conducted as part of the HARDIC project in the Jhaukhel-Duwakot Health Demographic Surveillance Site in two urbanizing villages near Kathmandu.

Participants: The study population included 777 respondents from six randomly selected clusters in both villages.

Results: Seventy percent of all participants were female and 26.9 % lacked formal education. The burden of cardiovascular risk factors was high; 20.1 % were current smokers, 43.3 % exhibited low physical activity, and 21.6 % were hypertensive. Participants showed only poor knowledge of heart disease causes; 29.7 % identified hypertension and 11% overweight and physical acitivity as causes, wheras only 2.2 % identified high blood sugar as causative. Around 60 % of respondents did not know any heart attack symptoms compared with 20% who knew 2–4 symptoms. Median % scores for knowledge, attitude, and practice/behavior were 79.3, 74.3, and 48.0, respectively. Nearly 44% respondents had insufficient knowledge and less than 20% had highly satisfactory knowledge. Among those with highly satisfactory knowledge, only 14.7% had highly satisfactory attitude and 19.5% and 13.9% had satisfactory and highly satisfactory practices, respectively.

Conclusions: Our study demonstrates a gap between cardiovascular health knowledge, attitude, and practice/behavior in a semi-urban community in a low-income nation, even among those already affected by CVD.

ARTICLE SUMMARY

Article focus

- We aimed to gain a deeper understanding of knowledge, attitude, and behavior/practice regarding cardiovascular health in a semi-urban community of Nepal, a low-income South Asian country.

Key messages

- We found a high burden of low physical activity and overweight/obesity; cardiovascular risk factors in our study site exceeded the national average.
- Low level of knowledge and attitude towards risk perception combined with poor relation between knowledge, attitude, and behavior scores imply that health promotion strategies that merely increase knowledge of cardiovascular health may be insufficient, thus emphasizing the need for a more holistic approach to disease prevention.

Strengths and limitations of the study

- This study is the first to extensively explore knowledge, attitude, and behavior/practice regarding cardiovascular health in Nepal and was conducted in a health demographic surveillance site which provides the logistic advantage of further follow-up and intervention.
- Using both prompted and unprompted questions in the same study illustrates the variation that may be obtained by these approaches.
- Unintentional oversampling of women respondents yielded unsegregated gender-wise results that are more likely to represent knowledge, attitude, and practice/behavior in women.

INTRODUCTION

Noncommunicable diseases (NCDs) pose increasingly important public health problems in low- and middle-income countries (LMICs). In 2008, 80% of global NCD deaths occurred in LMICs,¹ and NCDs recently accounted for 7.9 million of 14.5 million deaths (54%) in Southeast Asia² and were responsible for two of every three deaths (34.5 million) worldwide in 2010.³ Poverty, illiteracy, poor health infrastructure, and demographic transition, including increasing life expectancy, are major contributors to rising burden of NCDs in LMICs. Other postulated causes include poor fetal and childhood nutrition.^{4,5} A similar pattern is evident in South Asian countries like Nepal, where NCDs currently occur more frequently than communicable diseases and behavioral risk factors such as tobacco smoking are high.⁶

Globally, cardiovascular diseases (CVDs) are responsible for one in four deaths (12.9 million).³ Eighty percent of CVDs result from tobacco smoking, unhealthy diet, physical inactivity, and harmful use of alcohol.⁷ The risk of all-cause mortality among individuals who score poorly in behavioral factors is four fold higher than those who score best.⁸ In high-income countries, declining CVD mortality is largely attributed to the reduction of behavioral risk factors.⁹ Underlying socioeconomic parameters and societal influences affect exposure and vulnerability to CVD risk factors. Therefore, surveillance of NCD risk factors should include social determinants of cardiovascular health.^{1,10} Currently, understanding the individual and social determinants of cardiovascular health behaviors is among the top 20 priority areas for NCD research in LMICs.¹¹

As a precursor of favorable health behavior, experts have advocated health literacy since community-based interventions began exploring the potential impact of behavioral changes on CVD.^{1,12} The World Health Organization (WHO) promotes public awareness of diet and physical activity as a “best buy” to improve cardiovascular behavior.⁹ However, achieving desired health behavior requires deeper understanding of the underlying social context of disease.^{13,14} Hence, studies on knowledge, attitude, and practice (KAP) increase our understanding of cardiovascular health literacy and community behavior.¹⁵ Currently, such studies have been limited in South Asia.

1
2
3 Earlier studies report low level of knowledge about heart attack¹⁶ and diabetes symptoms in
4 Nepal.¹⁷ Nepal's high prevalence of underlying behavioral risk factors, particularly tobacco and
5 alcohol consumption, and poor cardiovascular health knowledge is a matter of concern.¹⁸
6
7 Nationwide in Nepal, the age group 25+ exhibits fasting blood glucose of 5.5 mmol/L for males
8 and 5.4 mmol/L for females, systolic blood pressure of 125.9 mmHg and 126.6 mmHg for males
9 and females respectively, and total cholesterol of 4.1 mmol/L (males) and 4.3 mmol/L
10 (females).¹⁹
11
12
13
14
15

16
17 Currently, attempts to improve cardiovascular health knowledge and behavior in Nepal focus
18 mainly on campaigns that transmit information but pay little attention to the population's actual
19 health literacy. We therefore undertook a community-based KAP study in the Jhaukhel-Duwakot
20 Health-Demographic Surveillance Site (JD-HDSS) outside Kathmandu where NCDs,
21 particularly CVDs, are the predominant cause of morbidity and mortality.²⁰ Additionally, our
22 pilot study in the Duwakot population demonstrated poor cardiovascular health knowledge.²¹ Our
23 study is part of the Heart-Health Associated Research and Dissemination in the Community
24 (HARDIC) project, which is the cardiovascular health component of JD-HDSS. In Nepalese,
25 hardic means "heartily" or "from the heart". This paper reports the baseline results from
26 HARDIC and is the first to conduct a detailed analysis of cardiovascular health knowledge,
27 attitude, and practice/behavior in an urbanizing community of Nepal.
28
29
30
31
32
33
34
35
36
37
38
39

40 **METHODS**

41 **Ethical considerations**

42
43 The Nepal Health Research Council and the Institutional Review Board of Kathmandu Medical
44 College approved this study. Following a pattern established at the outset of JD-HDSS, we
45 consulted local leaders and sought their informal permission for HDSS activities. In addition, we
46 periodically consulted with health and administrative authorities in both villages. All participants
47 gave informed consent, and privacy was maintained during the interview process. To maintain
48 confidentiality, all data were secured in the HDSS office. Computer data were accessible only to
49 the research team. Although respondents received no monetary incentives, we provided a referral
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that included discounted consultation fees at Kathmandu Medical College or Nepal Medical
4
5 College for self-reported illness and newly diagnosed hypertension.
6
7

8 **Study site**

9
10 We conducted our descriptive, community-based, cross-sectional study between September and
11
12 November 2011 in the JD-HDSS of Bhaktapur district in the Kathmandu Valley, 13 kilometers
13
14 from Kathmandu. An earlier report described the establishment of the surveillance site.²⁰
15

16 **Study population and sample size**

17
18 This study is the first step in the larger HARDIC project that will be conducted in the whole JD-
19
20 HDSS. Duwakot and Jhaukhel each contain nine administrative wards and we randomly selected
21
22 three wards from each village. Based on the JD-HDSS 2010 baseline census,²⁰ we compiled a
23
24 complete list of households containing adults aged 25-59 years in the selected wards. Next, we
25
26 selected one adult, of either sex, from all interview households and surveyed 789 of 840
27
28 households (non-response rate = 6.07%). We applied the Kish technique to select the
29
30 respondent.²²
31

32 **Study tools**

33
34 We structured the basic questionnaire and tailored it for site specific responses in Nepal using the
35
36 WHO-NCD STEPS (STEP 1 and 2 questionnaires) for demographic information, behavioral and
37
38 physical measurements²³ and other sources for addressing knowledge, attitude and
39
40 practice/behavior issues.^{15, 24-39} Responses to questions on cardiovascular health knowledge,
41
42 attitude and practice/behavior were given scores in order to compare and correlate the
43
44 components. A detailed description of the scoring system is given as an appendix to this article.
45

46 **Interviewers**

47
48 Twelve local female enumerators (two per ward) conducted the interviews. All enumerators had
49
50 completed at least grade 10 in school. Supervisors included four public health graduates; a field
51
52 coordinator with Master and PhD degrees in public health; and two PhD students/authors (AV
53
54 and URA).
55

56 **Data Collection**

Interview process

The interviews were conducted at the participants' homes. The initial part of the interview consisted of questions related to demographic profile. Questions then followed relating to knowledge of cardiovascular health (e.g., causes and risk factors, warning signs, and action for suspected heart attack); heart-healthy diet; and disease prevention. The first two areas included unprompted (spontaneous, open-ended) and prompted (clues provided, closed-ended) responses, and the latter two areas included only prompted responses. We asked respondents about their attitude toward different aspects of cardiovascular health: did they perceive themselves at risk for CVD? Did they think behavior change would reduce their risk of developing CVD? The responses were recorded in Likert-scale (strongly agree, somewhat agree, don't know, somewhat disagree, strongly agree).

Behavioral factors

Behavior-related questions were based on Step 1 of the WHO STEPwise approach to surveillance (STEPS), which uses a questionnaire to collect data on risk factors.²³ Our questions probed current and past smoking behavior; smokeless tobacco; alcohol consumption in the past 30 days (current drinker), and drinking frequency among current drinkers. The physical activity section asked respondents to report the number of days and amount of time they participated in vigorous and moderate activities at work, leisure, and travel. Additional practice and behavioral questions explored whether participants had engaged in any heart-friendly behavior in the previous year and for what reason.

Physical measurements

Enumerators recorded anthropometric measurements (i.e., weight, height, and waist and hip circumferences) according to STEPs guidelines.²³ Weight was measured to one decimal place using a Microlife BR-9201 Weighing Scale (Microlife AG Swiss Corporation, Widnau, Switzerland). Height and waist and hip circumferences were measured using non-stretchable tapes (Jonson Tapes Ltd, New Delhi, India). Blood pressure was determined digitally by Microlife BP 3AP1-3E (Microlife AG Swiss Corporation, Widnau, Switzerland). Enumerators recorded three readings for each respondent, taken at 5-minute intervals. We averaged the readings to attain the final blood pressure measurements.

Diagnostic criteria for sub-classification

Respondents with known history of hypertension (diagnosed cases) and those diagnosed during the study according to Joint National Committee -VII⁴⁰ were termed as having “hypertension”.

Data management

Supervisors checked all completed questionnaires and, if necessary, asked enumerators to revisit households. After omitting incomplete questionnaires, we included 777 individuals in the analysis. Except for missing recordings for height, weight, or blood pressure, data reported in our KAP-study covers all 777 respondents. We outsourced data entry to a team of data entry operators, public health graduates by qualification. Data was coded and entered in Epidata version 2.1.

Statistical analyses

Data was analyzed with SPSS version 17.0 (IBM, Armonk, New York, USA) and STATA version 10.0 (StataCorp, Texas, USA). After calculating the Metabolic Equivalent of Task (MET) minutes/week, we grouped participants as highly active, moderately active, or inactive (low physical activity) in accord with the STEPS manual.

For scoring knowledge, attitude, and practice/behavior questions and to compare and correlate the three components, we scored responses regarding KAP/behavior for cardiovascular health. The maximum possible score for KAP/behavior was 53, 56 and 25, respectively. Further, total knowledge score was classified into the following five categories based on the quintile scores: ≤ 20 % ‘highly insufficient’; 21–40 % ‘insufficient’; 41–60 % ‘sufficient’; 61–80 % ‘satisfactory’; > 80 % ‘highly satisfactory’.⁴¹

Categorical data is presented as number, percentages, and continuous data as median and inter-quartile range (IQR). KAP percentage scores are presented as median. We used Chi-square test to compare proportions among demographic variables and Mann-Whitney U test to compare scores from males and females. $P < 0.05$ was considered statistically significant.

Cronbach’s alpha, which measures question reliability, for the overall knowledge and attitude was 0.941 (range: 0.775–0.913) and 0.824 (range: 0.791–0.822), respectively. Cronbach’s alpha

for practice and behavior ranged from low (0.257) for physical activity questions to high (0.929) for blood pressure measurements.

RESULTS

Demographic characteristics and cardiovascular risk factors

Table 1 describes the sex-wise demographic characteristics of the respondents. Seventy percent of all participants were female, and 25% lacked formal education. The median age (IQR) for male and female respondents was 40 (33-49) years and 38 (32-46) years, respectively. Two third of the female respondents were housewives and 3 out of 5 male were either serviceholders or self-employed. We have shown previously that 59.2% belonged to the upper-lower class.²⁰

Table 1: Demographic characteristics of the study population.

Demographic variables	Male (n=229) Number (%)	Female (n=548) Number (%)	Total (n=777) Number (%)
Age (years)			
25–34	69 (26.8)	188 (73.2)	257 (100)
35–44	78 (27.9)	202 (72.1)	280 (100)
45–59	82 (34.2)	158 (65.8)	240 (100)
Caste/Ethnicity			
Brahmin	85 (28.9)	209 (71.1)	294 (100)
Chhetri	55 (28.4)	139 (71.6)	194 (100)
Newar	67 (34.5)	127 (65.5)	194 (100)
Others*	22 (23.2)	73 (76.8)	95 (100)
Education			
No formal schooling	27 (12.9)	182 (87.1)	209 (100)
Primary school (Grade:1-5)	74 (30.1)	172 (69.9)	246 (100)
Secondary school (Grades: 6-10)	74 (37.4)	124 (62.6)	198 (100)
High secondary and above (Grade:11 and above)	54 (43.5)	70 (56.5)	124 (100)

Major Occupation[§]			
Service (government and private)	70 (60.3)	46 (39.7)	116 (100)
Self-employed	72 (66.7)	36 (33.3)	108 (100)
Agriculture	48 (38.1)	78 (61.9)	126 (100)
Housewife**	N/A	369 (100)	369 (100)
Others [†]	39 (67.2)	19 (32.8)	58 (100)

* Other caste/ethnicity includes Rai, Magar, Tamang, Dalit, Gurung, Mandal, Chaudary, Pariayar, Purkutti. [§]The definition of occupation was adopted from the Nepal Non-Communicable Disease Survey, 2007 and modified to the local context. ⁴² ** A woman involved in her own household activities like cooking, washing, cleaning etc but does not earn money. [†] Other occupation includes unemployed individuals, able or unable to work, retired, and non-paid workers.

Figure 1 shows the sex-wise prevalence of cardiovascular risk factors in the study population. Overall, three in ten respondents was a past or current smoker and one in five respondents had consumed at least one alcoholic beverage in the previous month. More than 40% of the respondents reported low physical activity. Prevalence of known and newly diagnosed hypertension was 21.6%. As measured by BMI, the prevalence of overweight and obesity was 29.7% and 9.4 %, respectively. Abdominal obesity (i.e., increased waist circumference and waist-hip ratio) was 46.3% and 56.3%, respectively. There was statistically significant difference in the prevalence rate among males and females in smoking, smokeless tobacco use, alcohol intake, overweight/obesity and increased waist circumference.

m=number of male, f=number of female, # p<0.001, ## p<0.01 (chi-square tests). Risk factor categories based on the WHO-NCD Risk Factor STEPS Survey manual.²³ Current smokers included those who responded “yes” to “Do you smoke?” Past smokers included those who replied “yes” to “Did you ever smoke in the past?”. “Ever drinkers” included respondents who had “consumed a drink that contained alcohol ever in lifetime,” and current drinkers included respondents who had “consumed alcohol within the previous month.” Percentage of drinking frequency represents current drinkers who consumed at least one standard drink per occasion. Blood pressure data exclude respondents who did not submit to all three readings. Likewise, data for body mass index, waist circumference, and waist-hip ratio exclude respondents whose weight, height, and waist and/or hip measurements were not taken. Increased waist circumference includes waist measurements of ≥ 80 cm (females) and ≥ 90 cm (males); increased waist-hip ratio is ≥ 0.85 (females) and ≥ 0.90 (males).

Knowledge towards cardiovascular disease

Knowledge regarding causes and warning signs of heart disease

Table 2 shows unprompted responses for the perceived causes of heart diseases. The responses ranged from 1.0% for high blood sugar to 29.1% for smoking. The median percentage scores for males and females were 13.8 and 10.4 %, respectively (p<0.001, Mann-Whitney U test).

Younger respondents (<35 years) knew more than older participants (median 13.7% vs. 10.3%, $p=0.01$). Respondents belonging to Brahmin, Newer and Chhetri ethnicities had less knowledge than those in the other caste/ethnicity group (median 13.8% vs. 6.9%, $p=0.001$) (Table 1). The respondents with high school or more education and working in government or non-government offices knew more than those with less education and in other employment situations (median 17.2% vs. 10.34%, $p=0.001$).

Table 2: Percentage of respondents citing various causes of cardiovascular disease and warning signs of heart attack.

Causes of Heart Attack	Male (n=229) Number (%)	Female (n=548) Number (%)	Total (n=777) Number (%)	P-value*
Hypertension	43 (18.8)	60 (10.9)	103 (13.3)	<0.01
High blood sugar	3 (1.3)	5 (0.9)	8 (1.0)	0.06
High cholesterol	47 (20.5)	74 (13.5)	121 (15.6)	0.01
Physical inactivity	16 (7.0)	22 (4.0)	38 (4.9)	0.08
Overweight	13 (5.7)	29 (5.3)	42 (5.4)	0.82
Cigarette/Bidi Smoking	80 (34.9)	147 (26.8)	227 (29.1)	0.02
Smokeless tobacco	19 (8.3)	26 (4.7)	45 (5.8)	0.06
Excessive alcohol	72 (31.4)	131 (23.9)	203 (26.1)	0.03
Excess stress	37 (16.2)	74 (13.5)	111 (14.3)	0.34
Warning signs	Male (n=93)[§] Number (%)	Female (n=222)[§] Number (%)	Total (n=315)[§] Number (%)	P value*
Difficulty in breathing	17 (11.1)	41 (15.1)	58 (13.6)	0.25
Pain in the chest	30 (19.6)	57 (21.0)	87 (20.5)	0.74
Pain or numbness in the arms	0 (0)	3 (1.1)	3 (0.7)	-
Pain in the teeth or jaw	0 (0)	1 (0.4)	1 (0.2)	-
Excessive sweating	15 (9.8)	30 (11.0)	45 (10.6)	0.69
Vomiting tendency	1 (0.7)	6 (2.2)	7 (1.6)	0.65
Dizziness or light headedness	24 (15.7)	40 (14.7)	64 (15.1)	0.78
Loss of consciousness	55 (35.9)	86 (31.6)	141 (33.2)	0.36

Numbers based on participants' unprompted responses. Only established CVD risk factors included and responses citing other non-established causes (e.g., food hygiene, air pollution, etc.) excluded. *Obtained from chi square test.

[§]Respondents who did not know any warning sign were excluded from analysis.

Table 2 also describes the warning signs of heart attack as perceived by the respondents. Around 60% of respondents did not know any of the heart attack signs. Among those who were familiar with those signs, nearly 50% of respondents knew only one sign, 37% knew any two signs, 11.42% knew any three signs and 2.53% knew four signs. The proportions of each warning sign told by the male and female respondents were not statistically different ($p>0.05$). The proportions of knowledge about warning signs were not statistically different in terms of age ($\chi^2=3.39$, $p=0.49$), caste/ethnicity ($\chi^2=6.25$, $p=0.4$), and education ($\chi^2=5.50$, $p=0.53$). Importantly, only 20% of respondents identified chest pain—the most prominent and important indicator of heart attack—as a symptom of heart attack.

Knowledge on action in case of a heart attack

When asked about what action the respondents would take for a suspected heart attack, more than 20% gave no response and 75% (81.2% male and 73% female, $\chi^2=4.05$, $p=0.04$) said they would immediately take the affected person to hospital. About 3% said they would try home therapy, consult a traditional healer, or stay home and wait for the symptoms to go away.

Knowledge on healthy foods for heart diseases

When asked to identify a particular heart-healthy food, most respondents identified green leafy vegetables (94.6%) and fruit (92.1%). Furthermore, 22% thought salty food like chips and around 7% thought fried food or saturated fat (e.g., ghee) are healthy foods.

Attitudes and practice towards cardiovascular disease

Attitude toward heart health in an urbanizing community

Table 3 describes opinions reflecting attitudes towards various aspects on heart diseases. Regarding lifestyle and preventive measures, 25% of males and 30.0% females perceived themselves to be at risk of CVD. More than half of the respondents did not want to change their current lifestyle (64.6% males vs. 55.4% females, $p<0.01$). Nearly 23% said they eat more and would like to cut down; approximately 82% considered themselves to be overweight - the perceived proportion being significantly higher among males ($p<0.05$); nearly 32% believe that they do not have time to take care themselves; and three fifths did not think that changing their

behavior would lower their cardiovascular risk. Further, 25.3% of males and 27.2% of females thought that God or a higher power determines their health.

Table 3: Responses to attitude-related questions on cardiovascular disease.

SN	Items	Male	Female	Total	P-value*
		Number (%)	Number (%)	Number (%)	
Attitude items on heart health					
1	I don't perceive myself to be at risk for heart disease (m=225/f=540) [§]	58 (25.8)	167 (30.9)	225 (29.0)	0.15
2	I don't want to change my lifestyle (activity, smoking, alcohol) (m=229/f=548)	148 (64.6)	298 (54.4)	446 (57.4)	p<0.01
3	I feel the amount of food I eat is too much, and I would like to cut down (m=229/f=543) [§]	35 (15.2)	140 (25.5)	175 (22.5)	p<0.01
4	I feel my present weight is too high for my health (m=229/f=548)	37 (14.9)	130 (23.7)	167 (21.5)	0.02
5	I don't think changing my behavior will reduce my risk of developing heart disease (m=229/f=522) [§]	145 (63.3)	312 (56.9)	457 (58.8)	0.23
6	I don't have time to take care of myself (m=229/f=548)	77 (33.6)	168 (30.7)	245 (31.5)	0.42
7	God or a higher power ultimately determines my health (m=229/f=548)	58 (25.3)	149 (27.2)	207 (26.6)	0.59
Opinions on various preventive aspects of heart health that are helpful					
8	Access to more fruits, vegetables, and other healthy foods (m=229/f=548)	226 (98.6)	533 (99.1)	759 (99.1)	- [†]
9	Greater access to indoor and outdoor public recreational facilities (m=229/f=548)	200 (87.3)	492 (89.7)	692 (89.1)	0.31
10	Smoking bans (m=218/f=535) [§]	209 (96.1)	515 (96.2)	724 (96.1)	0.89
11	More awareness programs for healthy heart (m=226/f=536) [§]	214 (96.1)	483 (90.1)	697 (98.0)	0.38
12	Availability of health facilities to detect and treat heart diseases (m=226/f=531) [§]	217 (96.1)	524 (98.6)	741 (97.9)	0.02

13	Local volunteers that shall try to change any adverse health behavior (m=226/f=530) [§]	186 (82.1)	522 (98.5)	708 (93.6)	p<0.01
----	--	------------	------------	------------	--------

m=number of males, f=number of females. Questions based on 5-scale Likert questions (strongly disagree, somewhat disagree, I do not know, somewhat agree and strongly agree (question 1-7)/very helpful; somewhat helpful; I do not know; not very helpful; not at all helpful (question 8-13). We classified five options into three categories as 'agree/helpful', 'disagree/not helpful' and 'I do not know'. Only 'strongly agree' or 'very helpful' and 'agreed/somewhat helpful' answers were merged and presented in the table and 'I do not know/cannot say' were excluded from analysis. *Obtained from chi square test. [§]'I do not know/cannot say' answers were excluded from analysis. [†]Cannot be computed due to expected frequency less than 5.

Table 3 also describes the respondents' opinions on various preventive aspects of heart health that would be helpful. Almost all respondents believed that increased access to fruits, vegetables, and other healthy foods would be helpful. Nearly 9 in 10 respondents thought that greater access to indoor and outdoor public recreational facilities would be helpful. Four percent of respondents did not believe smoking bans would be helpful to reduce heart diseases. Eight in 10 male respondents believed that local volunteers should try to change any adverse health behaviors while nearly 9 in 10 females believed the same (p<0.01).

Practice towards heart diseases

Half of all the respondents had their blood pressure measured and 10% reported getting a diagnostic test for CVD in the previous year. About 60% attempted to reduce their consumption of unhealthy foods and 25% tried to lose weight or increase physical activity. Half of the smokers said they had attempted to quit smoking.

When enumerators asked the respondents who reported one or more positive health behaviors the main reason for trying to change their lifestyle or practice/behavior, the reasons given were: they wanted to feel better (70%); had seen, read, or heard information (65.4%); did it for their family (45.9%); or were encouraged to do so by a healthcare professional (41.3%), family member (39.0%), or friend (31.8%). Other reasons for changing lifestyle or practice/behavior included wanting to avoid taking medications (20.6%); a family member, relative, or friend who developed CVD, became ill, or died (16.5%); and developing possible cardiovascular symptoms (14.8%).

Comparison between level of knowledge, attitude and behavior/practice

Overall, the average prompted knowledge of heart diseases (median 79.3%) was 11.5% lower than average attitudes (median 67.8%) and nearly 31.1% higher than average practices (median 48.2%). Next, average level of attitudes score was nearly 19.6% higher than practices. Similarly, average spontaneous knowledge of heart diseases (median 10.3%) was 57.5% lower than average attitudes (median 67.8%) and nearly 40% lower than average practice (median 48.18%).

Table 4 describes level of KAP based on quintile scores. Regarding knowledge, 20.4 % of the respondents had highly insufficient knowledge, 23.2% had insufficient knowledge, 22.6% had sufficient knowledge, 16.5% had satisfactory knowledge and 17.2% had highly satisfactory knowledge. Similarly, attitude levels were as follows: highly insufficient 20.6%, insufficient 26.9%; sufficient 15.4%; satisfactory 11.2% and highly satisfactory 19.8%. Regarding the behaviour/practices, it was highly insufficient in 12.1% of the respondents, insufficient in 24.3%; sufficient in 22.7%; satisfactory in 27.5% and highly satisfactory in 13.4%. Among those with highly insufficient knowledge, 20.3% of them had highly satisfactory attitudes and 6.9% had highly satisfactory practice. Conversely, among the respondents with highly satisfactory knowledge, 14.7% had highly satisfactory attitude and 19.5% and 13.9% had satisfactory and highly satisfactory practices respectively. Likewise, one in ten respondents with satisfactory knowledge had satisfactory attitude while two in ten had highly satisfactory attitude. Nearly four out of ten had satisfactory/highly satisfactory behavior. As the level of knowledge increased, the highly satisfactory practice also increased from 6.9% to 13.4%. However, when the level of knowledge increased, the highly satisfactory attitude decreased from 20.3% to 14.7%.

Table 4: Percentage distribution of level of knowledge, attitude and behavior/practice regarding cardiovascular health.

Knowledge (n=777)	Attitude (n=777)				
	HI (n = 160)	I (n = 209)	Suff (n=120)	Sat (n=134)	HS (n=154)
HI (n = 159)	26.0	27.6	13.8	12.5	20.3
I (n=180)	17.9	28.4	19.1	12.5	22.2
Suff (n=176)	17.9	29.8	17.2	11.9	26.2
Sat (n=128)	20.8	33.7	15.8	6.9	23.8
HS	33.7	24.2	12.6	12.6	14.7

(n=134)					
Knowledge (n=777)	Behavior (n=777)				
	HI (n = 94)	I (n=189)	Suff (n=176)	Sat (n=214)	HS (n=104)
HI (n = 159)	16.4	29.6	21.4	25.8	6.9
I (n=180)	10.4	26.1	19.4	30.6	13.3
Suff (n=176)	11.4	20.5	20.5	31.2	16.5
Sat (n=128)	10.9	18.8	33.6	19.5	17.2
HS (n=134)	11.2	26.1	20.9	28.5	13.4
Attitude (n=777)	Behavior (n=777)				
	HI	I	Suff	Sat	HS
HI (n = 132)	15.0	23.8	20.0	28.1	13.1
I (n=101)	12.0	22.0	18.7	30.1	17.2
Suff (n=256)	10.0	24.2	25.0	28.3	12.5
Sat (n=134)	13.8	33.3	17.2	29.9	5.7
HS (n=154)	10.4	20.8	33.8	24.0	11.0

HI: Highly insufficient; I: Insufficient; Suff: Sufficient; Sat: Satisfactory; HS: Highly Satisfactory.

The total score was 53, 64 and 25 for KAP, respectively. Then, the quintile scores were obtained. The quintile score (1st-4th) was for prompted knowledge: 34, 40, 43, 45; for attitudes: 33, 37, 39, 42; for behavior: 9, 11, 13, 15. Next, the levels of survey respondents on KAP were classified into the following five subdivisions based on quintile score: $\leq 20\%$ 'highly insufficient'; 21–40 % 'insufficient'; 41–60 % 'sufficient'; 61–80 % 'satisfactory'; $> 80\%$ 'highly satisfactory'.⁴¹ Median percentage scores for knowledge, attitude and practice were 79.3, 74.2 and 48 for all respondents.

Regarding attitude, among those with the level of attitudes of highly insufficient, 41.2% had satisfactory/highly satisfactory behaviour/practice. When the level of attitude was highly satisfactory, 35% had satisfactory/highly satisfactory practice. Similarly, when the attitude was sufficient, 25% had insufficient practice and nearly 41% had satisfactory practice. When the level of attitude increased, the proportion of those with highly satisfactory practice remained unchanged.

1
2
3 A comparison of male and female respondents showed no difference in level of knowledge
4 ($\chi^2=4.4$, $p=0.4$) attitudes ($\chi^2=3.6$, $p=0.5$) and practice ($\chi^2=2.1$, $p=0.7$). However, there were
5 significant differences in the level of knowledge according to age ($\chi^2=18.1$, $p=0.02$),
6 caste/ethnicity ($\chi^2=43.8$, $p<0.001$) and education level ($\chi^2=48.7$, $p<0.001$). In contrast, the level
7 of attitude was not statistically significant for these variables ($p>0.05$). On the other hand, the
8 level of practice was also statistically different according to age ($\chi^2=18.7$, $p=0.01$),
9 caste/ethnicity ($\chi^2=21.1$; $p=0.004$) and education status ($\chi^2=45.7$, $p<0.001$).
10
11

12
13
14
15
16 Respondents <35 years of age were more likely to have highly satisfactory knowledge (37.3%
17 vs. 29.2%) and practice (38.5% vs. 19.2%) than those >45 years. Brahmins (30.5%) were more
18 likely to have highly satisfactory knowledge than Newars (27.6%), Chhetris (20.4%), and others
19 (21.5%). Regarding highly satisfactory practice, the proportion was highest among Brahmins
20 (41.3%), followed by Chhetris (30.8%), Newars (17.3%) and others (10.6%). Those who had no
21 formal schooling exhibited higher proportion of highly satisfactory knowledge than those who
22 had high school or more education (28.3% vs. 24.6%, but highly satisfactory practice is low
23 among them (12.5% vs. 28.8%, data not shown).
24
25
26
27
28
29
30
31
32
33

34 DISCUSSION

35
36
37 The present study is the first to conduct a detailed analysis of the relationship between
38 KAP/health literacy and cardiovascular health in a semi-urban setting near Kathmandu.
39
40

41 **Cardiovascular risk factors are abundant in the study population**

42
43
44 Our respondents showed a high burden of behavioral and biological risk factors. While the
45 prevalence of tobacco smoking was similar to the national average, the alcohol consumption was
46 lower.⁴² National figures on tobacco smoking and alcohol consumption for Bangladesh, Nepal,
47 India and Pakistan show that current smoking of any tobacco product (age-standardized rate,
48 both sexes) is 24%, 32%, 15% and 20%, respectively.¹⁹ The total adult (recorded and
49 unrecorded) per capita consumption of pure alcohol according to WHO-available projected
50 estimates for 2008 indicates that consumption in Bangladesh, Nepal, India and Pakistan is 0.17,
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2.42, 2.69, 0.05 litres of pure alcohol, respectively, over a calendar year.¹⁹ Numbers have been increasing in these countries over the last years.¹⁹

In accord with our earlier review on obesity in Nepal, the effect of urbanization is evident in low physical activity and overweight/obesity that exceeds the national average.^{21,42} The prevalence of hypertension in our study population is similar to the national average (22%) but less by about one third than a study in Bhadrabas, a similar semi-urban Kathmandu community,⁴³ most probably due to different age groups. The 2008 national reported age-standardized level of hypertension in Nepal is similar to levels in Bangladesh but lower than in Pakistan (38.6% and 39.5%, respectively) and higher than in India (35.2%).⁴⁴ Self-reported prevalence of diabetes mellitus (3.9%) confirms the HDSS baseline figure reported earlier.²⁰

Knowledge of cardiovascular health is limited

Our study population showed low overall knowledge of the causes of CVD. Unprompted responses suggested that respondents identified behavioral factors (e.g., smoking) more frequently than physiological causes (e.g., hypertension). Other studies have reported similar findings for unprompted³⁸ and prompted responses.³³ Conversely, unhealthy diet, obesity, and family history were common responses in Taiwan.³¹ In Vietnam, knowledge on the negative influence of tobacco smoking on stroke and heart attack is lower than general knowledge regarding health risks of active smoking and exposure to second hand smoke.⁴⁵

Many of our respondents also attributed CVD to environmental factors such as food hygiene and air pollution. Indeed, acute inflammatory reactions to air pollutants accelerate coronary atherosclerosis and trigger arrhythmia or myocardial infarction.⁴⁶ Unlike the Strong Heart Study³⁹ in American populations, our results did not exhibit a higher probability of knowledge of a risk factor in persons who have that risk factor, demonstrating poor cardiovascular health literacy even among those already affected.

Similar to Pakistan,¹⁵ a large majority (60%) of our respondents could not identify even one heart attack symptom. However, similar to an earlier study in Nepal,¹⁶ 6%–22% of individuals in our demographic subsets recognized chest pain as a symptom, which is less compared to 60% of Vietnamese Americans who spontaneously mentioned chest pain.⁴⁷ Importantly, two thirds of

1
2
3 heart attack patients delayed their arrival at a Karachi hospital because they did not recognize
4 heart attack symptoms.¹⁵ Similar to a previous study,¹⁸ three quarters of our respondents believed
5 they would immediately take a suspected heart attack patient to a hospital but one fifth did not
6 answer the question or opted instead for home therapy or traditional healing. Combined with our
7 earlier finding of not knowing heart attack symptoms, this lack of knowledge about appropriate
8 action underscores the need to educate the Nepalese population about this issue. In contrast, 85%
9 of Vietnamese Americans knew that they should immediately call 911 if they suspect a heart
10 attack.⁴⁵ Despite inadequate knowledge about the causes of CVD, our respondents were much
11 better informed about prevention measures. This finding concurs with the perception level of
12 American women,³⁵ probably because enumerators asked prompted questions in this section of
13 the questionnaire. In Taiwan, only 25% of respondents thought that CVD is preventable and 50%
14 were “not sure.”³¹

25 **Inappropriate attitude toward cardiovascular health**

26
27
28 In contrast to their opinion that behavioral modifications can prevent CVD, most of our
29 respondents did not consider themselves at risk and did not want to alter their lifestyle. This may
30 be explained by the Health Belief Model.⁴⁸ Others have described a similarly optimistic bias.⁴⁹
31 Risk indicators (e.g., number of cigarettes smoked; death of a parent due to CVD; and self-
32 assessment of blood pressure, weight, and cholesterol levels) account for a 24% variance in
33 perceived risk.⁴⁹ The likelihood of perceiving oneself at higher risk increases when the presence
34 of a risk factor is known.³¹ However, respondents often underestimate their cardiovascular risk.⁴⁹

41 **Attempts for better cardiovascular health practice/behavior**

42
43
44 Compared to Nepal’s national average (69%), only half of our respondents underwent blood
45 pressure measurement in the previous year.⁴² Ninety percent of the known hypertensive
46 population had their blood pressure under control, exceeding the 6% and 9.5% control rate
47 reported in two other suburban Kathmandu studies^{43,50} and an urban study in Eastern Nepal
48 (58.5%).⁵¹ Half of the smokers reported here that they had tried to quit smoking. Our
49 questionnaire did not explore whether they had actually quit and did not identify the barriers
50 against quitting, information that might be important for public health. The theory of reasoned
51 action⁴⁸ argues that intention is a critical component of healthy behavior. Although confirming
52
53
54
55
56
57
58
59
60

1
2
3 the genuineness of intention is difficult, such intention may provide an opportunity for public
4 health intervention. Importantly, we identified our respondents' reasons for attempting positive
5 health behavior. Many constructs of the Health Belief Model,⁴⁸ such as perceived benefit
6 ("wanted to feel better"), cues to action ("encouraged by a health professional, etc."), fear ("death
7 of a family member, etc."), appear to affect behavioral decisions.
8
9
10
11

12 **Influence of gender**

13
14 Although females were unintentionally oversampled, our analysis includes a gender-wise
15 stratification and we present data separately whenever we observed marked gender differences.
16 In our study, more men than women smoked tobacco or drank alcohol. The prevalence of
17 hypertension and self-reported diabetes was similar in both groups. On the other hand, low
18 physical activity and obesity were higher in the women. Although men had better knowledge of
19 heart disease, their attitude was similar to those reported by women. However, more women than
20 men strongly agreed that their health was ultimately determined by God or a higher power.
21 Female gender is usually significantly related to good knowledge of cardiovascular disease.⁵²
22 One major factor for the gender discrepancy in our study is that more women (33.2%) than men
23 (11.8%) lacked formal education, which in itself is alarming. In fact, illiteracy is one of the main
24 risk factors along with behavioral factors in rural/urban Asian Indian women.⁵³
25
26
27
28
29
30
31
32
33
34
35

36 **Effect of other socio-demographic variables on KAP**

37
38 Age-wise, KAP/behavior scores decreased around 4% from the youngest to the oldest age group.
39 An earlier study reported a higher level of knowledge among young respondents.³⁶ Interestingly,
40 a study in adolescents reported a very poor level of cardiovascular knowledge.⁵⁴ Otherwise,
41 knowledge levels rise linearly from childhood until middle age, when the level starts to plateau.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

⁵⁵ In our study, better-educated respondents ate more fruit and consumed less tobacco and alcohol, but they also reported less physical activity and more hypertension. The highest education category (high school or more) scored higher than the lowest category (no formal schooling) but only by 3%–8%. Other studies observed a consistent link between better KAP scores and higher education.^{31,36,52,56} Occupation-wise, employed respondents knew more about biological causes (e.g., high blood pressure, diabetes, and high cholesterol), possibly due to more

1
2
3 education, wider exposure to information, and better access to screening and treatment. A
4 Canadian study reported similar findings.³⁶

8 **Prompted vs. unprompted responses**

9
10 Although we used unprompted responses to evaluate general knowledge, we used prompted
11 responses to calculate scores, thus providing a denominator (“full marks”) to calculate median
12 score %. In other settings, KAP studies have variably used prompted^{33,39,47,57} or unprompted
13 questions;^{16, 31, 38} while some studies used both.³⁵ Other studies did not identify their chosen
14 method.³² Although the accuracy of prompted vs. unprompted questions in knowledge
15 assessment has been a topic of debate, prompted questions confer a higher level of knowledge.⁵⁸
16 When we analyzed knowledge responses based on prompted responses, the percent of correct
17 answers increased drastically. The psychological explanation is that respondents must simply
18 “recognize” answers to prompted questions, but they must engage in the more difficult mental
19 process of “recalling” answers to unprompted questions.⁵⁹ We believe that unprompted responses
20 more accurately reflect knowledge level because they emerge from a thought process that in
21
22
23
24
25
26
27
28
29
30

31 **Comparison between levels of knowledge, attitude, and practice towards cardiovascular** 32 **disease**

33
34
35 The findings of the study indicate that there is gap in knowledge attitude and practices. Poor
36 relation between KAP scores was unsurprising because a linear relationship between variables
37 may not exist in cardiovascular health (e.g. highly satisfactory knowledge=high satisfactory
38 attitude and behavior). Valente argued that each of six possible behavior change models contains
39 a varying order of KAP/behaviour and that a population can be described in terms of the
40 percentage that fits into each model.⁶⁰ This implies that no universal intervention can improve
41 the cardiovascular health KAP of every individual in a community. The challenge lies in
42 identifying individuals who fit into different linear models. Furthermore, health behavior theories
43 (e.g., social cognitive theory and the theory of reasoned action)⁴⁸ suggest that knowledge and
44 attitude are only two components of the pathways that lead to behavior. Constructs (e.g., self-
45 efficacy) and other personal and environmental factors (in social cognitive theory) and perceived
46 behavioral control, subjective norms, and ultimately intention (in the theory of reasoned action)
47 play crucial roles in shaping one’s health behavior.⁴⁸
48
49
50
51
52
53
54
55
56
57
58
59
60

Study limitations

Females were unintentionally oversampled (72.7%) although the male:female ratio of the study population is 1:1.15.²⁰ Women were more likely to be available in the households when enumerators visited as most of the females in our study were housewives. The age-range is limited to 25-59 years as the age groups 60-64 and 70-74 years only constitute 1-2% of the total JD-HDSS population. Most of the respondents belonged to the caste/ethnic groups Brahmin, Chhetri, and Newar which are not representative of all caste/ethnic groups in Nepal. Further, respondents' and interviewers' mood as well as place and time of interview may influence the study and a recall bias may occur when questions are asked about history of CVD risk factors.

CONCLUSION

Our community-based cross-sectional HARDIC study demonstrated poor knowledge, inappropriate attitude, and highly adverse behavior regarding cardiovascular health in a semi-urban community of Nepal even among those already affected by CVD. We aimed to determine a baseline KAP in the community to aid the planning of health promotion activities. Our results show the potential and possibility that community health promotion can improve cardiovascular health literacy.

Figure legend

Figure 1: Sex-wise distribution of cardiovascular risk factor status in the study population.

ACKNOWLEDGMENTS

The authors are grateful to the study participants. We acknowledge all enumerators; field supervisors (Rachana Shrestha, Vishal Bhandari, Chandra Shova Khaitu, and Shova Poudel); and our field coordinator, Prof. Dr. Muni Raj Chhetri. We thank scientific editor Karen Williams (Kwills Editing Services, Weymouth, MA, USA) for providing professional English-language editing of this article.

CONTRIBUTORS

AV designed the study, performed statistical analysis, and drafted the manuscript. URA performed statistical analysis and revised the manuscript. AK helped design the study and provided critical revision of the manuscript. All authors have read and approved the final manuscript.

FUNDING STATEMENT

This study was supported by grants from the Wilhelm & Martina Lundgren's Foundation (vet1-367/2011 and vet1-379/2012) and the University of Gothenburg, Sweden through a "Global University" grant (A11 0524/09).

COMPETING INTERESTS

None declared.

DATA SHARING

No additional data available.

REFERENCES

1. Alwan A. *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organization; 2011.
2. World Health Organization. *Noncommunicable diseases in the South-East Asia Region: Situation and response 2011*. New Delhi: World Health Organization , Regional Office for South-East Asia; 2011.
3. Lozano R, Naghavi M, Foreman K, *et al*. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128.
4. Reddy KS. Cardiovascular diseases in the developing countries: dimensions, determinants, dynamics and directions for public health action. *Public Health Nutr* 2002;5:231-37.
5. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998;97:596-601.
6. Engelgau MM, El-Saharty S, Kudesia P, *et al*. *Capitalizing on the demographic transition: tackling noncommunicable diseases in South Asia*: Washington, DC: World Bank Publications; 2011.
7. Ezzati M, Lopez AD, Rodgers A, *et al*. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. Geneva: World Health Organization; 2004.

- 1
2
3 8. Khaw KT, Wareham N, Bingham S, *et al.* Combined impact of health behaviours and
4 mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS medicine*
5 2008;5:e12.
6
7
- 8
9 9. Mendis S, Puska P, Norrving B. *Global atlas on cardiovascular disease prevention and*
10 *control.* Geneva: World Health Organization; 2011.
11
12
- 13
14 10. Alwan A, Maclean DR, Riley LM, *et al.* Monitoring and surveillance of chronic non-
15 communicable diseases: progress and capacity in high-burden countries. *Lancet* 2010;376:1861-
16 68.
17
18
- 19
20 11. Mendis S, Alwan A. *Prioritized research agenda for prevention and control of*
21 *noncommunicable diseases.* Geneva: World Health Organization; 2011.
22
23
- 24
25 12. Puska P. From Framingham to North Karelia: from descriptive epidemiology to public health
26 action. *Prog Cardiovasc Dis* 2010;53:15-20.
27
28
- 29
30 13. Pearson TA, Bazzarre TL, Daniels SR, *et al.* American Heart Association guide for
31 improving cardiovascular health at the community level: a statement for public health
32 practitioners, healthcare providers, and health policy makers from the American Heart
33 Association Expert Panel on Population and Prevention Science. *Circulation* 2003;107:645-51.
34
35
- 36
37 14. Lloyd-Jones DM, Hong Y, Labarthe D, *et al.* Defining and Setting National Goals for
38 Cardiovascular Health Promotion and Disease Reduction The American Heart Association's
39 Strategic Impact Goal Through 2020 and Beyond. *Circulation* 2010;121:586-613.
40
41
- 42
43 15. Khan MS, Jafary FH, Faruqui AM, *et al.* High prevalence of lack of knowledge of symptoms
44 of acute myocardial infarction in Pakistan and its contribution to delayed presentation to the
45 hospital. *BMC Public Health* 2007;7:284.
46
47
- 48
49 16. Limbu YR, Malla R, Regmi SR, *et al.* Public knowledge of heart attack in a Nepalese
50 population survey. *Heart & Lung: The Journal of Acute and Critical Care* 2006;35:164-69.
51
52
- 53
54 17. Upadhyay DK, Palaian S, Shankar PR, *et al.* Knowledge, attitude and practice about diabetes
55 among diabetes patients in Western Nepal. *Rawal Med J* 2008;33:8-11.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
18. Vaidya A. Tackling cardiovascular health and disease in Nepal: epidemiology, strategies and implementation. *Heart Asia* 2011;3:87-91.
 19. World Health Organization. Global Health Observatory Data Repository. <http://apps.who.int/ghodata> (accessed 28 Jun 2013).
 20. Aryal UR, Vaidya A, Shakya-Vaidya S, *et al.* Establishing a health demographic surveillance site in Bhaktapur district, Nepal: initial experiences and findings. *BMC Research Notes* 2012;5:489.
 21. Vaidya A, Shakya S, Krettek A. Obesity prevalence in nepal: public health challenges in a low-income nation during an alarming worldwide trend. *International Journal of Environmental Research and Public Health* 2010;7:2726-44.
 22. Kish L. Sampling organizations and groups of unequal sizes. *Am Sociol Rev* 1965; 30:564-72.
 23. WHO STEPS Instrument (Core and Expanded): The WHO STEPwise approach to chronic disease risk factor surveillance (STEPS). Available at: http://www.who.int/chp/steps/STEPS_Instrument_v2.1.pdf.
 24. Lin W, Yang H, Hang C, *et al.* Nutrition knowledge, attitude, and behavior of Taiwanese elementary school children. *Asia Pac J Clin Nutr* 2007;16:534-46.
 25. Torabi MR, Yang J, Li J. Comparison of tobacco use knowledge, attitude and practice among college students in China and the United States. *Health Promot Internation* 2002;17:247-53.
 26. Kosaryan M, Vahidshahi K, Siami R, *et al.* Knowledge, attitude, and practice of reproductive behavior in Iranian minor thalassemia couples. *Saudi Med J* 2009;30:835-39.
 27. Neamatollahi H, Ebrahimi M, Talebi M, *et al.* Major differences in oral health knowledge and behavior in a group of Iranian pre-university students: a cross-sectional study. *J Oral Sci* 2011;53:177-84.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
28. Sharda AJ, Shetty S. A comparative study of oral health knowledge, attitude and behaviour of first and final year dental students of Udaipur city, Rajasthan, India. *International Journal of Dental Hygiene* 2008;6:347-53.
29. Yap J, Lee V, Yau T, *et al.* Knowledge, attitudes and practices towards pandemic influenza among cases, close contacts, and healthcare workers in tropical Singapore: a cross-sectional survey. *BMC Public Health* 2010;10(1):442.
30. Maimaiti N, Shamsuddin K, Abdurahim A, *et al.* Knowledge, Attitude and Practice regarding HIV/AIDS among University students in Xinjiang. *Global Journal of Health Science* 2010;2:51.
31. Chen W, Yu Y, Glaser K. The Knowledge and Attitudes of Coronary Heart Disease Prevention among Middle and Older Aged People in a Community in Taipei. *Taiwan Geriatrics & Gerontology* 2009;4:251-62.
32. Dodani S, Mistry R, Khwaja A, *et al.* Prevalence and awareness of risk factors and behaviours of coronary heart disease in an urban population of Karachi, the largest city of Pakistan: a community survey. *Journal of Public Health* 2004;26:245-49.
33. Saeed O, Gupta V, Dhawan N, Streja L, Shin JS, Ku M, Bhoi S, Verma S: Knowledge of modifiable risk factors of Coronary Atherosclerotic Heart Disease (CASHD) among a sample in India. *BMC international health and human rights* 2009, 9(1):2.
34. Gazmararian JA, Williams MV, Peel J, *et al.* Health literacy and knowledge of chronic disease. *Patient Educ Couns* 2003;51(3):267-75.
35. Mosca L, Jones WK, King KB, *et al.* Awareness, perception, and knowledge of heart disease risk and prevention among women in the United States. *Arch Fam Med* 2000;9(6):506.
36. Potvin L, Richard L, Edwards AC. Knowledge of cardiovascular disease risk factors among the Canadian population: relationships with indicators of socioeconomic status. *Can Med Assoc J* 2000;162:S5-S11.
37. Lim K, Sumarni M, Amal N, *et al.* Tobacco use, knowledge and attitude among Malaysians age 18 and above. *Trop Biomed* 2009;26:92-99.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
38. Wong BM, Garcia Y, Barr A, *et al.* Cardiovascular risk factor awareness in a disadvantaged inner-city population—implications for preventive strategies. *Can J Cardiol* 2008;24:677-82.
39. Schweigman K, Eichner J, Welty TK, *et al.* Cardiovascular disease risk factor awareness in American Indian communities: The Strong Heart Study. *Ethnicity and disease* 2006;16:647.
40. Lenfant C, Chobanian AV, Jones DW, *et al.* Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7). *Circulation* 2003;107:2993-94.
41. United Nations Children Fund. *Baseline Survey of the Knowledge, Attitude and Practice (KAP) of Parents/Guardians on Early Childhood Development and Primary Education in Nepal.* Lalitpur, Nepal: UNICEF Nepal Country Office; 2009.
42. Ministry of Health and Population, Nepal. *Nepal non-communicable diseases risk factors survey 2007.* Kathmandu, Nepal: Ministry of Health and Population, Nepal; 2008.
43. Vaidya A, Pathak RP, Pandey MR. Prevalence of hypertension in Nepalese community triples in 25 years: a repeat cross-sectional study in rural Kathmandu. *Indian Heart J* 2012;64:128-31
44. World Health Organization. WHO Risk Factor Database. <https://apps.who.int/infobase/Indicators.aspx> (accessed 4 Aug 2013).
45. An DTM, Minh HV, Huong LT, *et al.* Knowledge of the health consequences of tobacco smoking: a cross-sectional survey of Vietnamese adults. *Global health action* 2013;6:18707.
46. Brook RD, Franklin B, Cascio W, *et al.* Air pollution and cardiovascular disease A statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association. *Circulation* 2004;109:2655-71.
47. Nguyen TT, Liao Y, Gildengorin G, *et al.* Cardiovascular risk factors and knowledge of symptoms among Vietnamese Americans. *Journal of General Internal Medicine* 2009;24:238-43.

- 1
2
3
4 48. Redding CA, Rossi S, Rossi R, *et al.* Health behavior models. *The International Electronic*
5 *Journal of Health Education* 2000;3:180-93.
6
7
8 49. Avis NE, Smith KW, McKinlay JB. Accuracy of perceptions of heart attack risk: what
9 influences perceptions and can they be changed?. *Am J Public Health* 1989; 79:1608-12.
10
11
12 50. Sharma D, Bkc M, Rajbhandari S, *et al.* Study of prevalence, awareness, and control of
13 hypertension in a suburban area of Kathmandu, Nepal. *Indian Heart J* 2006;58:34-37.
14
15
16 51. Vaidya A, Pokharel P, Karki P, *et al.* Exploring the iceberg of hypertension: a community
17 based study in an eastern Nepal town. *Kathmandu Univ Med J* 2007;5:349-59.
18
19
20
21 52. Attarchi M, Mohammadi S, Nojomi M, *et al.* Knowledge and Practice Assessment of
22 Workers in a Pharmaceutical Company about Prevention of Coronary Artery Disease. *Acta Med*
23 *Iran* 2012;50:697-703.
24
25
26
27 53. Pandey RM, Agrawal A, Misra A, *et al.* Population-based intervention for cardiovascular
28 diseases related knowledge and behaviours in Asian Indian women. *Indian Heart J* 2012;65:40-
29 7.
30
31
32
33 54. Vanhecke TE, Miller WM, Franklin BA, *et al.* Awareness, knowledge, and perception of
34 heart disease among adolescents. *European Journal of Cardiovascular Prevention &*
35 *Rehabilitation* 2006;13:718-23.
36
37
38
39 55. White CW, Albanese MA. Changes in cardiovascular health knowledge occurring from
40 childhood to adulthood. A cross-sectional study. *Circulation* 1981;63:1110-15.
41
42
43
44 56. Yahya R, Muhamad R, Yusoff HM. Association between Knowledge, Attitude and Practice
45 on Cardiovascular Disease among Women in Kelantan, Malaysia. *International Journal of*
46 *Collaborative Research on Internal Medicine & Public Health* 2012;4:1507-23.
47
48
49
50 57. Muhamad R, Yahya R, Yusoff HM, *et al.* Knowledge, Attitude and Practice on
51 Cardiovascular Disease among Women in North-Eastcoast Malaysia. *International Journal of*
52 *Collaborative Research on Internal Medicine & Public Health* 2012;4:85-98.
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

58. Waller J, McCaffery K, Wardle J. Measuring cancer knowledge: comparing prompted and unprompted recall. *Br J Psychol* 2004;95:219-34.

59. Glass AL, Holyoak KJ. *Cognition* . Second ed. New York: McGraw-Hill; 1986.

60. Valente TW, Paredes P, Poppe PR. Matching the message to the process the relative ordering of knowledge, attitudes, and practices in behavior change research. *Human Communication Research* 1998;24:366-85.

For peer review only

SCORE SYSTEM FOR KNOWLEDGE, ATTITUDE & PRACTICE/BEHAVIOUR

Responses to questions on cardiovascular health knowledge, attitude and practice/behaviour were given scores, in order to compare and correlate the three components.

Knowledge score

Prompted responses of the five sections on knowledge were considered for calculating knowledge score. The total score on knowledge was calculated by combining scores of the below-mentioned five sections of knowledge: cause of heart diseases, warning signs of heart attack, action in case of a heart attack, knowledge regarding foods that are good for heart and knowledge regarding prevention of heart diseases. Maximum possible score for the Knowledge part was thus $11+15+7+10+10= 53$.

Cause of heart diseases: Eleven conventional causes were read out to the respondent one by one. Score 1 was given if the answer was 'yes'. No negative marking was given for 'no' or 'don't know' responses. Maximum score possible for the section was 11.

Warning signs of heart attack: Nine common symptoms which are considered as warning signs of a heart attack by the American Heart Association (ref) and two incorrect signs (pain in the abdomen and pain in the leg) were mingled up and read to the respondent who were asked if the symptoms are correct. If answered 'yes', scores were allocated as follows: 3 for chest pain; 2 each for breathlessness, sweating and vomiting; 1 each for pain in the arm, teeth or jaw, light-headedness and unconsciousness; 0 for pain in the abdomen and pain in the leg. Additionally, if 'no' response came for pain in the abdomen and pain in the leg, 1 mark was given. No marks were given if the response was 'don't know'. Maximum score possible was 15 for the section.

Action in case of a heart attack: Respondents were asked about actions that they would do or would not do in case of a heart attack. If the responses were 'yes', then scores were given as follows: treat at home (0), immediately take the person to a hospital (2), take to a pharmacy/medicine shop (0), give home-made therapy (0), wait for appointment with a heart specialist (0), take the person to a traditional healer (0). No mark was given if the answer was 'no' or 'don't know'. Maximum score possible for the section was 7.

Foods that are good for heart: Each respondent was asked if they considered a particular food item from a pre-selected list to be good for the heart. The response was recorded as 'yes', 'no' or 'don't know'. Depending on whether the particular food is actually good (e.g. fruits) for heart or not (e.g. fat), 1 mark was given for each correct answer. No mark was given if the response was 'don't know'. Maximum score possible for the section was 10.

Prevention of heart diseases: Each respondent was asked about the various preventive actions that are considered good for improving heart-health. The response was recorded as 'yes', 'no' or 'don't know'. One mark was given if the response was 'yes'. Maximum score possible for the section is 10 for the ten questions.

Attitude score

Each respondent was asked a 14-item questions related to his/her attitude towards heart-health. Responses were recorded as strongly agree or disagree, or somewhat agree or disagree, or don't know/can't say. Depending on whether it was a proper attitude or not, scores from 1 to 4 were allotted. A score of '0' was given for don't know/can't say. Similarly, participants were also asked about their opinions on various preventive aspects of heart health and the responses were recorded with 'very helpful' and 'not at all helpful' as the extremes. Maximum possible score for the attitude section was $14 \times 4 = 56$.

Practice and behaviour score

Each respondent was asked if s/he undertook any particular action in the previous year that would help towards having a better heart health. Scores were given accordingly. Additionally, participants were also given scores according to their risk factor status. No negative marking was given. Maximum score possible was 25.

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	Comment [AV1]: Yes Comment [AV2]: Yes
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	Comment [AV3]: Yes
Objectives	3	State specific objectives, including any prespecified hypotheses	Comment [AV4]: Yes
Methods			
Study design	4	Present key elements of study design early in the paper	Comment [AV5]: Yes
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Comment [AV6]: Yes
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	Comment [AV7]: Yes
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Comment [AV8]: Yes
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Comment [AV9]: Yes
Bias	9	Describe any efforts to address potential sources of bias	Comment [AV10]: Yes
Study size	10	Explain how the study size was arrived at	Comment [AV11]: Purposive
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Comment [AV12]: Yes
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) If applicable, describe analytical methods taking account of sampling strategy (e) Describe any sensitivity analyses	Comment [AV13]: Yes Comment [AV14]: Yes Comment [AV15]: Not applicable
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	Comment [AV16]: Done, mentioned in the tables, etc as well Comment [AV17]: Yes Comment [AV18]: Not relevant
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest	Comment [AV19]: Yes Comment [AV20]: Yes
Outcome data	15*	Report numbers of outcome events or summary measures	Comment [AV21]: Yes
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Comment [AV22]: Yes Comment [AV23]: Yes
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion		
Key results	18	Summarise key results with reference to study objectives
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence
Generalisability	21	Discuss the generalisability (external validity) of the study results
Other information		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based

- Comment [AV24]: Yes
- Comment [AV25]: Yes
- Comment [AV26]: Yes
- Comment [AV27]: Yes
- Comment [AV28]: Yes

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Cardiovascular health knowledge, attitude, and practice/behavior in an urbanizing community of Nepal: a population-based cross-sectional study from Jhaukhel-Duwakot Health Demographic Surveillance Site

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Swedish (Sweden)

Abhinav Vaidya^{1,2}, Umesh Raj Aryal^{1,2}, and Alexandra Krettek^{2,3}

Formatted: Swedish (Sweden)

Formatted: Swedish (Sweden)

Formatted: None

Formatted: Space After: 0 pt

¹Department of Community Medicine, Kathmandu Medical College, Kathmandu, Nepal

²Nordic School of Public Health NHV, Gothenburg, Sweden, and

Formatted: None, Space After: 0 pt

³Department of Internal Medicine and Clinical Nutrition, Institute of Medicine, Sahlgrenska Academy at University of Gothenburg, Gothenburg, Sweden

Formatted: Space After: 0 pt

Correspondence to

Formatted: None, Space After: 0 pt

Abhinav Vaidya; Abhinav.Vaidya@nhv.se

Keywords: knowledge, attitude, and practice/behavior; Nepal; cardiovascular health; health literacy

Word count: 4,894,337 (excludes title page, abstract, references, tables, and figures, per author guidelines)

Formatted: Font: Not Bold

Formatted: Font: Not Bold

1

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

ABSTRACT

Objectives: This study determined the knowledge, attitude, and practice/behavior of cardiovascular health in residents of a semi-urban community of Nepal.

Design: To increase understanding of knowledge, attitude, and practice/behavior toward cardiovascular health, we conducted in-home interviews using a questionnaire based on WHO STEPS and other resources, scoring all responses.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

~~We also recorded blood pressure and took anthropometric measurements.~~

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Setting: Our study was conducted as part of the HARDIC project in the Jhaukel-Duwakot Health Demographic Surveillance Site in two urbanizing villages near Kathmandu.

Participants: The study population included 777 respondents from six randomly selected clusters in both villages.

Results: Seventy percent of all participants were female and 26.9% lacked formal education. The burden of cardiovascular risk factors was high; 20.1% were current smokers, 43.3% exhibited low physical activity, and 21.6% were hypertensive. Participants showed only poor knowledge of heart disease causes; 29.7% identified hypertension and 11% overweight and physical activity as causes, whereas only 2.2% identified high blood sugar as causative. Around 60% of respondents did not know any heart attack symptoms compared with 20% who knew 2-4 symptoms. Median % scores for knowledge, attitude, and practice/behavior were 79.3, 74.3, and 48.0, respectively. Nearly 44% respondents had insufficient knowledge and less than 20% had highly satisfactory knowledge. Among those with highly satisfactory knowledge, only 14.7% had highly satisfactory attitude and 19.5% and 13.9% had satisfactory and highly satisfactory practices, respectively.

Conclusions: Our study demonstrates a gap between cardiovascular health knowledge, attitude, and practice/behavior in a semi-urban community in a low-income nation, even among those already affected by CVD.

Formatted: None

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

ARTICLE SUMMARY

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Article focus

- We aimed to gain a deeper understanding of knowledge, attitude, and behavior/practice regarding cardiovascular health in a semi-urban community of Nepal, a low-income South Asian country.

Key messages

Formatted: None

- ~~We found~~ The study site has a high burden of cardiovascular risk factors. In particular, low physical activity and overweight/obesity; ~~cardiovascular risk factors in our study site exceeded~~ exceed the national average.
- Low-Unprompted responses to cardiovascular health questions revealed a poor level of knowledge, and respondents' attitude toward cardiovascular health showed low risk perception ~~combined with poor relation.~~
- Poor correlation between knowledge, attitude, and behavior scores imply that commonly applied health promotion strategies that merely increase knowledge of cardiovascular health may be insufficient, thus emphasizing the need for a more holistic approach to disease prevention.

Strengths and limitations of the study

Formatted: None

- This study is the first to extensively explore knowledge, attitude, and behavior/practice regarding cardiovascular health in Nepal ~~and was conducted in,~~
- Random sampling derived from a sampling frame generated from a health demographic surveillance site ~~which~~ provides the logistic advantage of further follow-up and intervention.
- Using both prompted and unprompted questions in the same study illustrates the variation that may be obtained by these approaches.
- Except for the WHO STEPS questionnaire on behavioral risk factors, we used questions from many unvalidated sources.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

- Unintentional oversampling of women respondents yielded unsegregated gender-wise results that are more likely to represent knowledge, attitude, and practice/behavior in women.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Abstract:

Objectives: This study aimed to determine the knowledge, attitude, and practice/behavior of cardiovascular health in residents of a semi-urban community of Nepal.

Design: To increase understanding of knowledge, attitude, and practice/behavior toward cardiovascular health, we conducted in-home interviews using a questionnaire based on WHO STEPS and other resources, scoring all responses. We also recorded blood pressure and took anthropometric measurements.

5

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Setting: Our study was conducted as part of the HARDIC project at the Jhaukhel-Duwakot Health Demographic Surveillance Site in two urbanizing villages near Kathmandu. Our results present baseline data from the project that aims to improve cardiovascular health literacy through health promotion.

Participants: The study population included 777 respondents from six randomly selected clusters in both villages.

Primary outcome measure: Proportion of the study population with correct knowledge about the causes, manifestation, immediate management, and preventability of cardiovascular diseases.

Our data reveals participants' attitude toward cardiovascular health and its preventability as well as cardiovascular health behavior, as reflected in the risk factor status and positive actions taken toward achieving them.

Secondary outcome measure: Our data calculated and correlated participants' scores regarding knowledge, attitude, and practice/behavior.

Results: Participants showed only poor knowledge of heart disease causes, heart attack symptoms, and appropriate action. In general, the participants' attitude toward heart health was inappropriate. The burden of cardiovascular risk factors among participants was high. Median percent scores for knowledge, attitude, and practice/behavior were 79.25, 74.29, and 48.00, respectively. We observed poor correlations between the Pearson coefficient for knowledge and attitude; attitude and behavior/practice; and knowledge and behavior/practice (0.025, 0.126, and 0.121, respectively).

Conclusions: Our study demonstrates poor level of cardiovascular health knowledge, attitude, and practice/behavior in a semi-urban community in a low-income nation. Tailoring health promotion activities to our findings might improve public health literacy for cardiovascular disease.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal, No bullets or numbering

Formatted: None

INTRODUCTION

Noncommunicable diseases (NCDs) pose increasingly important public health problems in low- and middle- income countries (LMICs). In 2008, 80% of global NCD deaths occurred in LMICs,¹ and NCDs recently accounted for 7.9 million of 14.5 million deaths (54%) in Southeast Asia,² and were responsible for two of every three deaths (34.5 million) worldwide in 2010.³

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Poverty, illiteracy, poor health infrastructure, and demographic transition, including increasing life expectancy, are major contributors to rising burden of NCDs in LMICs the region. Other postulated causes include poor fetal and childhood nutrition.^{4-6,3,4} A similar pattern is evident in South Asian countries like Nepal, where NCDs currently occur more frequently than communicable diseases and behavioral risk factors such as tobacco smoking are high.^{6,5}

Globally, cardiovascular diseases (CVDs) are responsible for one in four comprise 39% of all NCD deaths (12.9 million).^{2,6} Eighty percent of CVDs result from tobacco smoking, unhealthy diet, physical inactivity, and harmful use of alcohol.⁷ The risk of all-cause mortality among individuals who score poorly in behavioral factors is four fold higher than those who score best.⁸

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

In high-income countries, declining CVD mortality is largely attributed to the reduction of behavioral risk factors.⁹ Underlying socioeconomic parameters and societal influences affect exposure and vulnerability to CVD risk factors. Therefore, surveillance of NCD risk factors should include social determinants of cardiovascular health.^{1,10} Currently, understanding the individual and social determinants of cardiovascular health behaviors is among the top 20 priority areas for NCD research in LMICs.¹¹

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

As a precursor of favorable health behavior, experts have advocated health literacy since community-based interventions (e.g., Finland's North Karelia Project) began exploring the potential impact of behavioral changes on CVD.^{1,12} The World Health Organization (WHO) promotes public awareness of diet and physical activity as a "best buy" to improve

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

7

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

cardiovascular behavior.⁹ ~~Similarly, the American Heart Association emphasizes the importance of improving community awareness of cardiovascular health issues.~~^{13, 14} However, achieving desired health behavior requires deeper understanding of the underlying social context of disease.^{43, 44, 15} Hence, studies on knowledge, attitude, and practice (KAP) increase our understanding of cardiovascular health literacy and community behavior.^{45, 16} Currently, such studies have been limited in South Asia.

~~Earlier studies report~~In Pakistan, inability to recognize the symptoms of heart attack associates independently with delayed hospitalization.¹⁷ ~~An earlier study reported a~~ low level of knowledge about heart attack^{46, 18} and diabetes symptoms in Nepal.^{47, 19} Nepal's high prevalence of underlying behavioral risk factors, particularly tobacco and alcohol consumption, and poor cardiovascular health knowledge is a matter of concern.^{48, 20} ~~Nationwide in Nepal, the age group 25+ exhibits fasting blood glucose of 5.5 mmol/L for males and 5.4 mmol/L for females, systolic blood pressure of 125.9 mmHg and 126.6 mmHg for males and females respectively, and total cholesterol of 4.1 mmol/L (males) and 4.3 mmol/L (females).~~⁴⁹

Currently, attempts to improve cardiovascular health knowledge and behavior in Nepal focus mainly on campaigns that transmit information but pay little attention to the population's actual health literacy. ~~We therefore~~ To gain greater insight into this important issue, we undertook a community-based KAP study in the Jhaukhel-Duwakot Health-Demographic Surveillance Site (JD-HDSS) outside Kathmandu ~~where~~.²¹ The baseline census in these villages suggested that NCDs, particularly CVDs, are the predominant cause of morbidity and mortality.^{20, 21} Additionally, our pilot study in the Duwakot population demonstrated poor cardiovascular health knowledge.²² ~~Our study is part of the~~

Heart-Health Associated Research and Dissemination in the Community (HARDIC) ~~project,~~ ~~which~~ is the cardiovascular health component of JD-HDSS. In Nepalese, hardic means "heartily" or "from the heart". ~~This~~ Using community-level interventions, HARDIC aims to improve cardiovascular health literacy and practice/behavior and this paper reports the baseline results from HARDIC ~~and~~. Our study is the first to conduct a detailed analysis of ~~cardiovascular health knowledge, attitude, and practice/behavior~~ KAP/health literacy in ~~an urbanizing community of Nepal~~ relation to cardiovascular health.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

Formatted: Pattern: Clear

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

METHODS

Formatted: None, Space After: 0 pt

Ethical considerations

~~The Nepal Health Research Council and the Institutional Review Board of Kathmandu Medical College approved this study. Following a pattern established at the outset of JD-HDSS, we consulted local leaders and sought their informal permission for HDSS activities. In addition, we periodically consulted with health and administrative authorities in both villages. All participants gave informed consent, and privacy was maintained during the interview process. To maintain confidentiality, all data were secured in the HDSS office. Computer data were accessible only to the research team. Although respondents received no monetary incentives, we provided a referral that included discounted consultation fees at Kathmandu Medical College or Nepal Medical College for self-reported illness and newly diagnosed hypertension.~~

Formatted: Space After: 0 pt

Study site and population

Formatted: None, Space After: 0 pt

We conducted our descriptive, community-based, cross-sectional study between September and November 2011 in the JD-HDSS of Bhaktapur district in the Kathmandu Valley, 13 kilometers from Kathmandu. An earlier report described the establishment of the surveillance site.²⁰²¹

Study population and sample size

~~This study is the first step in the larger HARDIC project that will be conducted in the whole JD-HDSS.~~ Duwakot and Jhaukhel each contain nine administrative wards ~~and we~~. We randomly selected three wards from each village. Based on the JD-HDSS 2010 baseline census,²⁰²¹ we compiled a complete list of households containing adults aged 25-59 years in the selected wards.

Formatted: Normal

Next, we selected one adult, of either sex, from all interview households ~~and surveyed 789 of 840 households (non-response rate = 6.07%). We applied.~~ If a household contained more than ~~one adult, we used~~ the Kish technique to select the respondent.²²²³

Study tools

Formatted: None

Interviews based on the WHO-NCD STEPS questionnaire

We structured the basic questionnaire ~~and tailored it for site-specific responses in Nepal~~ using the WHO-NCD STEPS (STEP 1 and 2 questionnaires) for demographic information, behavioral and physical measurements²²²⁴ and other sources ~~for addressing knowledge, attitude and~~

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

~~practice/behavior issues.~~^{15, 24, 17, 25-39} ~~Responses to questions on cardiovascular health knowledge,~~
attitude and practice/behavior were given scores in order to compare and correlate the
components. A detailed description of the scoring system is given as an appendix to this article.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

Interviewers

~~Questions were related to respondents'~~ ~~Twelve local female enumerators (two per ward)~~
~~conducted the interviews. All enumerators had completed at least grade 10 in school. Supervisors~~
~~included four public health graduates; a field coordinator with Master and PhD degrees in public~~
~~health; and two PhD students/authors (AV and URA).~~

Data Collection

Interview process

~~The interviews were conducted at the participants' homes. The initial part of the interview~~
~~consisted of questions related to demographic profile. Questions then followed relating to~~
knowledge of cardiovascular health (e.g., causes and risk factors, warning signs, and action for
suspected heart attack); heart-healthy diet; and disease prevention. The first two areas included
unprompted (spontaneous, open-ended) and prompted (clues provided, closed-ended) responses,
and the latter two areas included only prompted responses. We asked respondents about their
attitude toward different aspects of cardiovascular health: did they perceive themselves at risk for
CVD? Did they think behavior change would reduce their risk of developing CVD? ~~The~~
~~responses were recorded in Likert scale (strongly agree, somewhat agree, don't know, somewhat~~
~~disagree, strongly agree).~~

Formatted: English (U.S.)

Behavioral factors

Behavior-related questions were based on Step 1 of the WHO STEPwise approach to
surveillance (STEPS), which uses a questionnaire to collect data on risk factors.²⁴ Our
questions probed current and past smoking behavior; smokeless tobacco; alcohol consumption in
the past 30 days (current drinker), and drinking frequency among current drinkers. The physical
activity section asked respondents to report the number of days and amount of time they
participated in vigorous and moderate activities at work, leisure, and travel. ~~Additional practice~~

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

~~and behavioral questions explored whether participants had engaged in any heart friendly behavior in the previous year and for what reason.~~

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

~~Physical measurements~~

Formatted: None, Space After: 0 pt

~~Enumerators recorded anthropometric measurements (i.e., weight, height, and waist and hip circumferences) according to STEPs guidelines.²³ Weight was measured to one decimal place using a Microlife BR 9201 Weighing Scale (Microlife AG Swiss Corporation, Widnau, Switzerland). Height and waist and hip circumferences were measured using non-stretchable tapes (Jonson Tapes Ltd, New Delhi, India). Blood pressure was determined digitally by Microlife BP 3API 3E (Microlife AG Swiss Corporation, Widnau, Switzerland). Enumerators recorded three readings for each respondent, taken at 5 minute intervals. We averaged the readings to attain the final blood pressure measurements.~~

~~Diagnostic criteria for sub-classification~~

~~Respondents with known history of hypertension (diagnosed cases) and those diagnosed during the study according to Joint National Committee –VII⁴⁰ were termed as having “hypertension”.~~

~~Data management~~

~~Supervisors checked all completed questionnaires and, if necessary, asked enumerators to revisit households. After omitting incomplete questionnaires, we included 777 individuals in the analysis. Except for missing recordings for height, weight, or blood pressure, data reported in our KAP study covers all 777 respondents. We outsourced data entry to a team of data entry operators, public health graduates by qualification. Data was coded and entered in Epidata version 2.1.~~

~~Statistical analyses~~

~~Data was analyzed with SPSS version 17.0 (IBM, Armonk, New York, USA) and STATA version 10.0 (StataCorp, Texas, USA). After calculating the Metabolic Equivalent of Task (MET) minutes/week, we grouped participants as highly active, moderately active, or inactive (low physical activity) in accord with the STEPS manual. Additional practice and behavioral questions explored whether participants had engaged in any heart-friendly behavior in the previous year and for what reason.~~

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

~~For scoring knowledge, attitude, and practice/behavior questions and to compare and correlate the three components, we scored responses regarding KAP/behavior for cardiovascular health. The maximum possible score for KAP/behavior was 53, 56 and 25, respectively. Further, total knowledge score was classified into the following five categories based on the quintile scores: ≤20 % ‘highly insufficient’; 21–40 % ‘insufficient’; 41–60 % ‘sufficient’; 61–80 % ‘satisfactory’; >80 % ‘highly satisfactory’.⁴¹~~

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

~~Categorical data is presented as number, percentages, and continuous data as median and inter-quartile range (IQR). KAP percentage scores are presented as median. We used Chi-square test to compare proportions among demographic variables and Mann-Whitney U test to compare scores from males and females. $P < 0.05$ was considered statistically significant.~~

Formatted: Font: Not Bold

Cronbach’s alpha, which measures question reliability, for the overall knowledge and attitude was 0.941 (range: 0.775–0.913) and 0.824 (range: 0.791–0.822), respectively. Cronbach’s alpha for practice and behavior ranged from low (0.257) for physical activity questions to high (0.929) for blood pressure ~~measurements~~ measurement.

Twelve local female enumerators (two per ward) conducted the interviews. All enumerators had completed at least grade 10 in school. Supervisors included two public health graduates: a field coordinator who holds both master’s and PhD degrees in public health; two PhD students including one author (AV).

Scoring knowledge, attitude, and practice/behavior questions

To compare and correlate the three components, we scored responses regarding KAP/behavior for cardiovascular health. The maximum possible score for KAP/behavior was 53, 70, and 25 respectively. See Supplement for further detail.

Physical measurements

Enumerators recorded anthropometric measurements (i.e., weight, height, and waist and hip circumferences) according to STEPs guidelines.²⁴ Weight was measured to one decimal place using a Microlife BR-9201 Weighing Scale (Microlife AG Swiss Corporation, Widnau, Switzerland). Height and waist and hip circumferences were measured using non-stretchable tapes (Jonson Tapes Ltd, New Delhi, India). Blood pressure was determined digitally by Microlife BP 3API-3E (Microlife AG Swiss Corporation, Widnau, Switzerland). Enumerators

Formatted: None, Space After: 0 pt

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

recorded three readings for each respondent, taken at 5-minute intervals. We averaged the readings to attain the final blood pressure measurements. Respondents with known history of hypertension (diagnosed cases) and those diagnosed during the study according to Joint National Committee -VII⁴¹ were termed as having “hypertension.”

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Data management and analysis

Supervisors checked all completed questionnaires and, if necessary, asked enumerators to revisit households. We outsourced data entry to a team of data entry operators, public health graduates by qualification. Data was coded and entered in Epidata version 2.1 and analyzed with SPSS version 17.0 (IBM, Armonk, New York, USA) and STATA version 10.0 (StataCorp, Texas, USA). Demographic variables were taken as independent variables; other variables, including KAP components, were analyzed as dependent variables. Categorical data is presented as percentages, and continuous data is presented as mean (standard deviation). KAP scores are presented as median score percents. We used the Chi-square test to compare proportions and the Kruskal–Wallis test to compare medians. $P < 0.05$ was considered statistically significant.

Formatted: Font: Not Bold

Ethical considerations

The Nepal Health Research Council and the Institutional Review Board of Kathmandu Medical College approved this study. Following a pattern established at the outset of JD-HDSS, we consulted local leaders and sought their informal permission for HDSS activities. In addition, we periodically consulted with health and administrative authorities in both villages. All participants gave informed consent, and privacy was maintained during the interview process. To maintain confidentiality, all data were secured in the HDSS office. Computer data were accessible only to the research team. Although respondents received no monetary incentives, we provided a referral that included discounted consultation fees at Kathmandu Medical College or Nepal Medical College for self-reported illness and newly diagnosed hypertension.

Formatted: None, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Font: Bold

RESULTS

Formatted: None, Space After: 0 pt

We surveyed 789 of 840 households in randomly selected six wards of Duwakot and Jhaukhel (non-response rate = 6.07%). After omitting incomplete questionnaires, we included 777 individuals in the analysis. Except for missing recordings for height, weight, or blood pressure, the data reported here cover all 777 respondents.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

Demographic characteristics and cardiovascular risk factors variables

Table 1 describes the sex-wise demographic characteristics of the respondents. Seventy percent of all participants were female, and 25% lacked formal education. The median age (IQR) for male and female respondents was 40 (33-49) years and 38 (32-46) years, respectively. Two third of the female respondents were housewives and 3 out of 5 male were either serviceholders or self employed. We have shown previously that 59.2% belonged to the upper lower class.²⁰

Table 1: Demographic characteristics and risk factor status of the study population- (n=777)

Demographic variables (n=777)	Male (n=229) Number (%)	Female (n=548) Number (%)	Total (n=777) Number (%)
Age (years)			
25-34	69 (26.8)	188 (73.2)	257 (100)
Age (years)		202 (72.1)	280 (100)
25-34	30.178 (27.9)	34.3 (36.9)	33.1 (36.0)
35-44	34.1	28.8	30.9
45-59	35.8		
45-59	82 (34.2)	158 (65.8)	240 (100)
Caste/Ethnicity			
Ethnic group	85 (28.8)	209 (71)	294 (100)
Brahmin	37.1	38.1)	37.8
Chhetri	24.0	25.4	25.0
Newar	29.3	23.2	25.0
Minorities	9.6	13.3	12.2
Chhetri	55 (28.4)	139 (71.6)	194 (100)
Newar	67 (34.5)	127 (65.5)	194 (100)
Others*	22 (23.2)	73 (76.8)	95 (100)

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None, Space After: 0 pt

Formatted: None, Space After: 0 pt, Line spacing: 1.5 lines

Split Cells

Formatted Table

Formatted: Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: Not Bold

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Split Cells

Formatted Table

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Tab stops: 0.13", Left

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted Table

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Tab stops: 0.13", Left

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013**Education-**

No formal schooling	27 (12.9)	182 (87.1)	209 (100)
Primary school (Grade:1-5)	74 (30.1)	172 (69.9)	246 (100)

Education*

	74 (37.4)	124 (62)	
No formal schooling	11.8	33.2	26.9 ¹⁹⁸
Primary school	32.3	31.4	(100)
Secondary school (Grades: 6-10)	32.3	22.6)	31.7
High school or more	23.6	12.8	25.5

High secondary and above (Grade:11 and above)Occupation†

	54 (43.5)	-70 (56)	124 (100)
Employee	30.6	8.4	14.9
Self-employed	32.4	6.6	13.9
Housewife	21.0	14.2	16.3
Agriculture	17.0	3.5)	17.5
Other			

Major Occupation§

Service (government and private)	-70 (60.3)	-46 (39.7)	116 (100)
----------------------------------	------------	------------	-----------

Self-employed Cardiovascular risk factors§

	-72 (66)		
Smoking (current) (n=777)	33.5	14.7-36	20.1
Smokeless tobacco (current) (n=777)	20.2	(33-	8.3
Alcohol drinkers (ever) (n=777)	36.7	3)3	20.7
Alcohol drinkers (current) (n=777)	34.5)	14.1	19.0
Low physical activity (n=777)	38.3	12.6	43.3 ¹⁰⁸
Hypertension (known and new) (n=670)	20.6	45.2	(100)
Self-reported high blood sugar (n=777)	4.8	22.1	21.6
Overweight/obese (n=762)	25.0/5.4	3.5	3.9
Increased waist circumference (cm) (n=773)	21.6	31.6/11.2	29.7/9.4
Increased waist/hip ration (n=750)	57.9	56.6	46.3

Agriculture	-48 (38.1)	-78 (61.9)	126 (100)
-------------	------------	------------	-----------

15

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted Table

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Tab stops: 0.13", Left

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Tab stops: 0.13", Left

Formatted: Font color: Black

Formatted Table

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Tab stops: 0.13", Left

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Housewife ^{††}	N/A	369 (100)	369 (100)
Others [†]	39 (67.2)	19 (32.8)	58 (100)

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

[†]Other caste/ethnicity includes Rai, Magar, Tamang, Dalit, Gurung, Mandal, Chaudary, Pariayar, Purkutti. ^{††}The definition Note: Classification of occupation was adopted from ethnic groups is based on the Nepal Non-Communicable Disease Survey, 2007 National Central Bureau of Statistics.⁴¹

*Primary and modified secondary schooling indicate education up to the local context.⁴² ^{††}A woman involved in her own household activities like cooking, washing, cleaning etc but does not earn money. [†]Other occupation includes grade 4 and 10, respectively.

[†]Employee describe individuals working for governmental or nongovernmental organizations. "Other" occupations include students, unemployed individuals, (able or unable to work), retired, and non-paid workers.

Formatted: Space After: 0 pt, Line spacing: 1.5 lines


Figure 1 shows the sex wise prevalence of cardiovascular risk factors in the study population.

Overall, three in ten respondents was a past or current smoker and one in five respondents had consumed at least one alcoholic beverage in the previous month. More than 40% of the respondents reported low physical activity. Prevalence of known and newly diagnosed hypertension was 21.6%. As measured by BMI, the prevalence of overweight and obesity was 29.7% and 9.4%, respectively. Abdominal obesity (i.e., increased waist circumference and waist hip ratio) was 46.3% and 56.3%, respectively. There was statistically significant difference in the prevalence rate among males and females in smoking, smokeless tobacco use, alcohol intake, overweight/obesity and increased waist circumference.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Figure 1: Sex-wise distribution of cardiovascular risk factor status in the study population.



Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: (Default) Times New Roman, 12 pt

m=number of male, f=number of female, # p<0.001, ## p<0.01 (chi-square tests).^{23,24} Risk factor categories are based on the WHO-NCD Risk Factor STEPS Survey manual.^{23,24} Current smokers included those who responded “yes” to “Do you smoke?” Past smokers included those who replied “yes” to “Did you ever smoke in the past?” “Ever drinkers” included respondents who had “consumed a drink that contained alcohol ever in lifetime,” and current drinkers included respondents who had “consumed alcohol within the previous month.” Percentage of drinking frequency represents current drinkers who consumed at least one standard drink per occasion. Blood pressure data exclude respondents who did not submit to all three readings. Likewise, data for body mass index, waist circumference, and waist-hip ratio exclude respondents whose weight, height, and waist and/or hip measurements were not taken. Increased waist circumference includes waist measurements of ≥ 80 cm (females) and ≥ 90 cm (males); increased waist-hip ratio is ≥ 0.85 (females) and ≥ 0.90 (males).

Formatted: Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: None, Space After: 0 pt

Formatted: Font: Not Bold

Knowledge towards

High burden of risk factors related to cardiovascular disease health

Table 1 also shows the prevalence of cardiovascular risk factors in the study population. One in four respondents was a past or current smoker. One in five respondents had consumed at least one alcoholic beverage in the previous month. Both males and females consumed fruit 2.6 ± 1.98 and 3.3 ± 2.17 days/week, respectively. Likewise, males and females consumed vegetables

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

5.33±1.69 and 5.64±1.55 days/week, respectively. More than 40% of the respondents reported low physical activity. Prevalence of known and newly diagnosed hypertension was 21.6%. As measured by BMI, the prevalence of overweight and obesity was 29.7% and 9.4 %, respectively. Abdominal obesity (i.e., increased waist circumference and waist-hip ratio) was 46.3% and 56.3%, respectively. Knowledge regarding causes and warning signs (Table 1).

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Poor knowledge of cardiovascular health

Cause of heart disease

Formatted: None, Space After: 0 pt

Table 2 shows When seeking unprompted responses for the perceived causes of heart diseases. The responses ranged, enumerators asked participants to spontaneously name the causes of CVD (Table 2). Respondents showed low overall knowledge of the factors that cause CVDs, ranging from 1.0% for (high blood sugar) to 29.1% for 2% (smoking. The median percentage scores for). Generally, males and knew more than females were 13.8 and 10.4 %, respectively (p; younger <0.001, Mann-Whitney U test). Younger respondents (<35 years) knew more than older participants (median 13.7% vs. 10.3%, p=0.01). Respondents belonging to Brahmin, Newer and Chhetri ethnicities had less knowledge than those in the other caste/ethnicity group (median 13.8% vs. 6.9%, p=0.001) (Table 1). The respondents with high school or more education and working in government or non-government offices knew more ones; ethnic minorities knew less; and better-educated individuals and governmental or nongovernmental employees were better informed than those with less education and in other employment situations (median 17.2% vs. 10.34%, p=0.001).

Table 2: Percentage of respondents citing various causes of cardiovascular heart disease and warning signs of heart attack. (n=777)

Formatted: Font color: Black

Formatted: Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font color: Black

Formatted: Font: Not Bold

Causes of Heart Attack	Male (n=229) Number (%)	Female (n=548) Number (%)	Total (n=777) Number (%)	P-value[*]
Hypertension	43 (18.8)	60 (10.9)	103 (13.3)	<0.01
High blood sugar	3 (1.3)	5 (0.9)	8 (1.0)	0.06
High cholesterol	47 (20.5)	74 (13.5)	121 (15.6)	0.01
Physical inactivity	16 (7.0)	22 (4.0)	38 (4.9)	0.08
Overweight	13 (5.7)	29 (5.3)	42 (5.4)	0.82
Cigarette/Bidi Smoking	80 (34.9)	147 (26.8)	227 (29.1)	0.02

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Smokeless tobacco	19 (8.3)	-26 (4.7)	-45 (5.8)	0.06
Excessive alcohol	72 (31.4)	131 (23.9)	203 (26.1)	0.03
Excess stress	37 (16.2)	-74 (13.5)	111 (14.3)	0.34
Warning signs	Male (n=93)[‡] Number (%)	Female (n=222)[‡] Number (%)	Total (n=315)[‡] Number (%)	P-value[‡]
Difficulty in breathing	17 (11.1)	41 (15.1)	58 (13.6)	0.25
Pain in the chest	30 (19.6)	57 (21.0)	87 (20.5)	0.74
Pain or numbness in the arms	-0 (0)	-3 (1.1)	-3 (0.7)	-
Pain in the teeth or jaw	-0 (0)	-1 (0.4)	-1 (0.2)	-
Excessive sweating	15 (9.8)	30 (11.0)	45 (10.6)	0.69
Vomiting tendency	-1 (0.7)	-6 (2.2)	-7 (1.6)	0.65
Dizziness or light headedness	24 (15.7)	40 (14.7)	-64 (15.1)	0.78
Loss of consciousness	55 (35.9)	86 (31.6)	141 (33.2)	0.36

Numbers

	<u>Hypertension</u>	<u>High blood sugar</u>	<u>High cholesterol</u>	<u>Physical inactivity</u>	<u>Overweight</u>	<u>Smoking</u>	<u>Smokeless tobacco</u>	<u>Excessive alcohol</u>	<u>Excess stress</u>
Sex									
<u>Male</u>	<u>18.8*</u>	<u>1.3</u>	<u>20.5*</u>	<u>7.0</u>	<u>5.7</u>	<u>34.9*</u>	<u>8.3</u>	<u>31.4*</u>	<u>16.2</u>
<u>Female</u>	<u>10.9*</u>	<u>0.9</u>	<u>13.5*</u>	<u>4.0</u>	<u>5.3</u>	<u>26.8*</u>	<u>4.7</u>	<u>23.9*</u>	<u>13.5</u>
Age (years)									
<u>25-34</u>	<u>14.8</u>	<u>1.6</u>	<u>17.5</u>	<u>5.4</u>	<u>6.2</u>	<u>31.1</u>	<u>6.2</u>	<u>28.4</u>	<u>16.0*</u>
<u>35-44</u>	<u>13.2</u>	<u>0.4</u>	<u>17.1</u>	<u>5.7</u>	<u>6.1</u>	<u>26.8</u>	<u>5.0</u>	<u>25.0</u>	<u>17.1*</u>
<u>45-59</u>	<u>11.7</u>	<u>1.3</u>	<u>11.7</u>	<u>3.3</u>	<u>3.8</u>	<u>30.0</u>	<u>6.3</u>	<u>25.0</u>	<u>9.2*</u>
Ethnicity									
<u>Brahmin</u>	<u>14.3*</u>	<u>1.0</u>	<u>21.1[#]</u>	<u>7.5*</u>	<u>5.8</u>	<u>29.3</u>	<u>8.2</u>	<u>25.2</u>	<u>12.9</u>
<u>Chhetri</u>	<u>19.1*</u>	<u>1.0</u>	<u>16.5[#]</u>	<u>4.6*</u>	<u>6.7</u>	<u>32.5</u>	<u>3.6</u>	<u>25.8</u>	<u>20.1</u>
<u>Newar</u>	<u>10.8*</u>	<u>.5</u>	<u>9.8[#]</u>	<u>3.6*</u>	<u>5.7</u>	<u>27.8</u>	<u>6.2</u>	<u>28.4</u>	<u>10.8</u>
<u>Minorities</u>	<u>3.2*</u>	<u>2.1</u>	<u>8.4[#]</u>	<u>0.0*</u>	<u>1.1</u>	<u>25.3</u>	<u>2.1</u>	<u>25.3</u>	<u>13.7</u>
Education									
<u>Primary school</u>	<u>13.0*</u>	<u>0.8</u>	<u>14.2[#]</u>	<u>5.3[#]</u>	<u>4.5*</u>	<u>30.5</u>	<u>8.1</u>	<u>27.6</u>	<u>15.0*</u>

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek—August 7, 2013

<u>Secondary school</u>	<u>15.7*</u>	<u>1.5</u>	<u>18.7[#]</u>	<u>8.6[#]</u>	<u>8.6*</u>	<u>29.8</u>	<u>3.0</u>	<u>27.8</u>	<u>16.7*</u>
<u>High school or more</u>	<u>21.0*</u>	<u>1.6</u>	<u>27.4[#]</u>	<u>4.8[#]</u>	<u>8.9*</u>	<u>33.9</u>	<u>8.1</u>	<u>26.6</u>	<u>18.5*</u>
<u>No formal schooling</u>	<u>6.7*</u>	<u>0.5</u>	<u>7.2[#]</u>	<u>1.0[#]</u>	<u>1.4*</u>	<u>24.4</u>	<u>4.3</u>	<u>22.5</u>	<u>8.6*</u>
<u>Occupation</u>									
<u>Employee</u>	<u>26.7[#]</u>	<u>0.9</u>	<u>32.8[#]</u>	<u>10.3</u>	<u>8.6</u>	<u>30.2</u>	<u>8.6</u>	<u>31.9</u>	<u>23.3*</u>
<u>Self-employed</u>	<u>17.6[#]</u>	<u>2.8</u>	<u>14.8[#]</u>	<u>1.9</u>	<u>7.4</u>	<u>29.6</u>	<u>2.8</u>	<u>25.9</u>	<u>15.7*</u>
<u>Housewife</u>	<u>10.6[#]</u>	<u>1.1</u>	<u>11.7[#]</u>	<u>3.3</u>	<u>5.4</u>	<u>25.7</u>	<u>3.5</u>	<u>23.8</u>	<u>13.0*</u>
<u>Agriculture</u>	<u>7.1[#]</u>	<u>0.0</u>	<u>14.3[#]</u>	<u>7.1</u>	<u>1.6</u>	<u>37.3</u>	<u>8.7</u>	<u>26.2</u>	<u>11.1*</u>
<u>Others</u>	<u>8.6[#]</u>	<u>0.0</u>	<u>10.3[#]</u>	<u>5.2</u>	<u>3.4</u>	<u>31.0</u>	<u>13.8</u>	<u>29.3</u>	<u>8.6*</u>
<u>Overall</u>	<u>13.3</u>	<u>1.0</u>	<u>15.6</u>	<u>4.9</u>	<u>5.4</u>	<u>29.2</u>	<u>5.8</u>	<u>26.1</u>	<u>14.3</u>

Note: Figures are based on participants' unprompted spontaneous participant responses. Only the table includes only established CVD risk factors included and excludes responses citing that cited other non-established causes (e.g., food hygiene, air pollution, etc.) excluded. *Obtained from chi-square test. # Respondents who did not know any warning sign were excluded from analysis. Calculated with Chi-square test, p-values compare all categories in the variables.

Table 2 also describes the warning signs* P-value <0.05; # p-value <0.01

Knowledge of heart attack as perceived by the respondents—symptoms

Around 60% of respondents did not know any of the heart attack signs. Among those who were familiar symptoms compared with those signs, nearly 50% of respondents knew only one sign, 37% 20% who knew any two signs, 11.42% knew any three signs and 2.53% knew four signs. The proportions of each warning sign told by 2–4 symptoms. This ratio was uniform across all the male and female respondents were not statistically different (p>0.05). The proportions of knowledge about warning signs were not statistically different in terms of age ($\chi^2=3.39$, p=0.49), caste/ethnicity ($\chi^2=6.25$, p=0.4), and education ($\chi^2=5.50$, p=0.53) demographic variables (Table 3). Importantly, only 20 less than 25% of respondents identified chest pain—the most prominent and important indicator of heart attack—as a symptom of heart attack (data not shown). This knowledge decreased with age (p=0.013) and level of education (p=0.055). Brahmin respondents (p<0.001) and governmental or nongovernmental employees (p=0.021) had slightly better knowledge about chest pain.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Space After: 0 pt

Formatted: Font: 10 pt

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Knowledge on action **Table 3: Correctly reported warning signs of heart attack (n=777)**

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

	<u>Signs reported (number)</u>			<u>P-value</u>
	<u>0</u> <u>(%)</u>	<u>1</u> <u>(%)</u>	<u>2-4</u> <u>(%)</u>	
<u>Sex</u>				
<u>Male</u>	<u>59.4</u>	<u>18.3</u>	<u>22.3</u>	<u>0.605</u>
<u>Female</u>	<u>59.5</u>	<u>20.8</u>	<u>19.7</u>	
<u>Age (years)</u>				
<u>25-34</u>	<u>63.4</u>	<u>16.7</u>	<u>19.8</u>	<u>0.494</u>
<u>35-44</u>	<u>57.1</u>	<u>21.4</u>	<u>21.4</u>	
<u>45-59</u>	<u>57.9</u>	<u>22.1</u>	<u>20.0</u>	
<u>Ethnicity</u>				
<u>Brahmin</u>	<u>56.8</u>	<u>22.1</u>	<u>21.1</u>	<u>0.395</u>
<u>Chhetri</u>	<u>62.9</u>	<u>18.0</u>	<u>19.1</u>	
<u>Newar</u>	<u>59.3</u>	<u>17.0</u>	<u>23.7</u>	
<u>Others</u>	<u>61.1</u>	<u>24.2</u>	<u>14.7</u>	
<u>Highest education</u>				
<u>No formal schooling</u>	<u>60.3</u>	<u>22.0</u>	<u>17.7</u>	<u>0.493</u>
<u>Up to primary school</u>	<u>58.1</u>	<u>18.7</u>	<u>23.2</u>	
<u>Up to secondary school</u>	<u>60.1</u>	<u>22.2</u>	<u>17.7</u>	
<u>High school or more</u>	<u>59.7</u>	<u>16.1</u>	<u>24.2</u>	
<u>Main job</u>				
<u>Employee</u>	<u>55.2</u>	<u>19.0</u>	<u>25.9</u>	<u>0.687</u>
<u>Self-employed (self)</u>	<u>57.4</u>	<u>20.4</u>	<u>22.2</u>	
<u>Housewife</u>	<u>59.1</u>	<u>21.4</u>	<u>19.5</u>	
<u>Agriculture</u>	<u>65.1</u>	<u>15.9</u>	<u>19.0</u>	
<u>Others</u>	<u>62.1</u>	<u>22.4</u>	<u>15.5</u>	

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Overall	59.5	20.1	20.5
----------------	-------------	-------------	-------------

Note: Warning signs were based on a list provided by American Heart Association.⁴²

P-values were calculated using the Chi-square test and compare all categories in the variables.

Action in case of a heart attack

When asked ~~about~~ what action ~~the respondents~~they would take for a suspected heart attack, ~~more~~ than 20% gave no response and 75% (81.2% male and 73% female, $\chi^2=4.05$, $p=0.04$) ~~75% of respondents~~ said they would immediately take the ~~affected~~ person to a hospital. About 3% said they would try home therapy, consult a traditional healer, or stay home and wait for the symptoms to go away. More than 20% gave no response.

Knowledge on Heart-healthy foods for heart diseases.

When asked to identify a particular heart-healthy food, most respondents identified green leafy vegetables (94.6%) and fruit (92.1%). ~~Furthermore, 22% thought salty food like chips and~~ around ~~Around~~ 7% thought fried food or saturated fat (e.g., ghee) are healthy foods. ~~Nearly 22% thought salty food (e.g., chips) is heart-healthy.~~

Attitudes and practice towards cardiovascular Heart disease prevention

After reading aloud a list of specific actions that help prevent CVD, enumerators asked respondents to comment if they thought a particular action would help. Actions included quitting smoking; getting physical exercise; losing weight; reducing intake of dietary fat, animal products, and salt; reducing stress; and maintaining healthy blood pressure. A majority of respondents thought these actions are helpful, ranging from 86.1% for weight reduction to 92.5% for smoking cessation.

Attitude toward heart health in an urbanizing community

Enumerators asked respondents about their attitude toward different issues of heart health (Table 3 describes opinions reflecting attitudes towards various aspects on heart diseases-4). Regarding lifestyle and preventive measures, ~~2572.9% of males and 3068.0%-females perceived~~ did not perceive themselves ~~to be~~ at risk of CVD. ~~More and more than half of the respondents did not want to change their current lifestyle (64.6% males vs. 55.4% = 63.3%; females, $p<0.01$). Nearly~~

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None, Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: None, Space After: 0 pt

Formatted: Font color: Black

Formatted: Font: Not Bold, Italic, Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: None, Space After: 0 pt

Formatted: Font: Not Bold, Font color: Black

Formatted: Font: Not Bold, Italic, Font color: Black

Formatted: Font: Bold

Formatted: None, Space After: 0 pt

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

23% = 53.1%). Seventy-five percent said they did not eat more and would like to cut down; approximately 82% considered than they need; 80% did consider themselves to be overweight—the perceived proportion being significantly higher among males (p<0.05); nearly 32% believe that they do not have time to take care themselves; and three fifths did not think that changing their behavior would lower their cardiovascular risk. Further, 25.352.4% of males and 27.260.8% of females thought that God or a higher power determines their health.

Table 3: Responses to attitude-related questions on cardiovascular disease/heart health

SN	Items	Male		Female		Total		P-value [‡]			
		Number (%)	Number (%)	Number (%)	Number (%)	Number (%)	Number (%)	Number (%)	Number (%)		
	Attitude items on heart health	Strongly disagree		Somewhat disagree		Somewhat agree		Strongly agree		Don't know/ Can't say	
		M	F	M	F	M	F	M	F	M	F
1	I don't perceive myself to be at risk for heart disease (m=22, f=54, n=76)*	58 (25.8)	167 (30.9)	225 (29.0)	37.2	15.7	21.9	9.6	8.6	1.7	1.5
2	I don't want to change my lifestyle	148 (64.6)	98 (54.4)	17.9	446 (57.2)	30.1	20.8	33.2	32.3	1.3	1.3

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None, Space After: 0 pt

Formatted: Font: Not Bold

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Centered, Right: 0.16", Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted Table

Inserted Cells

Formatted: Font: Times New Roman, 12 pt, Italic, Font color: Text 1

Formatted: Font: Times New Roman, 12 pt, Italic, Font color: Text 1

Formatted ... [2]

Formatted ... [3]

Formatted ... [1]

Deleted Cells

Formatted: Font color: Black

Formatted ... [4]

Formatted ... [5]

Inserted Cells

Formatted ... [6]

Formatted: Font color: Black

Inserted Cells

Inserted Cells

Inserted Cells

Inserted Cells

Formatted: Font color: Black

Inserted Cells

Formatted: Font color: Black

Formatted ... [7]

Formatted ... [8]

Formatted ... [9]

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

	for my health (m=22 9/f=54 8)											
5	I don't think changing my behavior will reduce my risk of developing heart disease (m=22 9/f=52 2)	145 (63.3) 3.1	312 (56.9) 15.0	457 (58.2) 8	0.23 4	38.4	32.5	24 9	24.5	1.7	4.7	
6	I don't have time to take care of myself (m=22 9/f=54 8)	44.1	39.4	22.3	29.9	13.5	14.4	18 3	-77 (33.14) 6	168 (30.1) 7	245 (31.5) 16	0.4 2
7	God or	25.8	23.7	20.1	13.5	-58	22.3	14	207	1.7	2.0	59

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font color: Black

Inserted Cells

Inserted Cells

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: ... [11]

Inserted Cells

Inserted Cells

Inserted Cells

Formatted: ... [12]

Deleted Cells

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: ... [13]

Formatted: ... [14]

Formatted: ... [15]

Inserted Cells

Formatted: ... [16]

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

8											
9	Greater	200	67.3	31.0	22.4	492	692	0.319	0.5	4.8	6.0
10	access	(87.5)				(89.7)	(89.1)				
11	to	6.3				0	3.6				
12	indoor										
13	and										
14	outdoor										
15	public										
16	recreati										
17	onal										
18	facilitie										
19	s										
20	(m=22										
21	9/f=54										
22	8)										
23											
24											
25											
26											
27											
28											
29											
30	Smokin	209	92.0	3.9	51.5	0.9	0.9	724	0.897	1.3	2.2
31	g bans	(96.9)			(96.4)			(96.1)			
32	(m=21	2.1			2			7			
33	8/f=53										
34	5)										
35											
36											
37											
38	More	214	87.0	9.6	8.6	483	697	0.389	0.4	13	31
39	awaren	(96.1)				(90.3)	(98.0)				
40	ess	85.2									
41	progra										
42	ms for										
43	healthy										
44	heart										
45											
46											
47	(m=22										
48	6/f=53										
49	6)										
50											
51											
52	Availa	217	524	7.0	7.7	1.7	0.7	741	0.027	1.3	3.3
53											

- Formatted ... [17]
- Formatted ... [18]
- Formatted ... [20]
- Formatted ... [21]
- Formatted ... [22]
- Inserted Cells ... [26]
- Formatted ... [24]
- Inserted Cells ... [27]
- Inserted Cells ... [28]
- Formatted ... [32]
- Formatted ... [29]
- Formatted ... [30]
- Formatted ... [25]
- Formatted ... [31]
- Formatted ... [23]
- Inserted Cells ... [40]
- Inserted Cells ... [41]
- Formatted ... [33]
- Formatted ... [34]
- Formatted ... [36]
- Formatted ... [38]
- Formatted ... [42]
- Formatted ... [44]
- Formatted ... [39]
- Formatted ... [43]
- Formatted ... [37]
- Formatted ... [35]
- Formatted ... [48]
- Inserted Cells ... [50]
- Formatted ... [51]
- Formatted ... [54]
- Inserted Cells ... [55]
- Formatted ... [45]
- Formatted ... [46]
- Formatted ... [52]
- Formatted ... [53]
- Formatted ... [49]
- Formatted ... [47]
- Formatted ... [56]
- Formatted ... [57]
- Formatted ... [58]
- Inserted Cells ... [59]
- Inserted Cells ... [60]
- Formatted ... [61]
- Formatted ... [62]
- Formatted ... [19]

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

2	bility	(968)	(9887)				(970.9)				
3	of	2.1	6								
4	health										
5	facilitie										
6	s to										
7	detect										
8	and										
9	treat										
10	heart										
11	disease										
12	s										
13	(m=22										
14	f=53										
15)										
16	Local	486	64.65	17.0	708	p<0.0	2.9	1.7	0.4	13.1	11.5
17	volunte	(826	22		(932	13.1					
18	ers that	5.1	(98.5)		0.6)						
19	shall										
20	try to										
21	change										
22	any										
23	adverse										
24	health										
25	behavi										
26	or										
27	(m=22										
28	f=53										
29)										

m=number of males, f=number of females. Questions based on 5-scale Likert questions (strongly disagree, somewhat disagree, I do not know, somewhat agree and strongly agree (question 1-7)/very helpful; somewhat helpful; I do not know; not very helpful; not at all helpful (question 8-13). We classified five options into three categories as 'agree/helpful', 'disagree/not helpful' and 'I do not know'. Only 'strongly agree' or 'very helpful'

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Inserted Cells

Formatted: Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and

Inserted Cells

Formatted: Font color: Black

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

and 'agreed/somewhat helpful' answers were merged and presented in the table and 'I do not know/cannot say' were excluded from analysis. *Obtained from chi-square test. [§]'I do not know/cannot say' answers were excluded from analysis. [¶]Cannot be computed due to expected frequency less than 5.

Table 3 also describes the respondents²Note: Figures indicate percentage of respondents among males (M) and females (F).

Next, enumerators asked for respondent opinions on various preventive aspects of heart health (e.g., improved access to fruits, vegetables, and other healthy foods) (Table 4). Most thought that would be helpful. Almost all respondents believed that smoking bans or increased access to fruits, vegetables, and other healthy foods would be very helpful. Nearly 9 in 10 About two thirds of all respondents thought that greater access to indoor and outdoor public recreational facilities would be very helpful. Four percent of respondents did not believe smoking bans and one quarter thought it would be somewhat helpful to reduce heart diseases. Eight in 10 male respondents believed that. Around 10% of females seemed unsure whether local health volunteers should try/trying to change any adverse health behaviors while nearly 9 in 10 females believed the same (p<0.01). community's health practice/behavior would be useful.

Practice towards/behavior in relation to heart diseasesdisease

Half of all theEnumerators elicited practice- and behavior-related responses in two ways. First, they asked WHO STEPs-based questions about four major behavior-related risk factors (i.e., tobacco smoking, alcohol consumption, physical activity, and fruit and vegetables) (Table 1). Second, they asked respondents if they had participated in any heart-friendly practice/behavior in the previous year (e.g., blood pressure, blood sugar, or other diagnostic tests; tobacco cessation; increased physical activity; weight loss). Half of all respondents had their blood pressure measured and 10% reported getting a diagnostic test for CVD in the previous year. About 60% attempted to reduce their consumption of unhealthy foods and 25% tried to lose weight or increase physical activity. Half of the smokers said they had attempted to quit smoking.

When enumerators asked the respondentswhy responders who reported one or more positive health behaviors the main reason for trying/ried to change their lifestyle or practice/behavior, the reasons given were: participants responded that they wanted to feel better (70%); had seen, read, or heard information (65.4%); did it for their family (45.9%); or were encouraged to do so by a healthcare professional (41.3%), family member (39.0%), or friend (31.8%). Other reasons for

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: Bold

Formatted: None, Indent: Left: 0", First line: 0", Space After: 0 pt

Formatted: Font: Bold

Formatted: Font: Times New Roman, 12 pt, Bold

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

changing lifestyle or practice/behavior included wanting to avoid taking medications (20.6%); a family member, relative, or friend who developed CVD, became ill, or died (16.5%); and developing possible cardiovascular symptoms (14.8%).

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font color: Black

Comparison between level of knowledge, attitude and behavior/practice

Overall, the average prompted knowledge of heart diseases (median 79.3%) was 11.5% lower than average attitudes (median 67.8%) and nearly 31.1% higher than average practices (median 48.2%). Next, average level of attitudes score was nearly 19.6% higher than practices. Similarly, average spontaneous knowledge of heart diseases (median 10.3%) was 57.5% lower than average attitudes (median 67.8%) and nearly 40% lower than average practice (median 48.18%).

Table 4 describes level of KAP based on quintile scores. Regarding knowledge, 20.4% of the respondents had highly insufficient knowledge, 23.2% had insufficient knowledge, 22.6% had sufficient knowledge, 16.5% had satisfactory knowledge and 17.2% had highly satisfactory knowledge. Similarly, attitude levels were as follows: highly insufficient 20.6%, insufficient 26.9%; sufficient 15.4%; satisfactory 11.2% and highly satisfactory 19.8%. Regarding the behaviour/practices, it was highly insufficient in 12.1% of the respondents, insufficient in 24.3%; sufficient in 22.7%; satisfactory in 27.5% and highly satisfactory in 13.4%. Among those with highly insufficient knowledge, **KAP scores and demographic variables**

Median percent scores were similar across the various demographic variables (Table 5). Males, younger people, or better-educated respondents scored 2%–8% higher than females, older respondents, and the less educated. Similarly, median percent scores for attitude tended to decrease about 2% in older respondents, Newars, and those without formal schooling. Behavioral median percent scores were 4%–8% higher among younger and more-educated respondents, Brahmins, Chhetris, and government employees or housewives (Table 5).

Table 5: Median percent score of knowledge, attitude, and practice/behavior according to demographic variables

~~20.3%~~ of them had highly satisfactory attitudes and 6.9% had highly satisfactory practice.

Formatted: Font color: Black

Conversely, among the respondents with highly satisfactory knowledge, 14.7% had highly satisfactory attitude and 19.5% and 13.9% had satisfactory and highly satisfactory practices respectively. Likewise, one in ten respondents with satisfactory knowledge had satisfactory

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek—August 7, 2013

attitude while two in ten had highly satisfactory attitude. Nearly four out of ten had satisfactory/highly satisfactory behavior. As the level of knowledge increased, the highly satisfactory practice also increased from 6.9% to 13.4%. However, when the level of knowledge increased, the highly satisfactory attitude decreased from 20.3% to 14.7%.

Table 4: Percentage distribution of level of knowledge, attitude and behavior/practice regarding cardiovascular health.

		Knowledge (n=777) median percent score		P-value	Attitude (n=777) median percent score		P-value	Behavior median percent score		
I (n=180)	HH (n=160)	10.4 (n=209)	26.1 (n=209)	10.7 (n=120)	20.6 (n=134)	13.4 (n=154)				
HH (n=159)	26.0	27.6	13.8	12.5	20.3					
I (n=180)	17.9	28.4	19.1	12.5	22.2					
Suff (n=176) Sex		17.9			17.2	11.9	26.2			
Male		79.2		<0.001	74.29	48.00	0.660			
Female		77.4			74.29	48.00				
Sat (n=128)		20.8	33.7	15.8	6.9	23.8				
HS (n=134)		33.7	24.2	12.6	12.6	14.7				
Knowledge (n=777)		Behavior (n=777)								
		HH (n=94)	I (n=189)	Suff (n=176)	Sat (n=214)	HS (n=104)				
HH (n=159) Age (years)		16.4			21.4	25.8	6.9			
25-34		79.79			74.29	48.00				
35-44		79.24	0.005		74.29	44.00	0.018			
45-59		75.47			72.86	44.00				

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Inserted Cells

Formatted: Level 9, Space Before: 10 pt, After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Keep with next, Keep lines together, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Inserted Cells

Formatted Table

Formatted: Level 9, Space Before: 10 pt, After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Keep with next, Keep lines together, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Inserted Cells

Split Cells

Inserted Cells

Inserted Cells

Formatted Table

Formatted: Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted: Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted ... [64]

Formatted ... [65]

Formatted ... [66]

Formatted ... [63]

Formatted Table

Inserted Cells

Formatted: Font color: Black

Formatted ... [70]

Formatted ... [72]

Formatted ... [68]

Formatted ... [69]

Formatted ... [71]

Formatted ... [67]

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

Attitude (n=777)	Behavior (n=777)							
	HI	I	Suff	Sat	HS			
Suff (n=176)	11.4	20.5	20.5	31.2	16.5			
Sat (n=128)	10.9	18.8	33.6	19.5	17.2			
HS (n=134)	11.2	26.1	20.9	28.5	13.4			
HI (n=132)	15.0	23.8	20.0	28.1	13.1			
I (n=101)	12.0	22.0	18.7	30.1	17.2			
Suff (n=256) Ethnic group		10.0		25	28.3	12.5		
Brahmin			75.47		74.29		48.0	
Chhetri			79.24	<0.001	74.29	<0.001	0	0.030
Newar			79.24		72.14		48.0	
Other			79.24		74.29		0	
							44.0	
							0	
							44.0	
							0	
Sat (n=134) Education			13.8		17.2			
No formal schooling			75.47		72.86		44.0	
Primary school			77.35		74.29		0	
Secondary school			79.24	<0.001	9	0.001	44.0	<0.001
High school or more			83.02		74.29	7	0	1
					75.71		48.0	
							0	
							52.0	
							0	
HS (n=154) Occupat ion		10.4	20.8	33.8		11		
Employed			79.24		74.29		48.0	
Self-employed			79.24	<0.001	74.29	0.940	44.0	0.156
Housewife			79.24		74.29		0	
Agriculture			75.47		74.29		48.0	
Other			79.24				0	
							44.0	
							0	
							48.0	
							0	

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Inserted Cells

Formatted Table

Deleted Cells

Inserted Cells

Formatted: Font color: Black

Formatted: Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted: Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted: Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted: Left, Level 9, Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Keep with next, Keep lines together, Tab stops: 0.13", Left, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Formatted ... [73]

Deleted Cells

Formatted: Font color: Black

Formatted ... [76]

Formatted: Font color: Black

Formatted ... [78]

Formatted ... [75]

Formatted ... [74]

Formatted ... [77]

Deleted Cells

Deleted Cells

Deleted Cells

Inserted Cells

Inserted Cells

Formatted: Font color: Black

Formatted ... [81]

Formatted: Font: +Body (Calibri)

Formatted ... [80]

Formatted ... [79]

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Overall	<u>79.24</u>		<u>74.29</u>		<u>48.00</u>	
----------------	--------------	--	--------------	--	--------------	--

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

HI: Highly insufficient; I: Insufficient; Suff: Sufficient; Sat: Satisfactory; HS: Highly Satisfactory. The total score was 53, 64 and 25 for KAP, respectively. Then, the quintile scores were obtained. The quintile score (1st-4th) was for prompted knowledge: 34, 40, 43, 45; for attitudes: 33, 37, 39, 42; for behavior: 9, 11, 13, 15. Next, the levels of survey respondents on KAP were classified into the following five subdivisions based on quintile score: $\leq 20\%$ 'highly insufficient'; 21-40% 'insufficient'; 41-60% 'sufficient'; 61-80% 'satisfactory'; $> 80\%$ 'highly satisfactory'.⁴⁴ Median percentage scores for knowledge, attitude and practice were 79.3, 74.2 and 48 for all respondents.

Regarding attitude, among those with the level of attitudes of highly insufficient, 41.2% had satisfactory/highly satisfactory behaviour/practice. When the level of attitude was highly satisfactory, 35% had satisfactory/highly satisfactory practice. Similarly, when the attitude was sufficient, 25% had insufficient practice and nearly 41% had satisfactory practice. When the level of attitude increased, the proportion of those with highly satisfactory practice remained unchanged.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

A comparison of male and female respondents showed no difference in level of knowledge ($\chi^2=4.4$, $p=0.4$) attitudes ($\chi^2=3.6$, $p=0.5$) and practice ($\chi^2=2.1$, $p=0.7$). However, there were significant differences in the level of knowledge according to age ($\chi^2=18.1$, $p=0.02$), easte/ethnicity ($\chi^2=43.8$, $p<0.001$). Note: Knowledge scores were based on prompted responses. Calculated with the Kruskal-Wallis test, p-values compared all categories in the variables.

Correlation between knowledge, attitude, and practice/behavior scores

Scatter-plots (Figure 1a–1c) show the correlations between the three components. We observed a poor positive correlation between knowledge and attitude median scores (Pearson correlation: 0.025, $p=0.480$) [Figure 1a] and a negative correlation between attitude and behavior (Pearson correlation: -0.126, $p<0.001$) [Figure 1b]. However, we observed a positive correlation between knowledge and behaviour (Pearson correlation: 0.121, ($p=0.001$) [Figure 1c]).

($p<0.001$) and education level ($\chi^2=48.7$, $p<0.001$). In contrast, the level of attitude was not statistically significant for these variables ($p>0.05$). On the other hand, the level of practice was also statistically different according to age ($\chi^2=18.7$, $p=0.01$), easte/ethnicity ($\chi^2=21.1$; $p=0.004$) and education status ($\chi^2=45.7$, $p<0.001$).

Respondents <35 years of age were more likely to have highly satisfactory knowledge (37.3% vs. 29.2%) and practice (38.5% vs. 19.2%) than those >45 years. Brahmins (30.5%) were more likely to have highly satisfactory knowledge than Newars (27.6%), Chhetris (20.4%), and others (21.5%). Regarding highly satisfactory practice, the proportion was highest among Brahmins (41.3%), followed by Chhetris (30.8%), Newars (17.3%) and others (10.6%). Those who had no formal schooling exhibited higher proportion of highly satisfactory knowledge than those who had high school or more education (28.3% vs. 24.6%, but highly satisfactory practice is low among them (12.5% vs. 28.8%, data not shown).

DISCUSSION

The present study is the first to conduct a detailed analysis of the relationship between KAP/health literacy and cardiovascular health in a semi-urban setting near Kathmandu.

Cardiovascular risk factors are abundant in the study population

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font color: Black

Formatted: Font color: Black

Formatted: None, Space After: 0 pt

Formatted: None, Space After: 0 pt, Tab stops: Not at 0.2"

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Our respondents showed a high burden of behavioral and biological risk factors. ~~While the~~ The prevalence of tobacco smoking ~~was~~ and vegetable and fruit consumption were similar to ~~the~~ national ~~average, the~~ averages and alcohol consumption was lower.^{43,43} ~~National figures on tobacco smoking and alcohol consumption for Bangladesh, Nepal, India and Pakistan show that current smoking of any tobacco product (age standardized rate, both sexes) is 24%, 32%, 15% and 20%, respectively.⁴⁹ The total adult (recorded and unrecorded) per capita consumption of pure alcohol according to WHO available projected estimates for 2008 indicates that consumption in Bangladesh, Nepal, India and Pakistan is 0.17, 2.42, 2.69, 0.05 litres of pure alcohol, respectively, over a calendar year.⁴⁹ Numbers have been increasing in these countries over the last years.⁴⁹~~

In accord with our earlier review on obesity in Nepal, the effect of urbanization is evident in low physical activity and overweight/obesity that exceeds the national average.^{21,42,22,43} The prevalence of hypertension in our study population is similar to the national average (22%) but less by about one third than a study in Bhadrabas, a similar semi-urban Kathmandu community,^{43,44} most probably due to different age groups. ~~The 2008 national reported age-standardized level of hypertension in Nepal is similar to levels in Bangladesh but lower than in Pakistan (38.6% and 39.5%, respectively) and higher than in India (35.2%).⁴⁴ Self-reported prevalence of diabetes mellitus (3.9%) confirms the HDSS baseline figure reported earlier.^{20,21}~~

Knowledge ~~Poor knowledge~~ of cardiovascular health is limited

Our study population showed low overall knowledge of the causes of CVD. Unprompted responses suggested that respondents identified behavioral factors (e.g., smoking) more frequently than physiological causes (e.g., hypertension). Other studies have reported similar findings for unprompted³⁸ and prompted responses.^{33,34} Conversely, unhealthy diet, obesity, and family history were common responses in Taiwan.^{34,32} ~~In Vietnam, knowledge on the negative influence of tobacco smoking on stroke and heart attack is lower than general knowledge regarding health risks of active smoking and exposure to second hand smoke.⁴⁵~~

American women frequently identified obesity.³⁵ Many of our respondents also attributed CVD to environmental factors such as food hygiene and air pollution. Indeed, acute inflammatory reactions to air pollutants accelerate coronary atherosclerosis and trigger arrhythmia or

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal, None, Tab stops: Not at 0.2"

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

myocardial infarction.^{46, 45} Unlike the Strong Heart Study³⁹ in American populations, our results did not exhibit a higher probability of knowledge of a risk factor in persons who have that risk factor, demonstrating poor cardiovascular health literacy even among those already affected.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

Formatted: Font color: Black

Formatted: Font color: Black

Similar to Pakistan,^{45, 17} a large majority (60%) of our respondents could not identify even one heart attack symptom. However, similar to an earlier study in Nepal,^{46, 18} 6%–22% of individuals in our demographic subsets recognized chest pain as a symptom, which is less compared to 60% of Vietnamese Americans who spontaneously mentioned chest pain.^{47, 47} Importantly, two thirds of heart attack patients delayed their arrival at a Karachi hospital because they did not recognize heart attack symptoms.^{45, 17} Similar to a previous study,¹⁸ three quarters of our respondents believed they would immediately take a suspected heart attack patient to a hospital but one fifth did not answer the question or opted instead for home therapy or traditional healing. Combined with our earlier finding of not knowing heart attack symptoms, this lack of knowledge about appropriate action underscores the need to educate the Nepalese population about this issue. In contrast, 85% of Vietnamese Americans knew that they should immediately call 911 if they suspect a heart attack.^{45, 46} Despite inadequate knowledge about the causes of CVD, our respondents were much better informed about prevention measures. This finding concurs with the perception level of American women,³⁵ probably because enumerators asked prompted questions in this section of the questionnaire. In Taiwan, only 25% of respondents thought that CVD is preventable and 50% were “not sure.”^{34, 32}

Formatted: Font: Times New Roman, 12 pt

Inappropriate attitude toward cardiovascular health

Formatted: None

In contrast to their opinion that behavioral modifications can prevent CVD, most of our respondents did not consider themselves at risk and did not want to alter their lifestyle. This may be explained by the Health Belief Model^{48, 47} (i.e., a person may be knowledgeable about disease prevention but is more likely to undertake prevention when s/he perceives herself/himself to be at risk of developing the disease). Others have described a similarly optimistic bias.^{49, 48} Risk indicators (e.g., number of cigarettes smoked; death of a parent due to CVD; and self-assessment of blood pressure, weight, and cholesterol levels) account for a 24% variance in perceived risk.^{49, 48} The likelihood of perceiving oneself at higher risk increases when the presence of a risk factor is known.^{34, 32} However, respondents often underestimate their cardiovascular risk.^{49, 48}

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Attempts ~~fortoward~~ better cardiovascular health practice/behavior

Compared to Nepal's national average (69%), only half of our respondents underwent blood pressure measurement in the previous year.^{42,44} Ninety percent of the known hypertensive population had their blood pressure under control, exceeding the 6% and 9.5% control rate reported in two other suburban Kathmandu studies^{42,50,44,49} and an urban study in Eastern Nepal (58.5%).^{54,50} Half of the smokers reported here that they had tried to quit smoking. Our questionnaire did not explore whether they had actually quit and did not identify the barriers against quitting, information that might be important for public health. The theory of reasoned action^{48,47} argues that intention is a critical component of healthy behavior. Although confirming the genuineness of intention is difficult, such intention may provide an opportunity for public health intervention. Importantly, we identified our respondents' reasons for attempting positive health behavior. Many constructs of the Health Belief Model,^{48,47} such as perceived benefit ("wanted to feel better"), cues to action ("encouraged by a health professional, etc."), fear ("death of a family member, etc."), appear to affect behavioral decisions.

Influence of gender

~~Although females were unintentionally oversampled, our analysis includes a gender-wise stratification~~**Gender**

~~Seventy percent of our respondents were women. We reported earlier a male: female ratio (1.15) in the population.²¹ Enumerators used the Kish technique to select respondents at the household level, expecting a proportionate sampling.²³ However, enumerators might have interviewed more women than men because women were more likely to be available in the households when enumerators visited. Nevertheless, our analysis includes a gender-wise stratification, and we present data separately whenever we observed marked gender differences. In our study, more men than women smoked tobacco or drank alcohol. The prevalence of hypertension and self-reported diabetes was similar in both groups. On the other hand, low physical activity and obesity were higher in the women. Although men had better knowledge of heart disease, their attitude was similar to those reported by women. However, more women than men strongly agreed that their health was ultimately determined by God or a higher power. ~~Female gender is usually significantly related to good knowledge of cardiovascular disease.⁵² A Canadian study³⁶ observed a similar knowledge level among women and men.~~ One major factor for the gender~~

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None, Space After: 0 pt

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

discrepancy in our study is that more women (33.2%) than men (11.8%) lacked formal education, ~~which in itself is alarming. In fact, illiteracy is one of the main risk factors along with behavioral factors in rural/urban Asian Indian women.~~⁵³

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Effect of other socio-demographic variables on KAP

Formatted: None, Space After: 0 pt

Age-wise, KAP/behavior scores decreased around 4% from the youngest to the oldest age group.

An earlier study reported a higher level of knowledge among young respondents.³⁶ Interestingly,

Formatted: Font: Times New Roman, 12 pt

a study in adolescents reported a very poor level of cardiovascular knowledge.^{54,51} Otherwise, knowledge levels rise linearly from childhood until middle age, when the level starts to plateau.

^{55, 52} In our study, better-educated respondents ate more fruit and consumed less tobacco and

alcohol, but they also reported less physical activity and more hypertension. The highest

education category (high school or more) scored higher than the lowest category (no formal

schooling) but only by 3%–8%. Other studies observed a consistent link between better KAP

scores and higher education.^{31,36,52,56,32, 36, 53} Occupation-wise, employed respondents knew more

about biological causes (e.g., high blood pressure, diabetes, and high cholesterol), possibly due

to more education, wider exposure to information, and better access to screening and treatment.

A Canadian study reported similar findings.^{36,36}

Prompted vs. unprompted responses

Formatted: Normal, None, Tab stops: Not at 0.1" + 0.2"

Although we used unprompted responses to evaluate general knowledge, we used prompted responses to calculate scores, thus providing a denominator (“full marks”) to calculate median

score ~~%-percent~~. In other settings, KAP studies have variably used prompted^{32,39,47,57,34, 39, 46, 54} or

unprompted questions.^{46,41,18,32, 38} ~~while~~ some studies used both.^{45,35} Other studies did not

Formatted: Not Superscript/ Subscript

identify their chosen method.^{32,33} Although the accuracy of prompted vs. unprompted questions in

knowledge assessment has been a topic of debate, prompted questions confer a higher level of

knowledge.^{58,55} When we analyzed knowledge responses based on prompted responses, the

percent of correct answers increased drastically. The psychological explanation is that

respondents must simply “recognize” answers to prompted questions, but they must engage in

the more difficult mental process of “recalling” answers to unprompted questions.^{59,56} We

believe that unprompted responses more accurately reflect knowledge level because they emerge

from a thought process that ~~involves respondents’ knowledge, experience, and perception~~

~~regarding a specific issue.~~

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Comparison between levels of knowledge, attitude, and practice towards cardiovascular disease

The findings of the study indicate that there is gap in knowledge attitude and practices. Poor relation **Poor correlation between KAP scores**

Poor correlation between KAP scores was unsurprising because a linear relationship between variables may not exist in cardiovascular health (e.g. highly satisfactory knowledge=high satisfactory attitude and behavior).² Valente argued that each of six possible behavior change models contains a varying order of KAP/**behaviourpractice** and that a population can be described in terms of the percentage that fits into each model.^{60,57} This implies that no universal intervention can improve the cardiovascular health KAP of every individual in a community. The challenge lies in identifying individuals who fit into different linear models. Furthermore, health behavior theories (e.g., social cognitive theory and the theory of reasoned action)^{48,47} suggest that knowledge and attitude are only two components of the pathways that lead to behavior. Constructs (e.g., self-efficacy) and other personal and environmental factors (in social cognitive theory) and perceived behavioral control, subjective norms, and ultimately intention (in the theory of reasoned action) play crucial roles in shaping one's health behavior.^{48,47}

Study limitations

Females were unintentionally oversampled (72.7%) although the male:female ratio of the study population is 1:1.15.²⁰ Women were more likely to be available in the households when enumerators visited as most of the females in our study were housewives. The age range is limited to 25-59 years as the age groups 60-64 and 70-74 years only constitute 1-2% of the total JD-HDSS population. Most of the respondents belonged to the caste/ethnic groups Brahmin, Chhetri, and Newar which are not representative of all caste/ethnic groups in Nepal. Further, respondents' and interviewers' mood as well as place and time of interview may influence the study and a recall bias may occur when questions are asked about history of CVD risk factors.

CONCLUSION

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

Our community-based cross-sectional HARDIC study demonstrated poor knowledge, inappropriate attitude, and highly adverse behavior regarding cardiovascular health in a semi-urban community of Nepal ~~even among those already affected by CVD.~~ We aimed to determine a baseline KAP in the community to aid the planning of health promotion activities. Our results show the potential and possibility that community health promotion can improve cardiovascular health literacy.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Space After: 0 pt

ACKNOWLEDGMENTS

Acknowledgments

The authors are grateful to the study participants. We acknowledge all enumerators; field supervisors (Rachana Shrestha, Vishal Bhandari, Chandra Shova Khaitu, and Shova Poudel); and our field coordinator, Prof. Dr. Muni Raj Chhetri. We thank Umesh Raj Aryal for field supervision and advice regarding statistical analysis. We thank scientific editor Karen Williams (Kwills Editing Services, Weymouth, MA, USA) for providing professional English-language editing of this article.

CONTRIBUTORS

Contributors

AV designed the study, performed statistical analysis, and drafted the manuscript. ~~URA performed statistical analysis and revised the manuscript.~~ AK helped design the study and provided critical revision of the manuscript. ~~AH~~Both authors have read and approved the final manuscript.

FUNDING STATEMENT

Funding Statement

This study was supported by grants from the Wilhelm & Martina Lundgren's Foundation (vet1-367/2011 and vet1-379/2012) and the University of Gothenburg, Sweden through a "Global University" grant (A11 0524/09).

Formatted: Font color: Accent 6

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

COMPETING INTERESTS

Competing interests

None declared.

Data sharing

No additional data available.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: None, Space After: 0 pt

Formatted: Font color: Accent 6

Formatted: Line spacing: Multiple 1.15 li

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

References

REFERENCES

1. Alwan A. *Global status report on noncommunicable diseases 2010*. Geneva: World Health Organization; 2011.
2. World Health Organization. *Noncommunicable diseases in the South-East Asia Region: Situation and response 2011*. New Delhi: World Health Organization, Regional Office for South-East Asia; 2011.
3. Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128.
3. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998;97:596-601.
4. Reddy KS. Cardiovascular diseases in the developing countries: dimensions, determinants, dynamics and directions for public health action. *Public Health Nutr* 2002;5:231-37.
5. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation* 1998;97:596-601.
6. Engलगau MM, El-Saharty S, Kudesia P, et al: Rajan V, Rosenhouse S, Okamoto K. *Capitalizing on the demographic transition: tackling noncommunicable diseases in South Asia*. Washington, DC: World Bank Publications; 2011.
6. Mathers C, Fat DM, Boerma J. *The global burden of disease: 2004 update*. Geneva: World Health Organization; 2008.
7. Ezzati M, Lopez AD, Rodgers A, et al. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*. Geneva: World Health Organization; 2004.
8. Khaw KT, Wareham N, Bingham S, et al. Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. *PLoS medicine* 2008;5:e12.
9. Mendis S, Puska P, Norrving B. *Global atlas on cardiovascular disease prevention and control*. Geneva: World Health Organization; 2011.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Not Italic

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

10. Alwan A, Maclean DR, Riley LM, *et al*. Monitoring and surveillance of chronic non-communicable diseases: progress and capacity in high-burden countries. *Lancet* 2010;376:1861-68.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Not Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

11. Mendis S, Alwan A. *Prioritized research agenda for prevention and control of noncommunicable diseases*. Geneva: World Health Organization; 2011.

12. Puska P. From Framingham to North Karelia: from descriptive epidemiology to public health action. *Prog Cardiovasc Dis* 2010;53:15-20.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

13. Pearson TA, Bazzarre TL, Daniels SR, *et al*. American Heart Association guide for improving cardiovascular health at the community level: a statement for public health practitioners, healthcare providers, and health policy makers from the American Heart Association Expert Panel on Population and Prevention Science. *Circulation* 2003;107:645-51.

Formatted: Font: Times New Roman, 12 pt, Italic

14. Lloyd-Jones DM, Hong Y, Labarthe D, *et al*. Defining and ~~Setting National Goals~~ setting national goals for Cardiovascular Health Promotion ~~cardiovascular health promotion and Disease Reduction~~ The disease reduction: the American Heart Association's Strategic Impact Goal Through strategic impact goal through 2020 and Beyond beyond. *Circulation* 2010;121:586-613.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

15. Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health Promot Internation* 2000;15:259-67.

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt

16. Gazmararian JA, Williams MV, Peel J, *et al*. Health literacy and knowledge of chronic disease. *Patient Educ Couns* 2003;51(3):267-75.

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

17. Khan MS, Jafary FH, Faruqui AM, *et al*. High prevalence of lack of knowledge of symptoms of acute myocardial infarction in Pakistan and its contribution to delayed presentation to the hospital. *BMC Public Health* 2007;7:284.

Formatted: Font: Times New Roman, 12 pt

18. Limbu YR, Malla R, Regmi SR, *et al*. Public knowledge of heart attack in a Nepalese population survey. *Heart & Lung: The Journal of Acute and Critical Care* 2006;35:164-69.

Formatted: Font: Times New Roman, 12 pt

19. Upadhyay DK, Palaian S, Shankar PR, *et al*. Knowledge, attitude and practice about diabetes among diabetes patients in Western Nepal. *Rawal Med J* 2008;33:8-11.

Formatted: Font: Times New Roman, 12 pt

20. Vaidya A. Tackling cardiovascular health and disease in Nepal: epidemiology, strategies and implementation. *Heart Asia* 2011;3:87-91.

Formatted: Font: Times New Roman, 12 pt

19. World Health Organization. ~~Global Health Observatory Data Repository~~. <http://apps.who.int/ghodata> (accessed 28 Jun 2013).

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

2021. Aryal UR, Vaidya A, Shakya-Vaidya S, *et al*. Establishing a health demographic surveillance site in Bhaktapur district, Nepal: initial experiences and findings. *BMC Research Notes* 2012;5:489.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

2122. Vaidya A, Shakya S, Krettek A. Obesity prevalence in nepal: public health challenges in a low-income nation during an alarming worldwide trend. *International Journal of Environmental Research and Public Health* 2010;7:2726-44.

2223. Kish L. Sampling organizations and groups of unequal sizes. *Am Sociol Rev* 1965; 30:564-72.

Formatted: Font: Times New Roman, 12 pt

2324. WHO STEPS Instrument (Core and Expanded): The WHO STEPwise approach to chronic disease risk factor surveillance (STEPS). Available at: http://www.who.int/chp/steps/STEPS_Instrument_v2.1.pdf.

Formatted: Font: Times New Roman, 12 pt

2425. Lin W, Yang H, Hang C, *et al*. Nutrition knowledge, attitude, and behavior of Taiwanese elementary school children. *Asia Pac J Clin Nutr* 2007;16:534-46.

Formatted: Font: Times New Roman, 12 pt, English (U.S.)

Formatted: Font: Times New Roman, 12 pt

2526. Torabi MR, Yang J, Li J. Comparison of tobacco use knowledge, attitude and practice among college students in China and the United States. *Health Promot Internation* 2002;17:247-53.

Formatted: Font: Times New Roman, 12 pt

2627. Kosaryan M, Vahidshahi K, Siami R, *et al*. Knowledge, attitude, and practice of reproductive behavior in Iranian minor thalassemia couples. *Saudi Med J* 2009;30:835-39.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, English (U.S.)

2728. Neamatollahi H, Ebrahimi M, Talebi M, *et al*. Major differences in oral health knowledge and behavior in a group of Iranian pre-university students: a cross-sectional study. *J Oral Sci* 2011;53:177-84.

Formatted: English (U.S.)

Formatted: Font: Times New Roman, 12 pt, English (U.S.)

Formatted: Font: Times New Roman, 12 pt

2829. Sharda AJ, Shetty S. A comparative study of oral health knowledge, attitude and behaviour of first and final year dental students of Udaipur city, Rajasthan, India. *International Journal of Dental Hygiene* 2008;6:347-53.

Formatted: Font: Times New Roman, 12 pt

2930. Yap J, Lee V, Yau T, *et al*. Knowledge, attitudes and practices towards pandemic influenza among cases, close contacts, and healthcare workers in tropical Singapore: a cross-sectional survey. *BMC Public Health* 2010;10(1):442.

Formatted: Font: Times New Roman, 12 pt

3031. Maimaiti N, Shamsuddin K, Abdurahim A, *et al*. Knowledge, Attitude and Practice regarding HIV/AIDS among University students in Xinjiang. *Global Journal of Health Science* 2010;2:51.

Formatted: Font: Times New Roman, 12 pt

3132. Chen W, Yu Y, Glaser K. The Knowledge and Attitudes of Coronary Heart Disease Prevention among Middle and Older Aged People in a Community in Taipei. *Taiwan Geriatrics & Gerontology* 2009;4:251-62.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

3233. Dodani S, Mistry R, Khwaja A, *et al*. Prevalence and awareness of risk factors and behaviours of coronary heart disease in an urban population of Karachi, the largest city of Pakistan: a community survey. *Journal of Public Health* 2004;26:245-49.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Font: Times New Roman, 12 pt

3334. Saeed O, Gupta V, Dhawan N, Streja L, Shin JS, Ku M, Bhoi S, Verma S: *et al*. Knowledge of modifiable risk factors of Coronary Atherosclerotic Heart Disease (CASHD) among a sample in India. *BMC international health/International Health, and human rights/Human Rights*, 2009; 9(1):2.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt, Italic

Formatted: Font: Times New Roman, 12 pt, Italic

34. Gazmararian JA, Williams MV, Peel J, *et al*. Health literacy and knowledge of chronic disease. *Patient Educ Couns* 2003;51(3):267-75.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

35. Mosca L, Jones WK, King KB, *et al*. Awareness, perception, and knowledge of heart disease risk and prevention among women in the United States. *Arch Fam Med* 2000;9(6):506.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

36. Potvin L, Richard L, Edwards AC. Knowledge of cardiovascular disease risk factors among the Canadian population: relationships with indicators of socioeconomic status. *Can Med Assoc J* 2000;162:S5-S11.

Formatted: Font: Times New Roman, 12 pt

37. Lim K, Sumarni M, Amal N, *et al*.. Tobacco use, knowledge and attitude among Malaysians age 18 and above. *Trop Biomed* 2009;26:92-99.

Formatted: Line spacing: Multiple 1.15 li

Formatted: Font: +Body (Calibri), 11 pt

38. Wong BM, Garcia Y, Barr A, *et al*. Cardiovascular risk factor awareness in a disadvantaged inner-city population—implications for preventive strategies. *Can J Cardiol* 2008;24:677-82.

Formatted: Font: Times New Roman, 12 pt

Formatted: Normal, Space After: 0 pt, Line spacing: single

39. Schweigman K, Eichner J, Welty TK, *et al*. Cardiovascular disease risk factor awareness in American Indian communities: The Strong Heart Study. *Ethnicity and disease* 2006;16:647.

Formatted: Font: Times New Roman, 12 pt, English (U.S.)

Formatted: English (U.S.)

40. Lenfant C, Chobanian AV, Jones DW, *et al*. Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7). *Circulation* 2003;107:2993-94.

Formatted: Font: Times New Roman, 12 pt, English (U.S.)

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

41. United Nations Children Fund. *Baseline Survey of the Knowledge, Attitude and Practice (KAP) of Parents/Guardians on Early Childhood Development and Primary Education in Nepal*. Lalitpur, Nepal: UNICEF Nepal Country Office; 2009.

42-41. Central Bureau of Statistics. *National Population and Housing Census 2011*. Kathmandu, Nepal: Government of Nepal, National Planning Commission Secretariat; 2012.

42. Warning Signs of a Heart Attack. Available at: http://www.heart.org/HEARTORG/Conditions/HeartAttack/WarningSignsofaHeartAttack/Warnng-Signs-of-a-Heart-Attack_UCM_002039_Article.jsp.

Formatted: Font: +Body (Calibri)

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

~~43~~ Ministry of Health and Population, Nepal. *Nepal non-communicable diseases risk factors survey 2007*. Kathmandu, Nepal: Ministry of Health and Population, Nepal; 2008.

~~44~~ Vaidya A, Pathak RP, Pandey MR. Prevalence of hypertension in Nepalese community triples in 25 years: a repeat cross-sectional study in rural Kathmandu. *Indian Heart J* 2012;64:128-31.

~~44~~ World Health Organization. WHO Risk Factor Database. <https://apps.who.int/infobase/Indicators.aspx> (accessed 4 Aug 2013).

~~45~~ An DTM, Minh HV, Huong LT, et al. Knowledge of the health consequences of tobacco smoking: a cross-sectional survey of Vietnamese adults. *Global health action* 2013;6:18707.

~~46~~ Brook RD, Franklin B, Cascio W, et al. Air pollution and cardiovascular disease A statement for healthcare professionals from the expert panel on population and prevention science of the American Heart Association. *Circulation* 2004;109:2655-71.

~~47~~ Nguyen TT, Liao Y, Gildengorin G, et al. Cardiovascular risk factors and knowledge of symptoms among Vietnamese Americans. *Journal of General Internal Medicine* 2009;24:238-43.

~~47~~ Redding CA, Rossi S, Rossi R, et al. Health behavior models. *The International Electronic Journal of Health Education* 2000;3:180-93.

~~48~~ Avis NE, Smith KW, McKinlay JB. Accuracy of perceptions of heart attack risk: what influences perceptions and can they be changed?. *Am J Public Health* 1989; 79:1608-12.

~~49~~ Sharma D, Bkc M, Rajbhandari S, et al. Study of prevalence, awareness, and control of hypertension in a suburban area of Kathmandu, Nepal. *Indian Heart J* 2006;58:34-37.

~~50~~ Vaidya A, Pokharel P, Karki P, et al. Exploring the iceberg of hypertension: a community based study in an eastern Nepal town. *Kathmandu Univ Med J* 2007;5:349-59.

~~52~~ Attarehi M, Mohammadi S, Nojomi M, et al. Knowledge and Practice Assessment of Workers in a Pharmaceutical Company about Prevention of Coronary Artery Disease. *Acta Med Iran* 2012;50:697-703.

~~53~~ Pandey RM, Agrawal A, Misra A, et al. Population-based intervention for cardiovascular diseases related knowledge and behaviours in Asian Indian women. *Indian Heart J* 2012;65:40-7.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek – August 7, 2013

5451. Vanhecke TE, Miller WM, Franklin BA, et al. Awareness, knowledge, and perception of heart disease among adolescents. *European Journal of Cardiovascular Prevention & Rehabilitation* 2006;13:718-23.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

5552. White CW, Albanese MA. Changes in cardiovascular health knowledge occurring from childhood to adulthood. A cross-sectional study. *Circulation* 1981;63:1110-15.

Formatted: Line spacing: Multiple 1.15 li

5653. Yahya R, Muhamad R, Yusoff HM. Association between Knowledge, Attitude and Practice on Cardiovascular Disease among Women in Kelantan, Malaysia. *International Journal of Collaborative Research on Internal Medicine & Public Health* 2012;4:1507-23.

Formatted: Font: +Body (Calibri), 11 pt

5754. Muhamad R, Yahya R, Yusoff HM, et al. Knowledge, Attitude and Practice on Cardiovascular Disease among Women in North-Eastcoast Malaysia. *International Journal of Collaborative Research on Internal Medicine & Public Health* 2012;4:85-98.

Formatted: Normal, Space After: 0 pt, Line spacing: single

Formatted: Font: Times New Roman, 12 pt

5855. Waller J, McCaffery K, Wardle J. Measuring cancer knowledge: comparing prompted and unprompted recall. *Br J Psychol* 2004;95:219-34.

Formatted: Font: Times New Roman, 12 pt

5956. Glass AL, Holyoak KJ. *Cognition*. Second ed. New York: McGraw-Hill; 1986.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: Times New Roman, 12 pt

6057. Valente TW, Paredes P, Poppe PR. Matching the message to the process the relative ordering of knowledge, attitudes, and practices in behavior change research. *Human Communication Research* 1998;24:366-85.

Formatted: Font: Times New Roman, 12 pt

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REVISION bmjopen-2013-002976
Vaidya, Aryal, Krettek - August 7, 2013

Figure legends:

Figure 1: Scattergrams showing correlation between median score percents for (a) knowledge and attitude, (b) attitude and behavior, and (c) knowledge and behavior (n=777).

Scores were obtained from the prompted knowledge questions, attitudinal questions, and practice/behavior responses. Median score percents were calculated and plotted. See Supplement for greater detail about score derivation.

Formatted: Font: +Body (Calibri), 11 pt

Formatted: Left

Formatted: Normal (Web)

Formatted: Font: +Body (Calibri)

1
2
3
4
5
6
7
8
9
10
11
12 **Page 23: [1] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

13 Font: Not Bold, Italic, Font color: Text 1

14
15
16 **Page 23: [2] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

17 Font: Times New Roman, 12 pt, Italic, Font color: Text 1

18
19
20 **Page 23: [3] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

21 Font: Times New Roman, 12 pt, Italic, Font color: Text 1

22
23
24 **Page 23: [4] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

25 Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust
26 space between Asian text and numbers

27
28
29 **Page 23: [5] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

30 Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text,
31 Don't adjust space between Asian text and numbers

32
33
34 **Page 23: [6] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

35 Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text,
36 Don't adjust space between Asian text and numbers

37
38
39 **Page 23: [7] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

40 Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust
41 space between Asian text and numbers

42
43
44 **Page 23: [8] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

45 Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text,
46 Don't adjust space between Asian text and numbers

47
48
49 **Page 23: [9] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

50 Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text,
51 Don't adjust space between Asian text and numbers

52
53
54 **Page 24: [10] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

55 Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text,
56 Don't adjust space between Asian text and numbers

57
58
59 **Page 25: [11] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

60 Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust
space between Asian text and numbers

Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 25: [13] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 25: [14] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 25: [15] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 25: [16] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Centered, Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 1: [17] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font: +Body (Calibri), 11 pt

Page 1: [18] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Left

Page 1: [19] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font: +Body (Calibri)

Page 27: [20] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font: Bold, Font color: Black

Page 27: [21] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [22] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 27: [23] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Centered, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 27: [25] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [25] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [26] Inserted Cells Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Inserted Cells

Page 27: [27] Inserted Cells Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Inserted Cells

Page 27: [28] Inserted Cells Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Inserted Cells

Page 27: [29] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Page 27: [30] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [30] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [31] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [32] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [33] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Font color: Black

Page 27: [34] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

1
2
3
4
5
6
7
8
9
10
11
12 **Page 27: [35] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

13 Font color: Black

14
15
16 **Page 27: [36] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

17 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
18 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
19 numbers
20

21
22 **Page 27: [37] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

23 Font color: Black

24
25
26 **Page 27: [37] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

27 Font color: Black

28
29
30 **Page 27: [38] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

31 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
32 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
33 numbers
34

35
36
37 **Page 27: [39] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

38 Font color: Black

39
40
41 **Page 27: [39] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

42 Font color: Black

43
44
45 **Page 27: [40] Inserted Cells** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

46 Inserted Cells

47
48
49 **Page 27: [41] Inserted Cells** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

50 Inserted Cells

51
52
53 **Page 27: [42] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

54 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
55 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
56 numbers
57

58
59 **Page 27: [43] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

60 Font color: Black

For peer review only - <http://bmjopen.bmj.com/site/about/guidelines.xhtml>

Page 27: [43] Formatted **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

Font color: Black

1
2
3
4
5
6
7
8
9
10
11
12 **Page 27: [44] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

13 Font color: Black

14
15
16 **Page 27: [45] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

17 Font color: Black

18
19
20 **Page 27: [46] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

21 Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust
22 space between Asian text and numbers
23

24
25 **Page 27: [47] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

26 Font color: Black

27
28
29 **Page 27: [48] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

30 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
31 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
32 numbers
33

34
35 **Page 27: [49] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

36 Font color: Black

37
38
39 **Page 27: [50] Inserted Cells** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

40 Inserted Cells

41
42
43 **Page 27: [51] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

44 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
45 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
46 numbers
47

48
49
50 **Page 27: [52] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

51 Font color: Black

52
53
54 **Page 27: [52] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

55 Font color: Black

56
57
58 **Page 27: [53] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

59 Font color: Black

60
Page 27: [53] Formatted **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

Font color: Black

1
2
3
4
5
6
7
8
9
10
11
12 Font color: Black

13
14 **Page 27: [55] Inserted Cells** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

15
16 Inserted Cells

17
18 **Page 27: [56] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

19
20 Font color: Black

21
22 **Page 27: [57] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

23
24 Space After: 0 pt, Line spacing: 1.5 lines, Don't adjust space between Latin and Asian text, Don't adjust
25 space between Asian text and numbers

26
27 **Page 27: [58] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

28
29 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
30 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
31 numbers

32
33 **Page 27: [59] Inserted Cells** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

34
35 Inserted Cells

36
37 **Page 27: [60] Inserted Cells** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

38
39 Inserted Cells

40
41 **Page 27: [61] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

42
43 Centered, Level 9, Right: 0", Space After: 0 pt, Line spacing: 1.5 lines, Keep with next, Keep lines
44 together, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and
45 numbers

46
47 **Page 27: [62] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

48
49 Font color: Black

50
51 **Page 31: [63] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

52
53 Left, Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Tab
54 stops: 0.13", Left, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin,
55 Horizontal: 0", Wrap Around

56
57 **Page 31: [64] Formatted** Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

58
59 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
60 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around

Page 31: [65] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM

1
2
3
4
5
6
7
8
9
10
11
12 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
13 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
14
15

16 **Page 31: [66] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

17 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
18 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
19
20

21 **Page 31: [67] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

22 Left, Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Tab
23 stops: 0.13", Left, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin,
24 Horizontal: 0", Wrap Around
25
26
27

28 **Page 31: [68] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

29 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
30 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
31
32

33 **Page 31: [69] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

34 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
35 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
36
37

38 **Page 31: [70] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

39 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
40 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
41
42

43 **Page 31: [71] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

44 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
45 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
46
47
48

49 **Page 31: [72] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

50 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
51 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
52
53

54 **Page 32: [73] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

55 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
56 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
57
58

59 **Page 32: [74] Formatted Ramos, Michael Angelo D. 8/8/2013 11:51:00 AM**

60 Left, Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Tab
stops: 0.13", Left, Position: Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin,
Horizontal: 0", Wrap Around
<http://bmjopen.bmj.com/site/about/guidelines.xhtml>

1
2
3
4
5
6
7
8
9
10
11
12 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
13 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
14
15

16 **Page 32: [76] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

17 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
18 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
19
20

21 **Page 32: [77] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

22 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
23 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
24
25

26 **Page 32: [78] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

27 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
28 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
29
30

31 **Page 32: [79] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

32 Left, Level 9, Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines,
33 Keep with next, Keep lines together, Tab stops: 0.13", Left, Position: Horizontal: Left, Relative to:
34 Column, Vertical: In line, Relative to: Mar
35
36
37

38 **Page 32: [80] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

39 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
40 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
41
42

43 **Page 32: [81] Formatted** **Ramos, Michael Angelo D.** **8/8/2013 11:51:00 AM**

44 Space After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1.5 lines, Position:
45 Horizontal: Left, Relative to: Column, Vertical: In line, Relative to: Margin, Horizontal: 0", Wrap Around
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60