

SUPPLEMENTARY TABLE S1 Selected laboratory values: baseline and changes during treatment

	Patients without cirrhosis						Patients with cirrhosis	
	Placebo (n=14)	100 mg (n=9)	200 mg (n=9)	400 mg (n=9)	800 mg (n=9)	1200 mg (n=10)	400 mg (n=8) <sup>a</sup>	600 mg (n=5)
<b>ALT (U/L)</b>								
Baseline, mean±SD	79±60	81±35	68±32	99±72	101±52	83±47	147±53	110±67
Change, median (range)	12 (-6 to 50)	-18 (-93 to 7)	-11 (-69 to 4)	-43 (-67 to -8)	-45 (-119 to 48)	-37 (-121 to 11)	-62 (-108 to 7)	-33 (-80 to 8)
<b>AST (U/L)</b>								
Baseline, mean±SD	59±27	59±23	46±21	78±64	61±24	47±15	117±46	94±21
Change, median (range)	-3 (-29 to 60)	-14 (-78 to -4)	-15 (-69 to 9)	-33 (-100 to -5)	-40 (-64 to 47)	-13 (-52 to 17)	-60 (-116 to 2)	-48 (-104 to 5)
<b>White blood cells (10<sup>9</sup>/L)</b>								
Baseline, mean±SD	6.1±1.9	6.4±1.5	5.6±1.7	6.2±1.3	6.2±1.4	6.6±1.7	4.9±0.9	5.0±1.3
Change, median (range)	0.1 (-2.8 to 1.3)	-0.2 (-1.1 to 1.2)	0.4 (-1.3 to 1.9)	0.4 (0.0 to 2.3)	0.3 (-0.3 to 1.2)	0.1 (-1.6 to 1.9)	0.0 (-1.2 to 0.4)	-0.6 (-1.1 to -0.3)
<b>Platelet count (10<sup>9</sup>/L)</b>								
Baseline, mean±SD	186±25	202±55	193±35	220±70	192±30	205±43	118±19	116±23
Change, median (range)	3 (-17 to 28)	9 (-40 to 27)	8 (-10 to 27)	10 (-80 to 21)	2 (-27 to 15)	11 (-4 to 21)	4 (-9 to 11)	-12 (-16 to -8)
<b>Hemoglobin (g/dL)</b>								
Baseline, mean±SD	14.8±1.2	15.1±1.1	15.0±1.1	14.3±2.1	15.6±0.7	15.4±0.5	14.9±2.0	13.7±1.2
Change, median (range)	-0.2 (-1.7 to 1.0)	0.0 (-0.8 to 0.6)	-0.8 (-1.7 to -0.1)	-0.3 (-1.1 to 0.0)	-0.6 (-1.3 to 0.5)	-0.6 (-1.5 to 0.3)	-0.6 (-1.8 to 0.9)	-1.6 (-1.9 to -0.6)
<b>Creatinine (mg/dL)</b>								
Baseline, mean±SD	0.8±0.1	0.7±0.1	0.8±0.2	0.9±0.2	0.8±0.1	0.8±0.1	0.7±0.3	0.9±0.2
Change, median (range)	-0.1 (-0.3 to 0.1)	0.1 (-0.2 to 0.4)	0.0 (-0.1 to 0.2)	-0.1 (-0.1 to 0.1)	0.0 (-0.2 to 0.1)	0.0 (-0.2 to 0.1)	-0.1 (-0.2 to 0.1)	-0.1 (-0.2 to 0.0)

<sup>a</sup> Includes one treatment-naïve patient.