

Table S1. List of designed proteins from previous experiments.

Sequence ID	PDB ID	Length	Failed step ^a	References
1lq7A	1lq7A	67	Success	Dai et al [1]
1mj0A	1mj0A	156	Success	Kohl et al [2]
1p68A	1p68A	102	Success	Wei et al [3]
1qysA	1qysA	91	Success	Kuhlman et al [4]
1vjqA	1vjqA	71	Success	Dantas et al [5]
2a3dA	2a3dA	73	Success	Walsh et al [6]
2cw1A	2cw1A	65	Success	Walsh et al [6]
2juaA	2juaA	102	Success	Go et al [7]
2jvfA	2jvfA	94	Success	Stordeur et al [8]
2kl8A	2kl8A	85	Success	Koga et al [9]
2ln3A	2ln3A	83	Success	Koga et al [9]
2ltaA	2ltaA	110	Success	Koga et al [9]
2lv8A	2lv8A	110	Success	Koga et al [9]
2lvbA	2lvbA	112	Success	Koga et al [9]
2qyjA	2qyjA	154	Success	Merz et al [10]
3b83A	3b83A	95	Success	Hu et al [11]
MSP2-FC27	/	237	Natively unfolded	Zhang et al [12]
alpha-synuclein	/	140	Natively unfolded	Eliezer et al [13]
4EbindingproteinI	/	118	Natively unfolded	Fletcher et al [14]
4EbindingproteinII	/	120	Natively unfolded	Fletcher et al [14]
Fold-I-11	/	76	Not soluble	Koga et al [9]
Fold-I-2	/	76	Not folded-CD	Koga et al [9]
Fold-I-4	/	76	Not folded-CD	Koga et al [9]
Fold-I-8	/	76	Not folded NMR	Koga et al [9]
Fold-II-8	/	99	Not folded-CD	Koga et al [9]
Fold-II-9	/	99	Not folded-CD	Koga et al [9]
Fold-II-1	/	99	Not folded NMR	Koga et al [9]
Fold-II-3	/	99	Not folded NMR	Koga et al [9]
Fold-II-11	/	99	Not folded NMR	Koga et al [9]
Fold-III-6	/	72	Not soluble	Koga et al [9]
Fold-III-7	/	72	Not soluble	Koga et al [9]
Fold-III-1	/	72	Incorrect CD	Koga et al [9]
Fold-III-13	/	72	Not folded-CD	Koga et al [9]
Fold-III-4	/	72	Not folded NMR	Koga et al [9]
Fold-IV-2	/	101	Not folded NMR	Koga et al [9]
Fold-V-8	/	99	Not soluble	Koga et al [9]
Fold-V-1	/	99	Not folded-CD	Koga et al [9]
Fold-V-2	/	99	Not folded-CD	Koga et al [9]
Fold-V-4	/	99	Not folded-CD	Koga et al [9]
Fold-V-9	/	99	Not folded-CD	Koga et al [9]
Fold-V-10	/	99	Not folded-CD	Koga et al [9]
Fold-V-11	/	99	Not folded-CD	Koga et al [9]

Fold-V-12	/	99	Not folded-CD	Koga et al [9]
MDM2a	/	85	Not folded-CD	Shultzis et al [15]
MDM2b	/	85	Not folded-CD	Shultzis et al [15]

^aSuccess: protein structure was solvable by NMR or X-ray crystallography; **Natively unfolded:** biologically expressed (i.e. not designed) protein experimentally known to be unfolded in the native state; **Not soluble:** protein was not soluble after expression (usually indicative of improperly exposed hydrophobic surfaces); **Not folded-CD:** largely random coil secondary structure by CD; **Incorrect CD:** secondary structure by CD not consistent with predicted structure; **Not folded NMR:** poorly resolved NMR spectra reflective of a dynamic (unstable) protein core.

References

1. Dai QH, Tommos C, Fuentes EJ, Blomberg MR, Dutton PL, et al. (2002) Structure of a de novo designed protein model of radical enzymes. *J Am Chem Soc* 124: 10952-10953.
2. Kohl A, Binz HK, Forrer P, Stumpp MT, Pluckthun A, et al. (2003) Designed to be stable: crystal structure of a consensus ankyrin repeat protein. *Proc Natl Acad Sci U S A* 100: 1700-1705.
3. Wei Y, Kim S, Fela D, Baum J, Hecht MH (2003) Solution structure of a de novo protein from a designed combinatorial library. *Proc Natl Acad Sci U S A* 100: 13270-13273.
4. Kuhlman B, Dantas G, Ireton GC, Varani G, Stoddard BL, et al. (2003) Design of a novel globular protein fold with atomic-level accuracy. *Science* 302: 1364-1368.
5. Dantas G, Corrent C, Reichow SL, Havranek JJ, Eletr ZM, et al. (2007) High-resolution structural and thermodynamic analysis of extreme stabilization of human procarboxypeptidase by computational protein design. *Journal of Molecular Biology* 366: 1209-1221.
6. Walsh ST, Cheng H, Bryson JW, Roder H, DeGrado WF (1999) Solution structure and dynamics of a de novo designed three-helix bundle protein. *Proc Natl Acad Sci U S A* 96: 5486-5491.
7. Go A, Kim S, Baum J, Hecht MH (2008) Structure and dynamics of de novo proteins from a designed superfamily of 4-helix bundles. *Protein Sci* 17: 821-832.
8. Stordeur C, Dalluge R, Birkenmeier O, Wienk H, Rudolph R, et al. (2008) The NMR solution structure of the artificial protein M7 matches the computationally designed model. *Proteins* 72: 1104-1107.
9. Koga N, Tatsumi-Koga R, Liu G, Xiao R, Acton TB, et al. (2012) Principles for designing ideal protein structures. *Nature* 491: 222-227.
10. Merz T, Wetzel SK, Firbank S, Pluckthun A, Grutter MG, et al. (2008) Stabilizing ionic interactions in a full-consensus ankyrin repeat protein. *Journal of Molecular Biology* 376: 232-240.
11. Hu X, Wang H, Ke H, Kuhlman B (2008) Computer-based redesign of a beta sandwich protein suggests that extensive negative design is not required for de novo beta sheet design. *Structure* 16: 1799-1805.

12. Zhang X, Perugini MA, Yao S, Adda CG, Murphy VJ, et al. (2008) Solution conformation, backbone dynamics and lipid interactions of the intrinsically unstructured malaria surface protein MSP2. *Journal of Molecular Biology* 379: 105-121.
13. Eliezer D, Kutluay E, Bussell R, Jr., Browne G (2001) Conformational properties of alpha-synuclein in its free and lipid-associated states. *Journal of Molecular Biology* 307: 1061-1073.
14. Fletcher CM, McGuire AM, Gingras AC, Li H, Matsuo H, et al. (1998) 4E binding proteins inhibit the translation factor eIF4E without folded structure. *Biochemistry* 37: 9-15.
15. Shultis D, Mitra P, Zhang Y (2013) Unpublished results.