

cgcaaaaacgttgactttatctcggctttaaaggcaaaaactaaagaaaaataatgcattgagttagtaaaaaaattaattgtttcaaa
 ttatagaaattattaaacgagcaagtcaggacgaaaftagaatcttggacctccaagatgcgcaaaaattatcttcatcttcttci
 caattcccgttcactcgcctcgcactttatcaac(agatct)gtcagcgacatacgaatgaatcgattatcttagaggagaggcagat
 gccaaaaaattacagtcaaaaagaagtattcgtgaaagagtgcgcaaatcgagagacgcagacagaatgcaaacgcgctccgct
 gaactgtggccgccaagcgtttcgtttaaaggagcggtgtctatcttctatcttctgcccagttttatcttgcgcaactatcttttacta
 cgatctcgaaaaatatttcaaagcaagaaagtaattatcttgcgttcagaaaccgagctcatgagtaaaaggagaagaactttcaactg
 gagtt

■ Nucleotides 1 to 228 upstream of the *lab-1* ATG

■ *baf-1* promoter

■ Plasmid backbone

■ GFP

() BglII site used to clone *lab-1* upstream sequence

Figure S3 Confirmation of seamless recombination at the *lab-1* locus. Sequence resulting from the PCR product generated by using primers 1100 base pairs upstream and downstream of *lab-1*. Sequences from the *lab-1* upstream region (violet), *baf-1* promoter (yellow), plasmid backbone (light blue), and *gfp* (green) are marked. The site by which the upstream fragment was cloned is marked with brackets.