Supplemental Table 1: Rep	ported Viru	s-ESCRT Interaction	ons	
Virus	Viral Protein	Late Assembly Domain	Early-Acting ESCRT Factor	References
		Enveloped Viruse	<u>es</u>	
RNA Viruses				
Retroviruses				
Human Immunodeficiency virus Type 1 (HIV-1)	Gag	P(T/S)AP YPXL unknown	TSG101 ALIX NEDD4L	[15,19,25,28,30,39,59, 82,92,93,95]
Simian Immunodeficiency Virus (and HIV-2)	Gag	P(T/S)AP YPXL	TSG101 ALIX	[5,6,61,82,110]
Feline Immunodeficiency Virus	Gag	P(T/S)AP unknown	TSG101 NEDD4L	[9,57]
Equine Infectious Anemia Virus (EIAV)	Gag	YPXL	ALIX	[13,53,67,68,75,82,10 9]
Human T Cell Leukemia Virus	Gag	P(T/S)AP PPXY unknown	TSG101 NEDD4 ALIX?	[8,22,35,36,51,91,96]
Prototypic Foamy Virus	Gag	P(T/S)AP PPXY?	TSG101 NEDD4 family member?	[63,81,107,108]
Rous Sarcoma Virus	Gag	PPXY YPXL	NEDD4 family ALIX	[20,27,47,60,80,94,99, 102]
Mason-Pfizer Monkey Virus	Gag	P(T/S)AP PPXY	TSG101 NEDD4 family	[31,103]
Murine Leukemia Virus (MLV)	Gag	P(T/S)AP YPXL PPXY unknown	TSG101 ALIX NEDD4 family NEDD4 family	[44,74,105,106]
Bovine Leukemia Virus	Gag	РРХҮ	NEDD4 family member?	[97]
Koala Retrovirus	Gag	PPXY P(T/S)AP?	NEDD4 family TSG101?	[76]

Filoviruses				
Ebola Virus	VP40	P(T/S)AP PPXY	TSG101 NEDD4 family	[33,41,56,59,86,87,10 4]
Marburg Virus	NP VP40	P(T/S)AP PPXY	TSG101 NEDD4 family member?	[21,48,88]
Arenaviruses				
Lymphocytic Choriomeningitis Virus	Z	P(T/S)AP PPXY	TSG101 NEDD4 family member?	[66]
Lassa Fever Virus	Z	P(T/S)AP PPXY	TSG101 NEDD4 family member?	[66,89]
Mopeia Virus	Z	unknown	ALIX	[77]
Tacaribe Virus	Z	unknown	unknown	[90]
Paramyxoviruses				
Human Parainfluenzavirus Type 5 (hPIV-5)	Μ	FPIV	unknown AMOT-L1	[73] [65]
Human Parainfluenzavirus Type 1	С	unknown	ALIX	[7]
Sendai Virus ¹	С	unknown	ALIX	[29,42,43,72]
	М	YPXL	ALIX	
Mumps Virus	unknown	FPIV	unknown	[54]
Rhabdoviruses				
Vesicular Stomatitis Virus ²	Μ	PPXY P(T/S)AP?	NEDD4 family TSG101?	[17,32,34,40,45,50,85]
Rabies Virus	М	РРХҮ	NEDD4 family	[34,101]
Flaviviruses				
Yellow Fever Virus	NS3	YPXL	ALIX	[11]
Dengue Fever Virus ²	unknown	unknown	ALIX	[62]
Hepatitis C Virus ⁴	unknown	unknown	unknown	[1,16,23,69,83]
Japanese Encephalitis Virus	NS3	unknown	TSG101	[14]

DNA Viruses

Her	npsviriisps
IICI.	9C3VII 43C3

Herpes Simples Virus	unknown	unknown	unknown	[10,18,64]
Epstein-Barr Virus	BFRF	unknown	ALIX	[52,71]
Human Cytomegalovirus ¹	unknown	unknown	unknown	[26,84]
Vaccinia virus	F13L	YPXL unknown	ALIX TSG101	[38]

Hepadnaviruses

Henatitis B Virus ¹	Core	ΡΡΧΥ	NEDD4 family	[1 37 16 19 70 98]
	unknown	unknown	TSG101	[1,57,70,75,70,96]
		unknown	ALIX	
			lpha-Taxilin	

Non-enveloped Viruses

RNA Viruses

Reoviruses

Bluetongue Virus ³	NS3/NS3A	P(T/S)AP PPXY	TSG101 NEDD4 family member?	[12,100]
Rotavirus ²	unknown	unknown	unknown	[78]
Picornaviruses				
Hepatitis A Virus ³	unknown	unknown	ALIX	[24]

Insect Viruses

Baculoviruses

Autographa californica Multiple Nucleopolyhedrovirus ²	unknown	unknown	unknown	[55]	
		Archaeal Viru	ises		
Sulfolobus Turreted Icosahedral Virus	unknown	unknown	unknown	[58,79]	
		Plant Viruse	<u>25</u>		
Tombusviruses					
Tomato Bushy Stunt Virus	p33	unknown	TSG101 ALIX	[2,3]	

¹Cases in which other studies have failed to confirm ESCRT involvement

²Cases in which ESCRT factors are (also) implicated in virus entry

³Non-enveloped viruses that can also be released as enveloped particles via ESCRT-dependent budding ⁴Case in which ESCRT pathway is also involved in RNA trafficking

Table References

[1] Ariumi, Y., Kuroki, M., Maki, M., Ikeda, M., Dansako, H., Wakita, T., and Kato, N. (2011). The ESCRT system is required for hepatitis C virus production. PloS one *6*, e14517.

[2] Barajas, D., Jiang, Y., and Nagy, P.D. (2009). A unique role for the host ESCRT proteins in replication of Tomato bushy stunt virus. PLoS pathogens *5*, e1000705.

[3] Barajas, D., and Nagy, P.D. (2010). Ubiquitination of tombusvirus p33 replication protein plays a role in virus replication and binding to the host Vps23p ESCRT protein. Virology *397*, 358-368.

[4] Bardens, A., Doring, T., Stieler, J., and Prange, R. (2011). Alix regulates egress of hepatitis B virus naked capsid particles in an ESCRT-independent manner. Cellular microbiology *13*, 602-619.

[5] Bello, N.F., Wu, F., Sette, P., Dussupt, V., Hirsch, V.M., and Bouamr, F. (2011). Distal leucines are key functional determinants of Alix-binding simian immunodeficiency virus SIV(smE543) and SIV(mac239) type 3 L domains. J Virol *85*, 11532-11537.

[6] Bibollet-Ruche, F., Bailes, E., Gao, F., Pourrut, X., Barlow, K.L., Clewley, J.P., Mwenda, J.M., Langat, D.K., Chege, G.K., McClure, H.M., *et al.* (2004). New simian immunodeficiency virus infecting De Brazza's monkeys (Cercopithecus neglectus): evidence for a cercopithecus monkey virus clade. J Virol *78*, 7748-7762.

[7] Boonyaratanakornkit, J., Schomacker, H., Collins, P., and Schmidt, A. (2013). Alix Serves as an Adaptor That Allows Human Parainfluenza Virus Type 1 to Interact with the Host Cell ESCRT System. PLoS One *8*, e59462.

[8] Bouamr, F., Melillo, J.A., Wang, M.Q., Nagashima, K., de Los Santos, M., Rein, A., and Goff, S.P. (2003). PPPYVEPTAP motif is the late domain of human T-cell leukemia virus type 1 Gag and mediates its functional interaction with cellular proteins Nedd4 and Tsg101 [corrected]. J Virol *77*, 11882-11895.

[9] Calistri, A., Del Vecchio, C., Salata, C., Celestino, M., Celegato, M., Gottlinger, H., Palu, G., and Parolin, C. (2009). Role of the feline immunodeficiency virus L-domain in the presence or absence of Gag processing: involvement of ubiquitin and Nedd4-2s ligase in viral egress. J Cell Physiol *218*, 175-182.

[10] Calistri, A., Sette, P., Salata, C., Cancellotti, E., Forghieri, C., Comin, A., Gottlinger, H., Campadelli-Fiume, G., Palu, G., and Parolin, C. (2007). Intracellular trafficking and maturation of herpes simplex virus type 1 gB and virus egress require functional biogenesis of multivesicular bodies. J Virol *81*, 11468-11478.

[11] Carpp, L.N., Galler, R., and Bonaldo, M.C. (2011). Interaction between the yellow fever virus nonstructural protein NS3 and the host protein Alix contributes to the release of infectious particles. Microbes and infection / Institut Pasteur *13*, 85-95.

[12] Celma, C.C., and Roy, P. (2009). A viral nonstructural protein regulates bluetongue virus trafficking and release. Journal of virology *83*, 6806-6816.

[13] Chen, C., Vincent, O., Jin, J., Weisz, O.A., and Montelaro, R.C. (2005). Functions of early (AP-2) and late (AIP1/ALIX) endocytic proteins in equine infectious anemia virus budding. J Biol Chem *280*, 40474-40480.

[14] Chiou, C.T., Hu, C.C., Chen, P.H., Liao, C.L., Lin, Y.L., and Wang, J.J. (2003). Association of Japanese encephalitis virus NS3 protein with microtubules and tumour susceptibility gene 101 (TSG101) protein. J Gen Virol *84*, 2795-2805.

[15] Chung, H.Y., Morita, E., von Schwedler, U., Muller, B., Krausslich, H.G., and Sundquist, W.I. (2008). NEDD4L overexpression rescues the release and infectivity of human immunodeficiency virus type 1 constructs lacking PTAP and YPXL late domains. J Virol *82*, 4884-4897.

[16] Corless, L., Crump, C.M., Griffin, S.D., and Harris, M. (2010). Vps4 and the ESCRT-III complex are required for the release of infectious hepatitis C virus particles. The Journal of general virology *91*, 362-372.

[17] Craven, R.C., Harty, R.N., Paragas, J., Palese, P., and Wills, J.W. (1999). Late domain function identified in the vesicular stomatitis virus M protein by use of rhabdovirus-retrovirus chimeras. J Virol *73*, 3359-3365.

[18] Crump, C.M., Yates, C., and Minson, T. (2007). Herpes simplex virus type 1 cytoplasmic envelopment requires functional Vps4. J Virol *81*, 7380-7387.

[19] Demirov, D.G., Ono, A., Orenstein, J.M., and Freed, E.O. (2002). Overexpression of the N-terminal domain of TSG101 inhibits HIV-1 budding by blocking late domain function. Proc Natl Acad Sci USA *99*, 955-960.

[20] Dilley, K.A., Gregory, D., Johnson, M.C., and Vogt, V.M. (2010). An LYPSL late domain in the gag protein contributes to the efficient release and replication of Rous sarcoma virus. J Virol *84*, 6276-6287.

[21] Dolnik, O., Kolesnikova, L., Stevermann, L., and Becker, S. (2010). Tsg101 is recruited by a late domain of the nucleocapsid protein to support budding of Marburg virus-like particles. J Virol *84*, 7847-7856.

[22] Dorweiler, I.J., Ruone, S.J., Wang, H., Burry, R.W., and Mansky, L.M. (2006). Role of the human T-cell leukemia virus type 1 PTAP motif in Gag targeting and particle release. J Virol *80*, 3634-3643.

[23] Dreux, M., Garaigorta, U., Boyd, B., Decembre, E., Chung, J., Whitten-Bauer, C., Wieland, S., and Chisari, F.V. (2012). Short-range exosomal transfer of viral RNA from infected cells to plasmacytoid dendritic cells triggers innate immunity. Cell Host Microbe *12*, 558-570.

[24] Feng, Z., Hensley, L., McKnight, K.L., Hu, F., Madden, V., Ping, L., Jeong, S.H., Walker, C., Lanford, R.E., and Lemon, S.M. (2013). A pathogenic picornavirus acquires an envelope by hijacking cellular membranes. Nature *496*, 367-371.

[25] Fisher, R.D., Chung, H.Y., Zhai, Q., Robinson, H., Sundquist, W.I., and Hill, C.P. (2007). Structural and biochemical studies of ALIX/AIP1 and its role in retrovirus budding. Cell *128*, 841-852.

[26] Fraile-Ramos, A., Pelchen-Matthews, A., Risco, C., Rejas, M.T., Emery, V.C., Hassan-Walker, A.F., Esteban, M., and Marsh, M. (2007). The ESCRT machinery is not required for human cytomegalovirus envelopment. Cellular microbiology *9*, 2955-2967.

[27] Garnier, L., Wills, J.W., Verderame, M.F., and Sudol, M. (1996). WW domains and retrovirus budding. Nature *381*, 744-745.

[28] Garrus, J.E., von Schwedler, U.K., Pornillos, O.W., Morham, S.G., Zavitz, K.H., Wang, H.E., Wettstein, D.A., Stray, K.M., Cote, M., Rich, R.L., *et al.* (2001). Tsg101 and the vacuolar protein sorting pathway are essential for HIV-1 budding. Cell *107*, 55-65.

[29] Gosselin-Grenet, A.S., Marq, J.B., Abrami, L., Garcin, D., and Roux, L. (2007). Sendai virus budding in the course of an infection does not require Alix and VPS4A host factors. Virology *365*, 101-112.

[30] Gottlinger, H.G., Dorfman, T., Sodroski, J.G., and Haseltine, W.A. (1991). Effect of mutations affecting the p6 gag protein on human immunodeficiency virus particle release. Proc Natl Acad Sci USA *88*, 3195-3199.

[31] Gottwein, E., Bodem, J., Muller, B., Schmechel, A., Zentgraf, H., and Krausslich, H.G. (2003). The Mason-Pfizer monkey virus PPPY and PSAP motifs both contribute to virus release. J Virol 77, 9474-9485.

[32] Harty, R.N., Brown, M.E., McGettigan, J.P., Wang, G., Jayakar, H.R., Huibregtse, J.M., Whitt, M.A., and Schnell, M.J. (2001). Rhabdoviruses and the cellular ubiquitin-proteasome system: a budding interaction. J Virol *75*, 10623-10629.

[33] Harty, R.N., Brown, M.E., Wang, G., Huibregtse, J., and Hayes, F.P. (2000). A PPxY motif within the VP40 protein of ebola virus interacts physically and functionally with a ubiquitin ligase: implications for filovirus budding. Proc Natl Acad Sci USA *97*, 13871-13876.

[34] Harty, R.N., Paragas, J., Sudol, M., and Palese, P. (1999). A proline-rich motif within the matrix protein of vesicular stomatitis virus and rabies virus interacts with WW domains of cellular proteins: implications for viral budding. J Virol *73*, 2921-2929.

[35] Heidecker, G., Lloyd, P.A., Fox, K., Nagashima, K., and Derse, D. (2004). Late assembly motifs of human T-cell leukemia virus type 1 and their relative roles in particle release. J Virol *78*, 6636-6648.

[36] Heidecker, G., Lloyd, P.A., Soheilian, F., Nagashima, K., and Derse, D. (2007). The role of WWP1-Gag interaction and Gag ubiquitination in assembly and release of human T-cell leukemia virus type 1. J Virol *81*, 9769-9777.

[37] Hoffmann, J., Boehm, C., Himmelsbach, K., Donnerhak, C., Roettger, H., T, S.W., Ploen, D., and Hildt, E. (2013). Identification of alpha-taxilin as an essential factor for the life cycle of Hepatitis B virus. Journal of hepatology, Epublication ahead of print.

[38] Honeychurch, K.M., Yang, G., Jordan, R., and Hruby, D.E. (2007). The vaccinia virus F13L YPPL motif is required for efficient release of extracellular enveloped virus. J Virol *81*, 7310-7315.

[39] Huang, M., Orenstein, J.M., Martin, M.A., and Freed, E.O. (1995). p6Gag is required for particle production from full-length human immunodeficiency virus type 1 molecular clones expressing protease. J Virol *69*, 6810-6818.

[40] Irie, T., Carnero, E., Okumura, A., Garcia-Sastre, A., and Harty, R.N. (2007). Modifications of the PSAP region of the matrix protein lead to attenuation of vesicular stomatitis virus in vitro and in vivo. J Gen Virol *88*, 2559-2567.

[41] Irie, T., Licata, J.M., and Harty, R.N. (2005). Functional characterization of Ebola virus L-domains using VSV recombinants. Virology *336*, 291-298.

[42] Irie, T., Nagata, N., Yoshida, T., and Sakaguchi, T. (2008). Recruitment of Alix/AIP1 to the plasma membrane by Sendai virus C protein facilitates budding of virus-like particles. Virology *371*, 108-120.

[43] Irie, T., Shimazu, Y., Yoshida, T., and Sakaguchi, T. (2007). The YLDL sequence within Sendai virus M protein is critical for budding of virus-like particles and interacts with Alix/AIP1 independently of C protein. J Virol *81*, 2263-2273.

[44] Jadwin, J.A., Rudd, V., Sette, P., Challa, S., and Bouamr, F. (2010). Late domain-independent rescue of a release-deficient Moloney murine leukemia virus by the ubiquitin ligase itch. J Virol *84*, 704-715.

[45] Jayakar, H.R., Murti, K.G., and Whitt, M.A. (2000). Mutations in the PPPY motif of vesicular stomatitis virus matrix protein reduce virus budding by inhibiting a late step in virion release. J Virol 74, 9818-9827.

[46] Kian Chua, P., Lin, M.H., and Shih, C. (2006). Potent inhibition of human Hepatitis B virus replication by a host factor Vps4. Virology *354*, 1-6.

[47] Kikonyogo, A., Bouamr, F., Vana, M.L., Xiang, Y., Aiyar, A., Carter, C., and Leis, J. (2001). Proteins related to the Nedd4 family of ubiquitin protein ligases interact with the L domain of Rous sarcoma virus and are required for gag budding from cells. Proc Natl Acad Sci USA *98*, 11199-11204.

[48] Kolesnikova, L., Strecker, T., Morita, E., Zielecki, F., Mittler, E., Crump, C., and Becker, S. (2009). Vacuolar protein sorting pathway contributes to the release of Marburg virus. J Virol *83*, 2327-2337.

[49] Lambert, C., Doring, T., and Prange, R. (2007). Hepatitis B virus maturation is sensitive to functional inhibition of ESCRT-III, Vps4, and gamma 2-adaptin. J Virol *81*, 9050-9060.

[50] Le Blanc, I., Luyet, P.P., Pons, V., Ferguson, C., Emans, N., Petiot, A., Mayran, N., Demaurex, N., Faure, J., Sadoul, R., *et al.* (2005). Endosome-to-cytosol transport of viral nucleocapsids. Nat Cell Biol *7*, 653-664.

[51] Le Blanc, I., Prevost, M.C., Dokhelar, M.C., and Rosenberg, A.R. (2002). The PPPY motif of human T-cell leukemia virus type 1 Gag protein is required early in the budding process. J Virol *76*, 10024-10029.

[52] Lee, C.P., Liu, P.T., Kung, H.N., Su, M.T., Chua, H.H., Chang, Y.H., Chang, C.W., Tsai, C.H., Liu, F.T., and Chen, M.R. (2012). The ESCRT machinery is recruited by the viral BFRF1 protein to the nucleus-associated membrane for the maturation of Epstein-Barr Virus. PLoS pathogens *8*, e1002904.

[53] Li, F., Chen, C., Puffer, B.A., and Montelaro, R.C. (2002). Functional replacement and positional dependence of homologous and heterologous L domains in equine infectious anemia virus replication. J Virol *76*, 1569-1577.

[54] Li, M., Schmitt, P.T., Li, Z., McCrory, T.S., He, B., and Schmitt, A.P. (2009). Mumps virus matrix, fusion, and nucleocapsid proteins cooperate for efficient production of virus-like particles. J Virol *83*, 7261-7272.

[55] Li, Z., and Blissard, G.W. (2012). Cellular VPS4 is required for efficient entry and egress of budded virions of Autographa californica multiple nucleopolyhedrovirus. Journal of virology *86*, 459-472.

[56] Licata, J.M., Simpson-Holley, M., Wright, N.T., Han, Z., Paragas, J., and Harty, R.N. (2003). Overlapping motifs (PTAP and PPEY) within the Ebola virus VP40 protein function independently as late budding domains: involvement of host proteins TSG101 and VPS-4. J Virol *77*, 1812-1819.

[57] Luttge, B.G., Shehu-Xhilaga, M., Demirov, D.G., Adamson, C.S., Soheilian, F., Nagashima, K., Stephen, A.G., Fisher, R.J., and Freed, E.O. (2008). Molecular characterization of feline immunodeficiency virus budding. J Virol *82*, 2106-2119.

[58] Maaty, W.S., Ortmann, A.C., Dlakic, M., Schulstad, K., Hilmer, J.K., Liepold, L., Weidenheft, B., Khayat, R., Douglas, T., Young, M.J., *et al.* (2006). Characterization of the archaeal thermophile Sulfolobus turreted icosahedral virus validates an evolutionary link among double-stranded DNA viruses from all domains of life. J Virol *80*, 7625-7635.

[59] Martin-Serrano, J., Zang, T., and Bieniasz, P.D. (2001). HIV-1 and Ebola virus encode small peptide motifs that recruit Tsg101 to sites of particle assembly to facilitate egress. Nat Med *7*, 1313-1319.

[60] Medina, G., Zhang, Y., Tang, Y., Gottwein, E., Vana, M.L., Bouamr, F., Leis, J., and Carter, C.A. (2005). The functionally exchangeable L domains in RSV and HIV-1 Gag direct particle release through pathways linked by Tsg101. Traffic *6*, 880-894.

[61] Myers, E.L., and Allen, J.F. (2002). Tsg101, an inactive homologue of ubiquitin ligase e2, interacts specifically with human immunodeficiency virus type 2 gag polyprotein and results in increased levels of ubiquitinated gag. J Virol *76*, 11226-11235.

[62] Pattanakitsakul, S.N., Poungsawai, J., Kanlaya, R., Sinchaikul, S., Chen, S.T., and Thongboonkerd, V. (2010). Association of Alix with late endosomal lysobisphosphatidic acid is important for dengue virus infection in human endothelial cells. Journal of proteome research *9*, 4640-4648.

[63] Patton, G.S., Morris, S.A., Chung, W., Bieniasz, P.D., and McClure, M.O. (2005). Identification of domains in gag important for prototypic foamy virus egress. J Virol *79*, 6392-6399.

[64] Pawliczek, T., and Crump, C.M. (2009). Herpes simplex virus type 1 production requires a functional ESCRT-III complex but is independent of TSG101 and ALIX expression. Journal of virology *83*, 11254-11264.

[65] Pei, Z., Bai, Y., and Schmitt, A.P. (2010). PIV5 M protein interaction with host protein angiomotinlike 1. Virology *397*, 155-166. [66] Perez, M., Craven, R.C., and De La Torre, J.C. (2003). The small RING finger protein Z drives arenavirus budding: Implications for antiviral strategies. Proc Natl Acad Sci USA *100*, 12978-12983.

[67] Puffer, B.A., Parent, L.J., Wills, J.W., and Montelaro, R.C. (1997). Equine infectious anemia virus utilizes a YXXL motif within the late assembly domain of the Gag p9 protein. J Virol *71*, 6541-6546.

[68] Puffer, B.A., Watkins, S.C., and Montelaro, R.C. (1998). Equine infectious anemia virus Gag polyprotein late domain specifically recruits cellular AP-2 adapter protein complexes during virion assembly. J Virol *72*, 10218-10221.

[69] Ramakrishnaiah, V., Thumann, C., Fofana, I., Habersetzer, F., Pan, Q., de Ruiter, P.E., Willemsen, R., Demmers, J.A., Stalin Raj, V., Jenster, G., *et al.* (2013). Exosome-mediated transmission of hepatitis C virus between human hepatoma Huh7.5 cells. Proc Natl Acad Sci USA *110*, 13109-13113.

[70] Rost, M., Mann, S., Lambert, C., Doring, T., Thome, N., and Prange, R. (2006). Gamma-adaptin, a novel ubiquitin-interacting adaptor, and Nedd4 ubiquitin ligase control hepatitis B virus maturation. J Biol Chem *281*, 29297-29308.

[71] Ruiss, R., Jochum, S., Wanner, G., Reisbach, G., Hammerschmidt, W., and Zeidler, R. (2011). A viruslike particle-based Epstein-Barr virus vaccine. Journal of virology *85*, 13105-13113.

[72] Sakaguchi, T., Kato, A., Sugahara, F., Shimazu, Y., Inoue, M., Kiyotani, K., Nagai, Y., and Yoshida, T. (2005). AIP1/Alix is a binding partner of Sendai virus C protein and facilitates virus budding. J Virol *79*, 8933-8941.

[73] Schmitt, A.P., Leser, G.P., Morita, E., Sundquist, W.I., and Lamb, R.A. (2005). Evidence for a new viral late-domain core sequence, FPIV, necessary for budding of a paramyxovirus. J Virol *79*, 2988-2997.

[74] Segura-Morales, C., Pescia, C., Chatellard-Causse, C., Sadoul, R., Bertrand, E., and Basyuk, E. (2005). Tsg101 and Alix interact with murine leukemia virus Gag and cooperate with Nedd4 ubiquitin ligases during budding. J Biol Chem *280*, 27004-27012.

[75] Shehu-Xhilaga, M., Ablan, S., Demirov, D.G., Chen, C., Montelaro, R.C., and Freed, E.O. (2004). Late domain-dependent inhibition of equine infectious anemia virus budding. J Virol *78*, 724-732.

[76] Shimode, S., Nakaoka, R., Hoshino, S., Abe, M., Shogen, H., Yasuda, J., and Miyazawa, T. (2013). Identification of cellular factors required for the budding of koala retrovirus. Microbiology and immunology *57*, 543-546.

[77] Shtanko, O., Watanabe, S., Jasenosky, L.D., Watanabe, T., and Kawaoka, Y. (2011). ALIX/AIP1 is required for NP incorporation into Mopeia virus Z-induced virus-like particles. J Virol *85*, 3631-3641.

[78] Silva-Ayala, D., Lopez, T., Gutierrez, M., Perrimon, N., Lopez, S., and Arias, C.F. (2013). Genome-wide RNAi screen reveals a role for the ESCRT complex in rotavirus cell entry. Proc Natl Acad Sci USA *110*, 10270-10275.

[79] Snyder, J.C., Samson, R.Y., Brumfield, S.K., Bell, S.D., and Young, M.J. (2013). Functional interplay between a virus and the ESCRT machinery in Archaea. Proc Natl Acad Sci USA.

[80] Spidel, J.L., Craven, R.C., Wilson, C.B., Patnaik, A., Wang, H., Mansky, L.M., and Wills, J.W. (2004). Lysines close to the Rous sarcoma virus late domain critical for budding. J Virol *78*, 10606-10616.

[81] Stange, A., Mannigel, I., Peters, K., Heinkelein, M., Stanke, N., Cartellieri, M., Gottlinger, H., Rethwilm, A., Zentgraf, H., and Lindemann, D. (2005). Characterization of prototype foamy virus gag late assembly domain motifs and their role in particle egress and infectivity. J Virol *79*, 5466-5476.

[82] Strack, B., Calistri, A., Craig, S., Popova, E., and Gottlinger, H.G. (2003). AIP1/ALIX Is a Binding Partner for HIV-1 p6 and EIAV p9 Functioning in Virus Budding. Cell *114*, 689-699.

[83] Tamai, K., Shiina, M., Tanaka, N., Nakano, T., Yamamoto, A., Kondo, Y., Kakazu, E., Inoue, J., Fukushima, K., Sano, K., *et al.* (2012). Regulation of hepatitis C virus secretion by the Hrs-dependent exosomal pathway. Virology *422*, 377-385.

[84] Tandon, R., AuCoin, D.P., and Mocarski, E.S. (2009). Human cytomegalovirus exploits ESCRT machinery in the process of virion maturation. Journal of virology *83*, 10797-10807.

[85] Taylor, G.M., Hanson, P.I., and Kielian, M. (2007). Ubiquitin depletion and dominant-negative VPS4 inhibit rhabdovirus budding without affecting alphavirus budding. Journal of virology *81*, 13631-13639.

[86] Timmins, J., Schoehn, G., Ricard-Blum, S., Scianimanico, S., Vernet, T., Ruigrok, R.W., and Weissenhorn, W. (2003). Ebola Virus Matrix Protein VP40 Interaction with Human Cellular Factors Tsg101 and Nedd4. J Mol Biol *326*, 493-502.

[87] Timmins, J., Scianimanico, S., Schoehn, G., and Weissenhorn, W. (2001). Vesicular release of ebola virus matrix protein VP40. Virology 283, 1-6.

[88] Urata, S., Noda, T., Kawaoka, Y., Morikawa, S., Yokosawa, H., and Yasuda, J. (2007). Interaction of Tsg101 with Marburg virus VP40 depends on the PPPY motif, but not the PT/SAP motif as in the case of Ebola virus, and Tsg101 plays a critical role in the budding of Marburg virus-like particles induced by VP40, NP, and GP. J Virol *81*, 4895-4899.

[89] Urata, S., Noda, T., Kawaoka, Y., Yokosawa, H., and Yasuda, J. (2006). Cellular factors required for Lassa virus budding. J Virol *80*, 4191-4195.

[90] Urata, S., Yasuda, J., and de la Torre, J.C. (2009). The z protein of the new world arenavirus tacaribe virus has bona fide budding activity that does not depend on known late domain motifs. J Virol *83*, 12651-12655.

[91] Urata, S., Yokosawa, H., and Yasuda, J. (2007). Regulation of HTLV-1 Gag budding by Vps4A, Vps4B, and AIP1/Alix. Virol J 4, 66.

[92] Usami, Y., Popov, S., and Gottlinger, H.G. (2007). Potent rescue of human immunodeficiency virus type 1 late domain mutants by ALIX/AIP1 depends on its CHMP4 binding site. J Virol *81*, 6614-6622.

[93] Usami, Y., Popov, S., Popova, E., and Gottlinger, H.G. (2008). Efficient and specific rescue of human immunodeficiency virus type 1 budding defects by a Nedd4-like ubiquitin ligase. J Virol *82*, 4898-4907.

[94] Vana, M.L., Tang, Y., Chen, A., Medina, G., Carter, C., and Leis, J. (2004). Role of Nedd4 and ubiquitination of Rous sarcoma virus Gag in budding of virus-like particles from cells. J Virol *78*, 13943-13953.

[95] VerPlank, L., Bouamr, F., LaGrassa, T.J., Agresta, B., Kikonyogo, A., Leis, J., and Carter, C.A. (2001). Tsg101, a homologue of ubiquitin-conjugating (E2) enzymes, binds the L domain in HIV type 1 Pr55Gag. Proc Natl Acad Sci USA *98*, 7724-7729.

[96] Wang, H., Machesky, N.J., and Mansky, L.M. (2004). Both the PPPY and PTAP motifs are involved in human T-cell leukemia virus type 1 particle release. J Virol *78*, 1503-1512.

[97] Wang, H., Norris, K.M., and Mansky, L.M. (2002). Analysis of bovine leukemia virus gag membrane targeting and late domain function. J Virol *76*, 8485-8493.

[98] Watanabe, T., Sorensen, E.M., Naito, A., Schott, M., Kim, S., and Ahlquist, P. (2007). Involvement of host cellular multivesicular body functions in hepatitis B virus budding. Proc Natl Acad Sci USA *104*, 10205-10210.

[99] Wills, J.W., Cameron, C.E., Wilson, C.B., Xiang, Y., Bennett, R.P., and Leis, J. (1994). An assembly domain of the Rous sarcoma virus Gag protein required late in budding. J Virol *68*, 6605-6618.

[100] Wirblich, C., Bhattacharya, B., and Roy, P. (2006). Nonstructural protein 3 of bluetongue virus assists virus release by recruiting ESCRT-I protein Tsg101. J Virol *80*, 460-473.

[101] Wirblich, C., Tan, G.S., Papaneri, A., Godlewski, P.J., Orenstein, J.M., Harty, R.N., and Schnell, M.J. (2008). PPEY motif within the rabies virus (RV) matrix protein is essential for efficient virion release and RV pathogenicity. J Virol *82*, 9730-9738.

[102] Xiang, Y., Cameron, C.E., Wills, J.W., and Leis, J. (1996). Fine mapping and characterization of the Rous sarcoma virus Pr76gag late assembly domain. J Virol *70*, 5695-5700.

[103] Yasuda, J., and Hunter, E. (1998). A proline-rich motif (PPPY) in the Gag polyprotein of Mason-Pfizer monkey virus plays a maturation-independent role in virion release. J Virol *72*, 4095-4103.

[104] Yasuda, J., Nakao, M., Kawaoka, Y., and Shida, H. (2003). Nedd4 regulates egress of Ebola virus-like particles from host cells. J Virol 77, 9987-9992.

[105] Yuan, B., Campbell, S., Bacharach, E., Rein, A., and Goff, S.P. (2000). Infectivity of moloney murine leukemia virus defective in late assembly events is restored by late assembly domains of other retroviruses. J Virol 74, 7250-7260.

[106] Yuan, B., Li, X., and Goff, S.P. (1999). Mutations altering the moloney murine leukemia virus p12 Gag protein affect virion production and early events of the virus life cycle. Embo J *18*, 4700-4710.

[107] Zhadina, M., and Bieniasz, P.D. (2010). Functional interchangeability of late domains, late domain cofactors and ubiquitin in viral budding. PLoS Pathog *6*, e1001153.

[108] Zhadina, M., McClure, M.O., Johnson, M.C., and Bieniasz, P.D. (2007). Ubiquitin-dependent virus particle budding without viral protein ubiquitination. Proc Natl Acad Sci USA *104*, 20031-20036.

[109] Zhai, Q., Fisher, R.D., Chung, H.Y., Myszka, D.G., Sundquist, W.I., and Hill, C.P. (2008). Structural and functional studies of ALIX interactions with YPX(n)L late domains of HIV-1 and EIAV. Nat Struct Mol Biol *15*, 43-49.

[110] Zhai, Q., Landesman, M.B., Robinson, H., Sundquist, W.I., and Hill, C.P. (2011). Identification and structural characterization of the ALIX-binding late domains of simian immunodeficiency virus SIVmac239 and SIVagmTan-1. J Virol *85*, 632-637.