

Supplementary table 1. Comparisons of the *c*-statistics of the prediction rule between the subgroups stratified by age or sex

Subgroups	Derivation cohort	Validation cohort	<i>P</i> value ^a
	(<i>n</i> = 698)	(<i>n</i> = 702)	
	<i>c</i> -statistics (95% CI)	<i>c</i> -statistics (95% CI)	
Overall population	0.87 (0.82 to 0.92)	0.89 (0.86 to 0.93)	0.39
Subgroup stratified by age			
Younger (<30 years)	0.86 (0.77 to 0.95)	0.90 (0.85 to 0.96)	0.41
Older (≥30 years)	0.87 (0.81 to 0.93)	0.90 (0.85 to 0.93)	0.56
Subgroup stratified by sex			
Men	0.88 (0.82 to 0.95)	0.88 (0.83 to 0.93)	0.90
Women	0.87 (0.81 to 0.94)	0.92 (0.88 to 0.96)	0.25

Abbreviation: CI, confidence interval.

^a*P* value for the differences in *c*-statistics between the derivation and the validation cohorts.

Supplemental Figure 1

Observed and predicted development of end-stage kidney disease by deciles of risk in the derivation cohorts (A) and the derivation cohorts.

Hosmer-Lemeshow χ^2 -statistic = 4.65, d.f. = 8, $P = 0.79$.

Supplementary figure 1

