

Supporting Information

Structure-Activity Relationships for Biodistribution, Pharmacokinetics and Excretion of Atomically Precise Nanoclusters in a Murine Model

O. Andrea Wong¹, Ryan J. Hansen², Thomas W. Ni¹, Christine Heinecke¹, W. Scott Compel¹,
Daniel L. Gustafson² and Christopher J. Ackerson^{1*}

1. Department of Chemistry, Colorado State University, Fort Collins, CO 80523

2. Department of Clinical Sciences, Colorado State University, Fort Collins, CO, 80523

Phone: 970-491-0521

Fax: 970-491-1801

Email: ackerson@mail.colostate.edu

Table of Contents

Fig. S1: Polyacrylamide gel of Au ₁₀₂ (<i>p</i> MBA) ₄₄	S2
Fig. S2: Graph of relative amounts of compounds 1-3 in urine and feces	S2
Fig. S3: Graph of relative amounts of compounds 4,5 , and the Au ₁₀₂ 1:1 exchange compound in urine and feces	S3
Fig. S4-S7: Atomistic models of Au ₂₅ -based compounds	S3-S5
Fig. S8-S10: Atomistic models of Au ₁₀₂ -based compounds	S5-S6
The chemical and biological properties of the compounds discussed in the main text are calculated and predicted from the atomistic models in Fig. S4-S10	
Fig. S11-S13: Percent dosed in tissues of the Au ₂₅ -based compounds	S7-S8
Fig. S14-S15: Percent dosed in tissues of the Au ₁₀₂ -based compounds	S8-S9
Fig. S16: ¹ H NMR of compound 5	S9
Table S1: Information on Au ₁₀₂ -based 1:1 ligand exchange compound	S10
Fig. S17: Blood drug concentration vs. time curves of Au ₁₀₂ -based compounds	S10



Fig. S1 Polyacrylamide gel electrophoresis of $\text{Au}_{102}(\text{pMBA})_{44}$ that shows the purity of the compound.




Fig. S2 Relative amounts of compounds 1-3 in urine and feces at 6, 12, and 24 hours post injection time.




Fig. S3 Relative amounts of compounds **4,5**, and the 1:1 exchange Au_{102} -based compound ($\text{Au}_{102}p\text{MBA}_{44}$ exchanged with $\text{HS-(CH}_2\text{)}_{11}\text{-EG}_4\text{-OH}$) in urine and feces at 6, 12, and 24 hours post injection time.




Fig. S4 $\text{Au}_{25}(\text{GSH})_{18}$, Compound **1**




Fig. S5 $\text{Au}_{25}(\text{GSH})_9(\text{OEG})_9$, ($\text{OEG} = \text{HS-(CH}_2\text{)}_6\text{-EG-OH}$), Compound **2**




Fig. S6 $\text{Au}_{25}(\text{GSH})_6(\text{OEG})_{12}$, ($\text{OEG} = \text{HS-(CH}_2\text{)}_6\text{-EG-OH}$), Compound **3**




Fig. S7 $\text{Au}_{25}(\text{OEG})_{18}$, (OEG = HS-(CH₂)₆-EG-OH)




Fig. S8 $\text{Au}_{102}(p\text{MBA})_{44}$, Compound 4




Fig. S9 Au₁₀₂(pMBA)₂₅(OEG)₁₉, (OEG = HS-(CH₂)₁₁-EG-OH), Compound **5**




Fig. S10 Au₁₀₂(OEG)₄₄, (OEG = HS-(CH₂)₁₁-EG-OH)




Fig. S11 Percent dosed in tissue for compound **1** at noted post-injection time points.




Fig. S12 Percent dosed in tissue for compound **2** at noted post-injection time points.




Fig. S13 Percent dosed in tissue for compound 3 at noted post-injection time points.




Fig. S14 Percent dosed in tissue for compound 4 at noted post-injection time points. Data points of liver are not included.




Fig. S15 Percent dosed in tissue for compound 5 at noted post-injection time points. Data points of liver are not included.




Fig. S16 ^1H NMR of compound 5

Table S1 Information on Au_{102} -based 1:1 exchanged compound

Calculated Stokes Diameter (nm) ^a	Net Surface Charge (e ⁻)	Surface Charge Density (Charge/ \AA^2)	Hydrophobic Surface Area (%)	Expected Clearance Mechanism % Renal / % RES
5.92	22 ⁻	1.8×10^{-4} (-)	74%	40 / 60

Ligand exchange reaction of $\text{Au}_{102}(p\text{MBA})_{44}$ with 23-mercaptop-3,6,9,12-tetraoxatricosan-1-ol [HS-(CH₂)₁₁-EG₄-OH]

A 500 μM solution of $\text{Au}_{102}p\text{MBA}_{44}$ (6.6 μmol , 178 mg in 13.37 mL H₂O) and a 0.1 M solution of HS-(CH₂)₁₁-EG₄-OH (0.79 mmol, 304 mg in 7.90 mL THF) were prepared. For the 1:1 incoming ligand:outgoing ligand reaction: 3 mL of $\text{Au}_{102}p\text{MBA}_{44}$ solution and 0.66 mL of HS-(CH₂)₁₁-EG₄-OH solution were mixed and diluted with H₂O to a final volume of 15 mL. Then the reaction was shaken at rt for 1 h, then the crude product was purified by ultrafiltration spin columns (5000 Da cutoff) and was washed with 3 x 10 mL 1:1 H₂O:MeOH. The remaining orange liquid was placed into a 15 mL conical and lyophilized until dry.

The Au_{102} -based 1:1 exchanged compound was dissolved in 5% DMSO, 5% Tween-80 in 90% D5W solution. The particle solutions were filtered through a 0.45 micron filter, and the dosage concentrations were determined after the filtration step. Approximate dosage concentration for this compound was 3.14×10^{-5} M.




Fig. S17 Blood drug concentration vs. time curves of Au_{102} -based compounds