

SUPPORTING INFORMATION

***O*-Phenyl Carbamate and Phenyl Urea Thiiranes as Selective MMP-2 Inhibitors that Cross the Blood-Brain Barrier**

Major Gooyit,[†] Wei Song,[†] Kiran V. Mahasenan,[†] Katerina Lichtenwalter,[†] Mark A. Suckow,[‡]
Valerie A. Schroeder,[‡] William R. Wolter,[‡] Shahriar Mobashery,[†] and Mayland Chang^{†*}

[†]Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, Indiana
46556, USA

[‡]Freimann Life Sciences Center and Department of Biological Sciences, University of Notre
Dame, Notre Dame, Indiana 46556, USA

Corresponding Author:

Tel: +1-574-631-2965. Fax: +1-574-631-6652. E-mail: mchang@nd.edu

Contents of Supporting Information

Supporting Tables

1. Kinetics parameters for the slow-binding inhibition of MMPs.....S1
2. Concentrations of compound **2** after a single sc dose of compound **5b** to miceS2
3. Concentrations of compound **6b** after a single sc dose of compound **6b** to miceS3

Table S1: Kinetic Parameters for the Slow-binding Inhibition of MMPs

| Inhibitor | Structure | Enzyme | $10^{-3} k_{\text{on}} (\text{M}^{-1}\text{s}^{-1})$ | $10^4 k_{\text{off}} (\text{s}^{-1})$ | $K_i (\mu\text{M})$ |
|-----------|-----------|-----------------------|--|---------------------------------------|---------------------|
| 4a | | MMP-2 | 3.1 ± 0.3 | 4.9 ± 0.2 | 0.16 ± 0.02 |
| | | MMP-9 _{cat} | 4.8 ± 0.4 | 11 ± 3 | 0.23 ± 0.06 |
| | | MMP-14 _{cat} | 0.47 ± 0.02 | 11 ± 1 | 2.4 ± 0.2 |
| 4b | | MMP-2 | 3.4 ± 0.4 | 7.8 ± 2.1 | 0.23 ± 0.07 |
| | | MMP-9 _{cat} | 1.5 ± 0.1 | 14 ± 1 | 0.93 ± 0.02 |
| | | MMP-14 _{cat} | 0.70 ± 0.06 | 19 ± 2 | 2.7 ± 0.3 |
| 4c | | MMP-2 | 1.8 ± 0.1 | 8.7 ± 3.1 | 0.31 ± 0.08 |
| | | MMP-9 _{cat} | 1.2 ± 0.2 | 12 ± 1 | 1.0 ± 0.2 |
| | | MMP-14 _{cat} | 0.54 ± 0.05 | 16 ± 1 | 2.8 ± 0.2 |
| 4d | | MMP-2 | 3.2 ± 0.8 | 9.9 ± 0.4 | 0.48 ± 0.02 |
| | | MMP-9 _{cat} | 1.1 ± 0.1 | 14 ± 2 | 1.3 ± 0.3 |
| | | MMP-14 _{cat} | 0.18 ± 0.01 | 5.0 ± 0.2 | 3.0 ± 0.3 |
| 5a | | MMP-2 | 1.7 ± 0.1 | 6.4 ± 0.3 | 0.39 ± 0.03 |
| | | MMP-9 _{cat} | 0.082 ± 0.007 | 8.6 ± 0.5 | 10 ± 1 |
| | | MMP-14 _{cat} | 0.048 ± 0.005 | 6.0 ± 0.5 | 12 ± 2 |
| 5b | | MMP-2 | 0.69 ± 0.05 | 3.8 ± 0.4 | 0.55 ± 0.08 |
| | | MMP-9 _{cat} | 0.068 ± 0.006 | 8.5 ± 0.7 | 12 ± 2 |
| | | MMP-14 _{cat} | 0.050 ± 0.002 | 8.5 ± 0.3 | 17 ± 1 |
| | | MMP-8 _{cat} | 0.17 ± 0.01 | 26 ± 1 | 15 ± 1 |
| 5c | | MMP-2 | 0.46 ± 0.03 | 3.5 ± 0.3 | 0.76 ± 0.09 |
| | | MMP-9 _{cat} | 0.021 ± 0.003 | 11 ± 1 | 52 ± 8 |
| | | MMP-14 _{cat} | 0.054 ± 0.005 | 14 ± 1 | 27 ± 2 |
| 5d | | MMP-2 | 0.21 ± 0.05 | 1.5 ± 0.6 | 0.69 ± 0.32 |
| 6a | | MMP-2 | 3.1 ± 0.4 | 7.5 ± 1.4 | 0.24 ± 0.05 |
| | | MMP-9 _{cat} | 0.12 ± 0.01 | 12 ± 1 | 9.9 ± 0.6 |
| | | MMP-14 _{cat} | 0.024 ± 0.003 | 10 ± 1 | 19 ± 2 |
| 6b | | MMP-2 | 1.5 ± 0.1 | 6.8 ± 0.7 | 0.44 ± 0.06 |
| | | MMP-8 _{cat} | 0.14 ± 0.01 | 24 ± 2 | 17 ± 2 |
| 6c | | MMP-2 | 1.5 ± 0.2 | 8.1 ± 0.8 | 0.56 ± 0.09 |
| 6d | | MMP-2 | 2.2 ± 0.1 | 8.7 ± 0.5 | 0.42 ± 0.03 |

| Inhibitor | Structure | Enzyme | $10^{-3} k_{\text{on}} (\text{M}^{-1} \text{s}^{-1})$ | $10^4 k_{\text{off}} (\text{s}^{-1})$ | $K_i (\mu\text{M})$ |
|-----------|-----------|-----------------------|---|---------------------------------------|---------------------|
| 7a | | MMP-2 | 5.6 ± 0.3 | 2.8 ± 1.1 | 0.050 ± 0.020 |
| | | MMP-9 _{cat} | 4.7 ± 0.5 | 8.5 ± 1.2 | 0.18 ± 0.03 |
| | | MMP-14 _{cat} | 0.29 ± 0.04 | 7.5 ± 0.7 | 2.6 ± 0.4 |
| | | MMP-8 _{cat} | 0.059 ± 0.005 | 7.7 ± 0.6 | 13 ± 2 |
| 7b | | MMP-2 | 2.5 ± 0.3 | 7.2 ± 1.3 | 0.29 ± 0.06 |
| | | MMP-9 _{cat} | 0.76 ± 0.05 | 6.5 ± 0.7 | 0.86 ± 0.11 |
| | | MMP-14 _{cat} | 0.43 ± 0.02 | 8.2 ± 0.6 | 1.9 ± 0.2 |
| | | MMP-8 _{cat} | 0.24 ± 0.01 | 35 ± 6 | 15 ± 3 |
| 7c | | MMP-2 | 0.91 ± 0.13 | 4.4 ± 0.3 | 0.49 ± 0.08 |
| | | MMP-9 _{cat} | 0.57 ± 0.06 | 6.2 ± 0.6 | 1.1 ± 0.1 |
| | | MMP-14 _{cat} | 0.30 ± 0.02 | 6.4 ± 0.4 | 2.1 ± 0.2 |
| 7d | | MMP-2 | 1.3 ± 0.2 | 9.1 ± 1.0 | 0.68 ± 0.13 |
| | | MMP-9 _{cat} | 0.30 ± 0.05 | 8.0 ± 0.1 | 2.7 ± 0.6 |
| | | MMP-14 _{cat} | 0.25 ± 0.04 | 6.2 ± 0.9 | 2.5 ± 0.4 |

Table S2. Concentrations of Compound **2** after a Single SC Dose of Compound **5b** to Mice at 25 mg/kg

| Time (min) | Compound 2 in Brain ^a | Compound 2 in Plasma ^b |
|------------|---|--|
| 10 | 0.405 ± 0.101 | 1.24 ± 0.676 |
| 20 | 1.35 ± 1.10 | 1.05 ± 0.318 |
| 30 | 0.726 ± 0.0997 | 0.574 ± 0.345 |
| 45 | 0.383 ± 0.0630 | 0.493 ± 0.0967 |
| 60 | 0.471 ± 0.0676 | 0.302 ± 0.0343 |
| 90 | 0.275 ± 0.0159 | 0.185 ± 0.0224 |
| 120 | 0.331 ± 0.0712 | 0.183 ± 0.0565 |
| 180 | 0.252 ± 0.0400 | 0.0949 ± 0.0186 |
| 240 | 0.255 ± 0.0154 | 0.114 ± 0.0147 |

^a Concentration in pmol/mg tissue

^b Concentration in μM

Table S3. Concentrations of Compound **6b** after a Single SC Dose of Compound **6b** to Mice at 25 mg/kg

| Time (min) | Compound 6b in Brain ^a | Compound 6b in Plasma ^b |
|------------|--|---|
| 10 | 0.114 ± 0.0193 | 4.39 ± 0.303 |
| 20 | 0.0779 ± 0.012 | 1.92 ± 0.472 |
| 40 | 0.0627 ± 0.0246 | 1.10 ± 0.475 |
| 90 | 0.0351 ± 0.00751 | 0.530 ± 0.285 |
| 180 | 0.0200 ± 0.00490 | 0.270 ± 0.0252 |
| 300 | 0.0133 ± 0.00337 | 0.0704 ± 0.0506 |

^a Concentration in pmol/mg tissue

^b Concentration in μM