

Appendix 5 (as supplied by the authors): Diagnostic accuracy parameters reported in included studies

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|-------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| Simultaneous and Separate studies | | | | | | | | | |
| Abraham et al. ⁽¹⁾ (2006) At-home | Sensitivity | - | 95.2% | 88.2% | 67% | - | - | - | - |
| | Specificity | - | 52.2% | 63.0% | 78% | - | - | - | - |
| | PPV | - | 64.5% | 60.0% | 53.3% | - | - | - | - |
| | NPV | - | 92.3% | 89.5% | 86.2% | - | - | - | - |
| | LR+ | - | 1.99 | 2.38 | 3.04 | - | - | - | - |
| | LR- | - | 0.09 | 0.19 | 0.423 | - | - | - | - |
| | T+ | - | 20 | 15 | 8 | - | - | - | - |
| | T- | - | 12 | 17 | 25 | - | - | - | - |
| | F+ | - | 11 | 10 | 7 | - | - | - | - |
| | F- | - | 1 | 2 | 4 | - | - | - | - |
| Ayappa et al. ⁽²⁾ (2008) At-home | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | 73% | 73% | 73% | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| | Sensitivity | - | 92.0% | 89.0% | 76.0% | - | - | - | - |
| | Specificity | - | 67.0% | 72.0% | 82.0% | - | - | - | - |
| | PPV | - | 88.0% | 77.0% | 81.0% | - | - | - | - |
| | NPV | - | 75.0% | 85.0% | 78.0% | - | - | - | - |
| | LR+ | - | 2.76 | 3.15 | 4.29 | - | - | - | - |
| | LR- | - | 0.12 | 0.16 | 0.30 | - | - | - | - |
| | T+ | - | 45 | 13 | 25 | - | - | - | - |
| Ayappa et al. ⁽²⁾ 2008 In-lab | T- | - | 12 | 23 | 8 | - | - | - | - |
| | F+ | - | 6 | 9 | 6 | - | - | - | - |
| | F- | - | 4 | 4 | 8 | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| | Sensitivity | - | 98.0% | 97.0% | 91.0% | - | - | - | - |
| | Specificity | - | 76% | 78.0% | 92.0% | - | - | - | - |
| | PPV | - | 91.0% | 82.0% | 92.0% | - | - | - | - |
| | NPV | - | 94.0% | 96.0% | 92.0% | - | - | - | - |
| Garcia-Diaz et al. ⁽³⁾ (2007) In-lab | LR+ | - | 4.12 | 4.38 | 11.85 | - | - | - | - |
| | LR- | - | 0.03 | 0.03 | 0.10 | - | - | - | - |
| | T+ | - | 51 | 36 | 31 | - | - | - | - |
| | T- | - | 16 | 28 | 36 | - | - | - | - |
| | F+ | - | 5 | 8 | 3 | - | - | - | - |
| | F- | - | 1 | 1 | 3 | - | - | - | - |
| | Prevalence | - | 71% | 51% | 47% | - | - | - | - |
| | Accuracy | - | 92% | 87.6% | 91.7% | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|--|-----------------------------|--|-------|----------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | AUC | - | - | 0.97 | 0.99 | - | - | 0.98 | - |
| Garcia-Diaz et al. ⁽³⁾ (2007) At-home | Sensitivity | - | - | 86.4% | 87.5% | - | - | 91.7% | - |
| | Specificity | - | - | 100% | 96.7% | - | - | 94.7% | - |
| | PPV | - | - | 100% | 97.0% | - | - | 92.0% | - |
| | NPV | - | - | 84.0% | 88.0% | - | - | 95.0% | - |
| | LR+ | - | - | Infinity | 27.0 | - | - | 17.0 | - |
| | LR- | - | - | 0.14 | 0.13 | - | - | 0.09 | - |
| | T+ | - | - | - | 30 | - | - | 23 | - |
| | T- | - | - | - | 27 | - | - | 35 | - |
| | F+ | - | - | - | 1 | - | - | 2 | - |
| | F- | - | - | - | 4 | - | - | 2 | - |
| Kuna et al. ⁽⁴⁾ (2005) At-home | Prevalence | - | - | - | 55% | - | - | 40% | - |
| | Accuracy | - | - | - | 91.6% | - | - | 93% | - |
| | AUC | - | - | 0.97 | 0.97 | - | - | 0.98 | - |
| | Sensitivity | - | - | - | 86.7% | - | - | - | - |
| | Specificity | - | - | - | 77.8% | - | - | - | - |
| | PPV | - | - | - | 93.0% | - | - | - | - |
| | NPV | - | - | - | 64.0% | - | - | - | - |
| | LR+ | - | - | - | 3.90 | - | - | - | - |
| | LR- | - | - | - | 0.17 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| Kuna et al. ⁽⁴⁾ (2005) In-lab | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| | Sensitivity | - | - | - | 96.6% | - | - | - | - |
| | Specificity | - | - | - | 100% | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| Polese et al. ⁽⁵⁾ (2009) In-lab | LR+ | - | - | - | 0.03 | - | - | - | - |
| | LR- | - | - | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Santos-Silva et al. ⁽⁶⁾ | Sensitivity | - | 98.0% | - | 97.0% | - | - | 96.0% | - |
| | Specificity | - | 62.0% | - | 74.0% | - | - | 92.0% | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|--------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| (2009) In-lab | PPV | - | 87.0% | - | 78.0% | - | - | 87.0% | - |
| | NPV | - | 93.0% | - | 96.0% | - | - | 98.0% | - |
| | LR+ | - | 2.58 | - | 3.73 | - | - | 12.0 | - |
| | LR- | - | 0.032 | - | 0.041 | - | - | 0.043 | - |
| | T+ | - | 57 | - | 38 | - | - | 28 | - |
| | T- | - | 14 | - | 30 | - | - | 47 | - |
| | F+ | - | 8 | - | 11 | - | - | 4 | - |
| | F- | - | 1 | - | 1 | - | - | 1 | - |
| | Prevalence | - | 72% | - | 49% | - | - | 36% | - |
| | Accuracy | - | 88% | - | 85% | - | - | 93% | - |
| Santos-Silva et al.(⁶) (2009) At-home | Sensitivity | - | 93.0% | - | 85.0% | - | - | 77.0% | - |
| | Specificity | - | 59.0% | - | 80.0% | - | - | 93.0% | - |
| | PPV | - | 85.0% | - | 80.0% | - | - | 81.0% | - |
| | NPV | - | 76.0% | - | 84.0% | - | - | 91.0% | - |
| | LR+ | - | 2.27 | - | 4.25 | - | - | 11.0 | - |
| | LR- | - | 0.12 | - | 0.19 | - | - | 0.25 | - |
| | T+ | - | 53 | - | 33 | - | - | 17 | - |
| | T- | - | 14 | - | 33 | - | - | 54 | - |
| | F+ | - | 9 | - | 8 | - | - | 4 | - |
| | F- | - | 4 | - | 6 | - | - | 5 | - |
| Smith et al.(⁷) (2007) At-home | Sensitivity | - | - | - | 58.3% | - | - | - | - |
| | Specificity | - | - | - | 62.5% | - | - | - | - |
| | PPV | - | - | - | 70.0% | - | - | - | - |
| | NPV | - | - | - | 50.0% | - | - | - | - |
| | LR+ | - | - | - | 1.55 | - | - | - | - |
| | LR- | - | - | - | 0.67 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| Tonelli de Oliveria et al.(⁸) (2009) At-home | Sensitivity | - | 96.15% | 90.7% | 81.3% | - | - | 80.0% | - |
| | Specificity | - | 64.7% | 82.9% | 82.6% | - | - | 92.1% | - |
| | PPV | - | 94.3% | 92.9% | 88.4% | - | - | 85.7% | - |
| | NPV | - | 73.3% | 68.4% | 73.1% | - | - | 88.6% | - |
| | LR+ | - | 2.7 | 5.2 | 4.6 | - | - | 10.1 | - |
| | LR- | - | 0.05 | 0.11 | 0.22 | - | - | 0.21 | - |
| | T+ | - | 100 | 78 | 55 | - | - | 36 | - |
| | T- | - | 11 | 29 | 44 | - | - | 70 | - |
| | F+ | - | 6 | 6 | 9 | - | - | 6 | - |
| | F- | - | 4 | 8 | 13 | - | - | 9 | - |
| Tonelli de Oliveria et al.(⁸) (2009) In-lab | Prevalence | - | 85% | 70% | 52% | - | - | 37% | - |
| | Accuracy | - | 92% | 88% | 82% | - | - | 88% | - |
| | AUC | - | 0.96 | 0.92 | 0.91 | - | - | 0.92 | - |
| | Sensitivity | 95.3% | - | - | - | - | - | - | - |
| | Specificity | 75.0% | - | - | - | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | 3.81 | - | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|--|-----------------------------|--|-------|--------|----------|--------|--------|----------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | LR- | 0.06 | - | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Simultaneous studies | | | | | | | | | |
| Amir et al. ⁽⁹⁾ (2010) | Sensitivity | - | 97.2% | - | 100% | - | - | - | - |
| | Specificity | - | 94.1% | - | 92.7% | - | - | - | - |
| | PPV | - | 92% | - | 80% | - | - | - | - |
| | NPV | - | 94% | - | 100% | - | - | - | - |
| | LR+ | - | 16.47 | - | 1.66 | - | - | - | - |
| | LR- | - | 0.029 | - | infinity | - | - | - | - |
| | T+ | - | 35 | - | 12 | - | - | - | - |
| | T- | - | 16 | - | 38 | - | - | - | - |
| | F+ | - | 1 | - | 3 | - | - | - | - |
| | F- | - | 1 | - | 0 | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Candela et al. ⁽¹⁰⁾ (2005) | Sensitivity | - | - | 97% | 96% | 96% | - | 98% | - |
| | Specificity | - | - | 82% | 94% | 97% | - | 98% | - |
| | PPV | - | - | 93% | 96% | 98% | - | 98% | - |
| | NPV | - | - | 92% | 94% | 95% | - | 98% | - |
| | LR+ | - | - | 5.39 | 16 | 32.0 | - | 49.0 | - |
| | LR- | - | - | 0.037 | 0.042 | 0.041 | - | 0.020 | - |
| | T+ | - | - | 63 | 53 | 53 | - | 45 | - |
| | T- | - | - | 22 | 35 | 35 | - | 45 | - |
| | F+ | - | - | 5 | 2 | 2 | - | 1 | - |
| | F- | - | - | 2 | 2 | 2 | - | 1 | - |
| | Prevalence | - | - | 71% | 60% | 60% | - | 50% | - |
| | Accuracy | - | - | 92.6% | 95% | 95% | - | 98% | - |
| | AUC | - | - | 0.971 | 0.975 | 0.995 | - | 0.996 | - |
| Cheliout-Heraut et al. ⁽¹¹⁾ (2011) | Sensitivity | - | 83.6% | - | - | - | - | - | - |
| | Specificity | - | 81.8% | - | - | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | 4.59 | - | - | - | - | - | - |
| | LR- | - | 0.002 | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Driver et al. ⁽¹²⁾ (2009) | Sensitivity | - | 97.0% | 84.0% | 80.0% | 80.0% | - | 70.0% | - |
| | Specificity | - | 67.0% | 91.0% | 97.0% | 95.0% | - | 100.0% | - |
| | PPV | - | 94.0% | 96.0% | 97.0% | 93.0% | - | 100.0% | - |
| | NPV | - | 80.0% | 71.0% | 76.0% | 84.0% | - | 88.0% | - |
| | LR+ | - | 2.94 | 9.33 | 22.7 | 16.0 | - | infinity | - |
| | LR- | - | 0.04 | 0.18 | 0.21 | 0.21 | - | 0.3 | - |
| | T+ | - | 59 | 44 | 35 | 26 | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|-------|--------|----------|--------|--------|----------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | T- | - | 8 | 19 | 28 | 38 | - | - | - |
| | F+ | - | 4 | 2 | 1 | 2 | - | - | - |
| | F- | - | 2 | 8 | 9 | 7 | - | - | - |
| | Prevalence | - | 84% | 71% | 60% | 45% | - | - | - |
| | Accuracy | - | 92% | 86% | 86% | 88% | - | - | - |
| | AUC | - | 0.94 | 0.944 | 0.926 | 0.889 | - | 0.87 | - |
| Ferre et al. ⁽¹³⁾ (2008) | Sensitivity | - | 93.3% | - | 84.2% | - | - | 55.6% | - |
| | Specificity | - | 86.7% | - | 100% | - | - | 96.4% | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | 7.01 | - | Infinity | - | - | 15.44 | - |
| | LR- | - | 0.08 | - | 0.16 | - | - | 0.46 | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Goodrich et al. ⁽¹⁴⁾ (2009) | Sensitivity | - | 85% | 92% | 87% | 85% | 100% | 88% | - |
| | Specificity | - | 67% | 88% | 82% | 94% | 97% | 100% | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | 2.58 | 7.67 | 4.83 | 14.16 | 33.33 | Infinity | - |
| | LR- | - | 0.22 | 0.09 | 0.16 | 0.16 | - | 0.12 | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Grant et al. ⁽¹⁵⁾ (2009) | Sensitivity | - | 97.0% | - | 97.0% | - | - | - | - |
| | Specificity | - | 73.0% | - | 81.0% | - | - | - | - |
| | PPV | - | - | - | 47.0% | - | - | - | - |
| | NPV | - | - | - | 99.0% | - | - | - | - |
| | LR+ | - | 3.6 | - | 5.1 | - | - | - | - |
| | LR- | - | 0.015 | - | 0.041 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Ng et al. ⁽¹⁶⁾ (2010) | Sensitivity | - | 92.4% | 90.0% | 87.8% | 85.3% | - | - | - |
| | Specificity | - | 85.7% | 86.7% | 94.9% | 95.7% | - | - | - |
| | PPV | - | 96.8% | 91.8% | 94.7% | 93.5% | - | - | - |
| | NPV | - | 70.6% | 83.9% | 88.1% | 89.8% | - | - | - |
| | LR | - | 6.46 | 6.77 | 17.22 | 19.84 | - | - | - |
| | LR | - | 0.089 | 0.115 | 0.126 | 0.154 | - | - | - |
| | T+ | - | 61 | 45 | 36 | 29 | - | - | - |
| | T- | - | 12 | 26 | 37 | 44 | - | - | - |
| | F+ | - | 2 | 4 | 2 | 2 | - | - | - |
| | F- | - | 5 | 5 | 5 | 5 | - | - | - |
| | Prevalence | - | 82% | 62% | 51% | 42% | - | - | - |
| | Accuracy | - | 91% | 89% | 91% | 91% | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|----------|----------|----------|----------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | AUC | | - | - | - | - | - | - | - |
| NG et al. ⁽¹⁷⁾ (2009) | Sensitivity | - | 100% | 97.7% | 94.7% | 96.9% | - | - | - |
| | Specificity | - | 100% | 100% | 100% | 100% | - | - | - |
| | PPV | - | 100% | 100% | 100% | 100% | - | - | - |
| | NPV | - | 100% | 87.5% | 85.7% | 94.7% | - | - | - |
| | LR | - | infinity | infinity | infinity | infinity | - | - | - |
| | LR | - | 0 | 0.23 | 0.05 | 0.031 | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | 1 | 1 | 0.99 | - | - | - | - |
| Nigro et al. ⁽¹⁸⁾ (2010) | Sensitivity | - | 75% | - | 85.7% | - | - | 100% | - |
| | Specificity | - | 86.7% | - | 91.4% | - | - | 98.8% | - |
| | PPV | - | 88.2% | - | 80% | - | - | 77.3% | - |
| | NPV | - | 72.2% | - | 94.1% | - | - | 100% | - |
| | LR+ | - | 5.6 | - | 10 | - | - | 9.8 | - |
| | LR- | - | 0.29 | - | 0.016 | - | - | - | - |
| | T+ | - | 28 | - | 16 | - | - | 17 | - |
| | T- | - | 25 | - | 43 | - | - | 44 | - |
| | F+ | - | 4 | - | 4 | - | - | 5 | - |
| | F- | - | 9 | - | 3 | - | - | 0 | - |
| | Prevalence | - | 57% | - | 29% | - | - | 26% | - |
| | Accuracy | - | 80% | - | 90% | - | - | 92% | - |
| | AUC | - | - | - | - | - | - | - | - |
| Orr et al. ⁽¹⁹⁾ (2006) | Sensitivity | - | - | 83.0% | 82.0% | - | 100% | - | - |
| | Specificity | - | 81.0% | 87.0% | 81% | - | - | 100% | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | 6.38 | 4.31 | - | - | - | - |
| | LR- | - | - | 0.195 | 0.222 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Su S, et al. ⁽²⁰⁾ (2004) | Sensitivity | - | 98.0% | 87.8% | 83.9% | - | - | - | - |
| | Specificity | - | 40.0% | 73.7% | 75.9% | - | - | - | - |
| | PPV | - | 89.1% | 87.8% | 78.8% | - | - | - | - |
| | NPV | - | 80.0% | 73.7% | 81.5% | - | - | - | - |
| | LR+ | - | 1.63 | 3.33 | 3.64 | - | - | - | - |
| | LR- | - | 0.05 | 0.165 | 0.159 | - | - | - | - |
| | T+ | - | 49 | 36 | 26 | - | - | - | - |
| | T- | - | 4 | 14 | 23 | - | - | - | - |
| | F+ | - | 6 | 5 | 7 | - | - | - | - |
| | F- | - | 1 | 5 | 4 | - | - | - | - |
| | Prevalence | - | 88% | 68% | 50% | - | - | - | - |
| | Accuracy | - | 88.3% | 83% | 82% | - | - | - | - |
| | AUC | - | 0.945 | 0.908 | 0.872 | - | - | - | - |
| Sullivan et al. ⁽²¹⁾ (2009) | Sensitivity | 70.0% | 86.0% | - | 57.0% | - | - | 75.0% | - |
| | Specificity | 100% | 82.0% | - | 88.0% | - | - | 97.0% | - |
| | PPV | - | - | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|--|-----------------------------|--|----------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | infinity | 4.78 | - | 4.75 | - | - | 25.0 | - |
| | LR- | 0.3 | 0.17 | - | 0.49 | - | - | 0.26 | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Takaman et al. ²² (2010) | Sensitivity | - | - | - | - | 81% | - | - | - |
| | Specificity | - | - | - | - | 86% | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | - | - | 5.79 | - | - | - |
| | LR- | - | - | - | - | 0.25 | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | 0.87 | - | - | - |
| To et al. ²³ (2009) | Sensitivity | - | 84.0% | - | - | - | - | - | - |
| | Specificity | - | 100% | - | - | - | - | - | - |
| | PPV | - | 100% | - | - | - | - | - | - |
| | NPV | - | 27% | - | - | - | - | - | - |
| | LR+ | - | Infinity | - | - | - | - | - | - |
| | LR- | - | 0.16 | - | - | - | - | - | - |
| | ROC | - | - | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | 0.96 | - | - | - | - | - | - |
| Yagi et al. ²⁴ (2008) | Sensitivity | - | - | - | 95% | - | - | - | - |
| | Specificity | - | - | - | - | - | - | - | - |
| | PPV | - | - | - | 95% | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | - | - | - | - | - | - |
| | LR- | - | - | - | - | - | - | - | - |
| | ROC | - | - | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Separate studies | | | | | | | | | |
| Alvarez et al. ²⁵ (2008) | Sensitivity | - | - | 61.9% | - | - | - | - | - |
| | Specificity | - | - | 95.8% | - | - | - | - | - |
| | PPV | - | - | 92.9% | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|-------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| Campbell et al. ⁽²⁶⁾ (2010) | NPV | - | - | 74.2% | - | - | - | - | - |
| | LR+ | - | - | 14.78 | - | - | - | - | - |
| | LR- | - | - | 0.40 | - | - | - | - | - |
| | T+ | - | - | 13 | - | - | - | - | - |
| | T- | - | - | 23 | - | - | - | - | - |
| | F+ | - | - | 1 | - | - | - | - | - |
| | F- | - | - | 8 | - | - | - | - | - |
| | Prevalence | - | - | 47% | - | - | - | - | - |
| | Accuracy | - | - | 80% | - | - | - | - | - |
| | AUC | - | - | 0.875 | - | - | - | - | - |
| Cilli et al. ⁽²⁷⁾ (2006) | Sensitivity | - | 88% | 90.5% | 93.7% | - | - | - | - |
| | Specificity | - | 50% | 88.9% | 96.9% | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | 1.76 | 8.14 | 4.06 | - | - | - | - |
| | LR- | - | 0.24 | 0.11 | 0.81 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| Danzi-Soares et al. ⁽²⁸⁾ (2012) | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| | Sensitivity | - | 93.0% | - | - | - | - | - | - |
| | Specificity | - | 44.4% | - | - | - | - | - | - |
| | PPV | - | 89.5% | - | - | - | - | - | - |
| | NPV | - | 57.1% | - | - | - | - | - | - |
| | + LR | - | 1.67 | - | - | - | - | - | - |
| | - LR | - | 0.16 | - | - | - | - | - | - |
| | T+ | - | 43 | - | - | - | - | - | - |
| Finkel et al. ⁽²⁹⁾ (2009) | T- | - | 4 | - | - | - | - | - | - |
| | F+ | - | 5 | - | - | - | - | - | - |
| | F- | - | 3 | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|--|-----------------------------|--|----------|--------|--------|----------|--------|----------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Fordyce et al. ⁽³⁰⁾ (2009) | Sensitivity | - | 80.0% | - | - | - | - | - | - |
| | Specificity | - | 100% | - | - | - | - | - | - |
| | PPV | - | 100% | - | - | - | - | - | - |
| | NPV | - | 80.0% | - | - | - | - | - | - |
| | LR+ | - | Infinity | - | - | - | - | - | - |
| | LR- | - | 0.20 | - | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Furukawa et al. ⁽³¹⁾ (2009) | Sensitivity | - | - | - | - | 88% | 77% | 72% | - |
| | Specificity | - | - | - | - | 61% | 81% | 90% | - |
| | PPV | - | - | - | - | 76% | 84.6% | 91.2% | - |
| | NPV | - | - | - | - | 79.1% | 71.4% | 70% | - |
| | LR+ | - | - | - | - | 2.25 | 4.05 | 7.2 | - |
| | LR- | - | - | - | - | 0.19 | 0.28 | 0.31 | - |
| | ROC | - | - | - | - | 38 | 33 | 31 | - |
| | T+ | - | - | - | - | 19 | 25 | 28 | - |
| | T- | - | - | - | - | 12 | 6 | 3 | - |
| | F+ | - | - | - | - | 5 | 10 | 12 | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | 75.7% | 78.4% | 79% | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Gjevre et al. ⁽³²⁾ (2011) | Sensitivity | - | 90% | 75% | 62.5% | 46% | - | 28.1% | - |
| | Specificity | - | 66% | 86.7% | 93.3% | 100% | - | 100% | - |
| | PPV | - | 83% | 92.3% | 95.2% | 100% | - | 100% | - |
| | NPV | - | 75% | 61.9% | 53.9% | 46.9% | - | 39.5% | - |
| | LR+ | - | 2.26 | 5.63 | 9.32 | Infinity | - | Infinity | - |
| | LR- | - | 0.156 | 0.288 | 0.402 | - | - | - | - |
| | T+ | - | 29 | 24 | 20 | - | - | - | - |
| | T- | - | 9 | 13 | 14 | - | - | - | - |
| | F+ | - | 6 | 2 | 1 | - | - | - | - |
| | F- | - | 3 | 8 | 12 | - | - | - | - |
| | Prevalence | - | 68% | 68% | 68% | - | - | - | - |
| | Accuracy | - | 80% | 78.7% | 72.3% | - | - | - | - |
| | AUC | - | - | - | - | - | - | 0.88 | - |
| Hernandez et al. ⁽³³⁾ (2007) | Sensitivity | - | - | 83.3% | - | - | - | - | - |
| | Specificity | - | - | 78.9% | - | - | - | - | - |
| | PPV | - | - | 91.8% | - | - | - | - | - |
| | NPV | - | - | 62.5% | - | - | - | - | - |
| | LR+ | - | - | 3.95 | - | - | - | - | - |
| | LR- | - | - | 0.21 | - | - | - | - | - |
| | T+ | - | - | 45 | - | - | - | - | - |
| | T- | - | - | 15 | - | - | - | - | - |
| | F+ | - | - | 4 | - | - | - | - | - |
| | F- | - | - | 9 | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|---|-----------------------------|--|-------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | Accuracy AUC | - | - | - | - | - | - | - | - |
| | - | - | - | - | - | - | - | - | - |
| Masa et al. ⁽³⁴⁾ (2011) | Sensitivity | - | - | - | - | - | - | 52% | - |
| | Specificity | - | - | - | - | - | - | 76% | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | - | - | - | - | 2.18 | - |
| | LR- | - | - | - | - | - | - | 0.63 | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Masdeu et al. ⁽³⁵⁾ (2010) | Sensitivity | - | - | 86% | 81% | - | - | - | - |
| | Specificity | - | - | 84% | 94% | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | 5.375 | 13.5 | - | - | - | - |
| | LR- | - | - | 0.166 | 0.202 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | 85% | 86% | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Quintana-Gallego et al. ⁽³⁶⁾ (2004) | Sensitivity | - | 82.5% | 79.3% | 68.4% | - | - | - | - |
| | Specificity | - | 88.6% | 97.8% | 94.6% | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | 7.24 | 36.04 | 12.67 | - | - | - | - |
| | LR- | - | 0.20 | 0.21 | 0.33 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | 0.986 | 0.907 | 0.862 | - | - | - | - |
| Shrivastava et al. ⁽³⁷⁾ (2006) | Sensitivity | - | - | - | 86.4% | - | - | - | - |
| | Specificity | - | - | - | 80.8% | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | - | 4.50 | - | - | - | - |
| | LR- | - | - | - | 0.17 | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| Skomro et al. ⁽³⁸⁾ (2005) | Sensitivity | - | 92.0% | - | - | - | - | - | - |
| | Specificity | - | 67.0% | - | - | - | - | - | - |
| | PPV | - | 88.0% | - | - | - | - | - | - |

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. CMAJ 2013; DOI:10.1503/cmaj.130952.

Copyright © 2013 Canadian Medical Association or its licensors

| Study | Diagnostic Accuracy Measure | Sleep Study Apnea-Hypopnea Index Cut-off | | | | | | | |
|--------------------------------------|-----------------------------|--|-------|--------|--------|--------|--------|--------|--------|
| | | ≤5/hr | ≥5/hr | ≥10/hr | ≥15/hr | ≥20/hr | ≥25/hr | ≥30/hr | ≥50/hr |
| | NPV | - | 75.0% | - | - | - | - | - | - |
| | LR+ | - | 2.79 | - | - | - | - | - | - |
| | LR- | - | 0.12 | - | - | - | - | - | - |
| | T+ | - | 22 | - | - | - | - | - | - |
| | T- | - | 6 | - | - | - | - | - | - |
| | F+ | - | 3 | - | - | - | - | - | - |
| | F- | - | 2 | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |
| To et al. ⁽³⁹⁾ (2012) | Sensitivity | - | - | 89% | - | - | - | - | - |
| | Specificity | - | - | 50% | - | - | - | - | - |
| | PPV | - | - | - | - | - | - | - | - |
| | NPV | - | - | - | - | - | - | - | - |
| | LR+ | - | - | 1.78 | - | - | - | - | - |
| | LR- | - | - | 0.22 | - | - | - | - | - |
| | T+ | - | - | - | - | - | - | - | - |
| | T- | - | - | - | - | - | - | - | - |
| | F+ | - | - | - | - | - | - | - | - |
| | F- | - | - | - | - | - | - | - | - |
| | Prevalence | - | - | - | - | - | - | - | - |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | 0.89 | - | - | - | - | - |
| Yin et al. ⁽⁴⁰⁾ (2006) | Sensitivity | - | 100% | - | 93.8% | - | - | 79.2% | 90.0% |
| | Specificity | - | - | - | 25.0% | - | - | 70.0% | 97.1% |
| | PPV | - | 93.2% | - | 76.9% | - | - | 76.0% | 90.0% |
| | NPV | - | - | - | 60.0% | - | - | 73.7% | 97.1% |
| | LR+ | - | - | - | 1.25 | - | - | 2.64 | 31.03 |
| | LR- | - | - | - | 0.228 | - | - | 0.297 | 0.103 |
| | T+ | - | - | - | 30 | - | - | 19 | 9 |
| | T- | - | - | - | 3 | - | - | 14 | 33 |
| | F+ | - | - | - | 9 | - | - | 6 | 1 |
| | F- | - | - | - | 2 | - | - | 5 | 1 |
| | Prevalence | - | - | - | 72% | - | - | 54% | 23% |
| | Accuracy | - | - | - | - | - | - | - | - |
| | AUC | - | - | - | - | - | - | - | - |

AUC = area under the curve; F+ = false positive; F- = false negative; LR+ = positive likelihood ratio; LR- = negative likelihood ratio; NPV = negative predictive value; PPV = positive predictive value; T+ = true positive; T- = true negative;

References

- (1) Abraham WT, Trupp RJ, Phillips B, Bourge RC, Bailey B, Harding SM, et al. Validation and clinical utility of a simple in-home testing tool for sleep-disordered breathing and arrhythmias in heart failure: results of the Sleep Events, Arrhythmias, and Respiratory Analysis in Congestive Heart Failure (SEARCH) study. *Congest Heart Fail* 2006;12(5):241-7.
- (2) Ayappa I, Norman RG, Seelall V, Rapoport DM. Validation of a self-applied unattended monitor for sleep disordered breathing. *J Clin Sleep Med* 2008;4(1):26-37.

- (3) Garcia-Diaz E, Quintana-Gallego E, Ruiz A, Carmona-Bernal C, Sanchez-Armengol A, Botebol-Benhamou G, et al. Respiratory polygraphy with actigraphy in the diagnosis of sleep apnea-hypopnea syndrome. *Chest* 2007;131(3):725-32.
- (4) Kuna ST, Seeger T, Brendel M. Intra-subject comparison of polysomnography and a type 3 portable monitor [abstract]. *Sleep* 2005;28(Suppl):A324 (Abstract no. 0956).
- (5) Polese JF, Santos SR, Sartori DE, Carlos VA, Kobayashi RF, Ferrari PMP, et al. Validation of a portable monitoring system for the diagnosis of obstructive sleep apnea in elderly patients [abstract]. *Sleep Medicine* 2009;10(Suppl 2):S22 (Abstract no. 078).
- (6) Santos-Silva R, Sartori DE, Truksinas V, Truksinas E, Alonso FF, Tufik S, et al. Validation of a portable monitoring system for the diagnosis of obstructive sleep apnea syndrome. *Sleep* 2009;32(5):629-36.
- (7) Smith LA, Chong DW, Vennelle M, Denvir MA, Newby DE, Douglas NJ. Diagnosis of sleep-disordered breathing in patients with chronic heart failure: evaluation of a portable limited sleep study system. *J Sleep Res* 2007;16(4):428-35.
- (8) Tonelli de Oliveira AC, Martinez D, Vasconcelos LF, Goncalves SC, Lenz MC, Fuchs SC, et al. Diagnosis of obstructive sleep apnea syndrome and its outcomes with home portable monitoring. *Chest* 2009;135(2):330-6.
- (9) Amir O, Barak-Shinar D, Amos Y, MacDonald M, Pittman S, White DP. An automated sleep-analysis system operated through a standard hospital monitor. *J Clin Sleep Med* 2010;6(1):59-63.
- (10) Candela A, Hernandez L, Asensio S, Sanchez-Paya J, Vila J, Benito N, et al. Validation of a respiratory polygraphy system in the diagnosis of sleep apnea syndrome. *Arch Bronconeumol* 2005;41(2):71-7.
- (11) Cheliout-Heraut F, Senny F, Djouadi F, Ouayoun M, Bour F. Obstructive sleep apnoea syndrome: comparison between polysomnography and portable sleep monitoring based on jaw recordings. *Neurophysiol Clin* 2011;41(4):191-8.
- (12) Driver HS, Pereira EJ, Bjerring K, Toop F, Stewart SC, Munt PW, et al. Validation of the MediByte(R) type 3 portable monitor compared with polysomnography for screening of obstructive sleep apnea. *Can Respir J* 2011;18(3):137-43.
- (13) Ferre A, Grau M, Jurado M, Romero O, Saheb A, Bartolome M, et al. Evaluation of a new simplified polysomnography system for the diagnosis of sleep disordered breathing [abstract]. *J Sleep Res* 2008;17(Suppl 1):234 (Abstract no. P441).

Appendix to: El Shayeb M, Topfer LA, Stafinski T, et al. A comparison of level 3 portable sleep tests to level 1 polysomnography in the diagnosis of sleep-disordered breathing: a systematic review and meta-analysis. *CMAJ* 2013; DOI:10.1503/cmaj.130952.
Copyright © 2013 Canadian Medical Association or its licensors

- (14) Goodrich S, Orr WC. An investigation of the validity of the Lifeshirt in comparison to standard polysomnography in the detection of obstructive sleep apnea. *Sleep Med* 2009;10(1):118-22.
- (15) Grant B. Diagnostic accuracy of home sleep studies for sleep apnea [abstract]. *Sleep* 2009;32(Suppl):A232 (Abstract no. 713).
- (16) Ng SS, Chan TO, To KW, Ngai J, Tung A, Ko FW, et al. Validation of Embletta portable diagnostic system for identifying patients with suspected obstructive sleep apnoea syndrome (OSAS). *Respirology* 2010;15(2):336-42.
- (17) Ng SS, Chan TO, To KW, Ngai J, Tung A, Ko FW, et al. Validation of a portable recording device (ApneaLink) for identifying patients with suspected obstructive sleep apnoea syndrome. *Intern Med J* 2009;39(11):757-62.
- (18) Nigro CA, Serrano F, Aimaretti S, Gonzalez S, Codinardo C, Rhodius E. Utility of ApneaLink for the diagnosis of sleep apnea-hypopnea syndrome. *Medicina (B Aires)* 2010;70(1):53-9.
- (19) Orr WC, Goodrich S. An investigation of the accuracy of the Lifeshirt in comparison to standard polysomnography [abstract]. *Sleep* 2006;29(Suppl):A348 (Abstract no. 1020).
- (20) Su S, Baroody FM, Kohrman M, Suskind D. A comparison of polysomnography and a portable home sleep study in the diagnosis of obstructive sleep apnea syndrome. *Otolaryngol Head Neck Surg* 2004;131(6):844-50.
- (21) Sullivan GE, Morehouse R, Savoy A. Comparison of synchronized level 1 and level 3 sleep studies [abstract]. *Vigilance* 2009;19(Suppl):(Abstract no. P074).
- (22) Takama N, Kurabayashi M. Effectiveness of a portable device and the need for treatment of mild-to-moderate obstructive sleep-disordered breathing in patients with cardiovascular disease. *J Cardiol* 2010;56(1):73-8.
- (23) To KW, Chan WC, Chan TO, Tung A, Ngai J, Ng S, et al. Validation study of a portable monitoring device for identifying OSA in a symptomatic patient population. *Respirology* 2009;14(2):270-5.
- (24) Yagi H, Nakata S, Tsuge H, Yasuma F, Noda A, Morinaga M, et al. Significance of a screening device (Apnomonitor 5) for sleep apnea syndrome. *Auris Nasus Larynx* 2009;36(2):176-80.

- (25) Alonso Alvarez MD, Santos JT, Guevara JC, Martinez MG, Pascual LR, Banuelos JL, et al. Reliability of home respiratory polygraphy for the diagnosis of sleep apnea-hypopnea syndrome: analysis of costs. *Archivos de Bronconeumologia* 2008;44(1):22-8.
- (26) Campbell AJ, Neill AM. Home set-up polysomnography in the assessment of suspected obstructive sleep apnea. *J Sleep Res* 2011;20(1 Pt 2):207-13.
- (27) Cilli A, Erogullari I, Turhan M, Ozbudak O, Ozdemir T, Dinc O. Comparison of Embletta portable device versus Embla for diagnosing the obstructive sleep apnea [abstract]. *J Sleep Res* 2006;15(Suppl 1):152.
- (28) Danzi-Soares NJ, Genta PR, Nerbass FB, Pedrosa RP, Soares FS, Cesar LA, et al. Obstructive sleep apnea is common among patients referred for coronary artery bypass grafting and can be diagnosed by portable monitoring. *Coron Artery Dis* 2012;23(1):31-8.
- (29) Finkel KJ, Searleman AC, Tymkew H, Tanaka CY, Saager L, Safer-Zadeh E, et al. Prevalence of undiagnosed obstructive sleep apnea among adult surgical patients in an academic medical center. *Sleep Med* 2009;10(7):753-8.
- (30) Fordyce L, Samuels CH, Oram C, Wallins B. A retrospective, observational study showing patients with a normal level III sleep study and normal OSA pretest probability factors may still require additional investigations [abstract]. *Vigilance* 2009;19(Suppl):(Abstract no. P075).
- (31) Furukawa T, Suzuki M, Funatogawa I, Isshiki T, Miyazawa Y, Teramoto T, et al. Screening method for severe sleep-disordered breathing in hypertensive patients without daytime sleepiness. *J Cardiol* 2009;53(1):79-85.
- (32) Gjever JA, Taylor-Gjever RM, Skomro R, Reid J, Fenton M, Cotton D. Comparison of polysomnographic and portable home monitoring assessments of obstructive sleep apnea in Saskatchewan women. *Can Respir J* 2011;18(5):271-4.
- (33) Hernandez L, Torrella M, Roger N, Llunell A, Ballester E, Quinto L, et al. Management of sleep apnea: concordance between nonreference and reference centers. *Chest* 2007;132(6):1853-7.
- (34) Masa JF, Corral J, Pereira R, Duran-Cantolla J, Cabello M, Hernandez-Blasco L, et al. Therapeutic decision-making for sleep apnea and hypopnea syndrome using home respiratory polygraphy: a large multicentric study. *Am J Respir Crit Care Med* 2011;184(8):964-71.

- (35) Masdeu MJ, Ayappa I, Hwang D, Mooney AM, Rapoport DM. Impact of clinical assessment on use of data from unattended limited monitoring as opposed to full-in lab PSG in sleep disordered breathing. *J Clin Sleep Med* 2010;6(1):51-8.
- (36) Quintana-Gallego E, Villa-Gil M, Carmona-Bernal C, Botebol-Benhamou G, Martinez-Martinez A, Sanchez-Armengol A, et al. Home respiratory polygraphy for diagnosis of sleep-disordered breathing in heart failure. *Eur Respir J* 2004;24(3):443-8.
- (37) Shrivastava D, Aurora NR, Gathoo M, Hardin K. Home diagnostic portable monitoring: an effective screening tool in a community-based primary care population [abstract]. *Sleep* 2006;29(Suppl):A192 (Abstract no. 565).
- (38) Skomro R, Cotton D, Gjevre J, Hill F, King L, Mink J, et al. Diagnosis of obstructive sleep apnea using unattended level III monitoring [abstract]. *Sleep* 2005;28(Suppl):A164 (Abstract no. 0483).
- (39) To KW, Chan WC, Chan TO, Ngai J, Tung A, Ng S, et al. Comparison of empirical continuous positive airway pressure (CPAP) treatment versus initial portable sleep monitoring followed by CPAP treatment for patients with suspected obstructive sleep apnoea. *Intern Med J* 2012;42(6):e107-e114.
- (40) Yin M, Miyazaki S, Ishikawa K. Evaluation of type 3 portable monitoring in unattended home setting for suspected sleep apnea: factors that may affect its accuracy. *Otolaryngol Head Neck Surg* 2006;134(2):204-9.

