

**A Mechanistic Framework for In Vitro – In Vivo Extrapolation of Liver
Membrane Transporters: Prediction of Drug-Drug Interaction between
Rosuvastatin and Cyclosporine.**

M. Jamei^{1*}, F. Bajot^{1,†}, S. Neuhoff¹, Z. Barter¹, J. Yang^{1,‡}, A. Rostami-Hodjegan^{1,2}
and K. Rowland-Yeo¹

¹*Simcyp Limited, Blades Enterprise Centre, Sheffield, UK;* ²*Centre of Applied Pharmacokinetic
Research, the School of Pharmacy and Pharmaceutical Sciences, the University of
Manchester, Manchester, UK*

[†] Current position: Computational Toxicology, British American Tobacco, Southampton, UK

[‡] Current position: Clinical Pharmacology Modelling and Simulation, GSK, Shanghai, China

Supplementary Materials 2

Simulated versus observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 10, 20, 40 and 80 mg, respectively.

Table S1 – Details of the single dose clinical studies used for Rosuvastatin performance verification

| PK Study | Dose [mg] | Age range [years] | n | % female | Reference |
|-----------------|------------------|--------------------------|----------|-----------------|------------------|
| 1 | 10 | 22-42 | 11 | 0 | [1] |
| 2 | 10 | 31-60 | 18 | 0 | [2] |
| 3 | 20 | 31-60 | 9 | 0 | [2] |
| 4 | 40 | 31-60 | 9 | 0 | [2] |
| 5 | 40 | 21-51 | 10 | 0 | [3] |
| 6 | 40 | 21-39 | 36 | 13.9 | [4] |
| 7 | 80 | 22-44 | 11 | 0 | [5] |
| 8 | 80 | 25-56 | 14 | 0 | [1] |
| 9 | 80 | 29-51 | 14 | 0 | [6] |
| 10 | 80 | 31-60 | 18 | 0 | [2] |
| 11 | 80 | 35-47 | 20 | 15 | [7] |

PK study 1




Fig. S1 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 10 mg. The grey thin lines represent simulated individual trials (10) of 11 male subjects (22-42 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=110). The circles denote mean values from the clinical study by Cooper *et al.* 2003 [1].

PK study 2




Fig. S2 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 10 mg. The grey thin lines represent simulated individual trials (10) of 18 male subjects (31-60 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=180). The circles denote mean values from the clinical study by Martin *et al.* 2003 [2].

PK study 3




Fig. S3 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 20 mg. The grey thin lines represent simulated individual trials (10) of 9 male subjects (31-60 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=90). The circles denote mean values from the clinical study by Martin *et al.* 2003 [2].

PK study 4




Fig. S4 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 40 mg. The grey thin lines represent simulated individual trials (10) of 9 male subjects (31-60 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=90). The circles denote mean values from the clinical study by Martin *et al.* 2003 [2].

PK study 5




Fig. S5 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 40 mg. The grey thin lines represent simulated individual trials (10) of 10 male subjects (21-51 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=100). The circles denote mean values from the clinical study by Martin *et al.* 2003 [3].

PK study 6




Fig. S6 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 40 mg. The grey thin lines represent simulated individual trials (10) of 36 subjects (0.139% female, 21-39 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=360). The circles denote mean values from the clinical study by Lee *et al.* 2005 [4].

PK study 7




Fig. S7 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 80 mg. The grey thin lines represent simulated individual trials (10) of 11 male subjects (22-44 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=110). The circles denote mean values from the clinical study by Cooper *et al.* 2003 [5].

PK study 8




Fig. S8 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 80 mg. The grey thin lines represent simulated individual trials (10) of 14 male subjects (25-56 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=140). The circles denote mean values from the clinical study by Cooper *et al.* 2003 [1].

PK study 9




Fig. S9 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 80 mg. The grey thin lines represent simulated individual trials (10) of 14 male subjects (29-51 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=140). The circles denote mean values from the clinical study by Cooper *et al.* 2002 [6].

PK study 10




Fig. S10 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 80 mg. The grey thin lines represent simulated individual trials (10) of 18 male subjects (31-60 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=180). The circles denote mean values from the clinical study by Martin *et al.* 2003 [2].

PK study 11




Fig. S11 - Simulated *versus* observed plasma concentration-time profiles of rosuvastatin in healthy volunteers following the oral administration of 80 mg. The grey thin lines represent simulated individual trials (10) of 20 subjects (0.15% female, 35-47 years) and the solid black line represents the simulated mean of the healthy volunteers population (n=200). The circles denote mean values from the clinical study by Schneck *et al.* 2004 [7].

References

1. Cooper KJ, Martin PD, Dane AL, Warwick MJ, Schneck DW, Cantarini MV. Effect of itraconazole on the pharmacokinetics of rosuvastatin. *Clin Pharmacol Ther.* 2003;73(4):322-9.
2. Martin PD, Warwick MJ, Dane AL, Cantarini MV. A double-blind, randomized, incomplete crossover trial to assess the dose proportionality of rosuvastatin in healthy volunteers. *Clin Ther.* 2003;25(8):2215-24.
3. Martin PD, Warwick MJ, Dane AL, Brindley C, Short T. Absolute oral bioavailability of rosuvastatin in healthy white adult male volunteers. *Clin Ther.* 2003;25(10):2553-63.
4. Lee E, Ryan S, Birmingham B, Zalikowski J, March R, Ambrose H, et al. Rosuvastatin pharmacokinetics and pharmacogenetics in white and Asian subjects residing in the same environment. *Clin Pharmacol Ther.* 2005;78(4):330-41.
5. Cooper KJ, Martin PD, Dane AL, Warwick MJ, Raza A, Schneck DW. The effect of erythromycin on the pharmacokinetics of rosuvastatin. *Eur J Clin Pharmacol.* 2003;59(1):51-6.
6. Cooper KJ, Martin PD, Dane AL, Warwick MJ, Schneck DW, Cantarini MV. The effect of fluconazole on the pharmacokinetics of rosuvastatin. *Eur J Clin Pharmacol.* 2002;58(8):527-31.
7. Schneck DW, Birmingham BK, Zalikowski JA, Mitchell PD, Wang Y, Martin PD, et al. The effect of gemfibrozil on the pharmacokinetics of rosuvastatin. *Clin Pharmacol Ther.* 2004;75(5):455-63.