

**Systemic delivery of miR-126 by miRNA-loaded Bubble liposomes for the
treatment of hindlimb ischemia**

Yoko Endo-Takahashi^a, Yoichi Negishi^{a}, Arisa Nakamura^a, Saori Ukai^a, Kotomi Ooku^a, Yusuke Oda^b, Katsutoshi Sugimoto^c, Fuminori Moriyasu^c, Norio Takagi^d, Ryo Suzuki^b, Kazuo Maruyama^b,
Yukihiko Aramaki^a*

^aDepartment of Drug Delivery and Molecular Biopharmaceutics, School of Pharmacy, Tokyo University of Pharmacy and Life Sciences, 1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan

^bLaboratory of Drug and Gene Delivery Research, Faculty of Pharma-Sciences, Teikyo University, 2-11-1 Kaga, Itabashi-ku, Tokyo 173-8605, Japan

^cDepartment of Gastroenterology and Hepatology, Tokyo Medical University, 6-7-1 Nishishinjuku, Shinjuku-ku, Tokyo 160-0023, Japan

^dDepartment of Applied Biochemistry, School of Pharmacy, Tokyo University of Pharmacy and Life Sciences, 1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan

Corresponding author: Yoichi Negishi

Tel. & Fax: +81-42-676-3183

E-mail: negishi@toyaku.ac.jp

Supplementary Figure S1.

Full scans of the blots shown in Figure 1b.

