

Supplemental Table 1. Baseline characteristics stratified by baseline body mass index in Korean HD patients^{a,b}

	Body mass index (kg/m ²)						
	Total	≤18.5	18.6–20.0	20.1–21.5	21.6–23.0	23.1–25.0	>25.0
n	20,818	3,086	3,817	4,401	3,758	3,318	2,438
Age at entry (yrs)	54±14	53±16	53±15	54±14	55±13	56±13	55±13
Female (%)	43	55	45	39	37	38	47
Diabetes (%)	39	31	35	39	42	42	44
HD vintage (yrs)	2.4±3.5	3.0±4.1	2.9±3.8	2.6±3.7	2.2±3.2	1.9±3.0	1.6±2.5
Hemoglobin (g/dL)	9.6±1.4	9.5±1.4	9.6±1.4	9.6±1.4	9.7±1.4	9.8±1.4	9.8±1.4
Albumin (g/dL)	3.6±0.6	3.5±0.6	3.6±0.6	3.6±0.6	3.6±0.6	3.6±0.6	3.6±0.6
Creatinine (mg/dL)	9.6±6.5	8.8±6.2	9.4±6.1	9.7±6.6	9.7±6.2	9.9±6.7	10.1±7.4
Single pool Kt/V	1.37±0.37	1.51±0.36	1.44±0.37	1.38±0.38	1.34±0.38	1.29±0.33	1.25±0.35
nPNA (g/kg/day)	0.93±0.30	0.86±0.28	0.91±0.29	0.92±0.29	0.94±0.29	0.96±0.30	1.01±0.32

^aHD = hemodialysis; nPNA = normalized protein nitrogen appearance

^bCategorical variables are given as percentage; continuous variables as mean ± standard deviation. Conversion factors for units:

hemoglobin and albumin in g/dL to g/L, ×10; creatinine in mg/dL to μmol/L, ×88.4