

## **Supplemental Material**

### **Associations of PM<sub>2.5</sub> Constituents and Sources with Hospital Admissions: Analysis of Four Counties in Connecticut and Massachusetts (USA) for Persons ≥ 65 Years of Age**

Michelle L. Bell, Keita Ebisu, Brian P. Leaderer, Janneane F. Gent, Hyung Joo Lee, Petros Koutrakis, Yun Wang, Francesca Dominici, and Roger D. Peng

#### **Table of contents**

<b>Supplemental Material, Table S1.</b> Summary statistics of data stratified by county.....	2
<b>Supplemental Material, Table S2.</b> Correlation of PM <sub>2.5</sub> sources and PM <sub>2.5</sub> chemical components.....	3
<b>Supplemental Material, Figure S1.</b> Location of monitors and county in Connecticut.....	4
<b>Supplemental Material, Figure S2.</b> Percent increase in risk of respiratory hospital admissions per IQR increase in exposure to PM <sub>2.5</sub> chlorine, with adjustment by other PM <sub>2.5</sub> components.....	5


**Supplemental Material, Table S1.** Summary statistics of data stratified by county.

County	Cause of Hospital Admissions	Mean $\pm$ SD (admissions /day)	Median (admissions/day)	IQR (admissions /day)	Total admissions across study period	Population at risk/day
Fairfield	CVD	18.8 $\pm$ 5.7	18	8	24,672	89,290
	Respiratory	5.6 $\pm$ 2.9	5	3	7,324	
Hartford	CVD	21.3 $\pm$ 5.9	21	8	27,846	104,558
	Respiratory	8.0 $\pm$ 3.7	7	5	10,512	
New Haven	CVD	21.6 $\pm$ 6.0	21	9	28,258	93,683
	Respiratory	8.8 $\pm$ 4.0	8	5	11,565	
Hampden	CVD	11.5 $\pm$ 4.2	11	5	15,055	46,368
	Respiratory	3.6 $\pm$ 2.2	3	3	4,768	

**Supplemental Material, Table S2.** Correlation of PM<sub>2.5</sub> sources and PM<sub>2.5</sub> chemical components.

	PM <sub>2.5</sub> Sources					PM <sub>2.5</sub> Elements										
	Motor vehicles	Oil	Road dust	Sea salt	Regional source	Al	BC	Br	Ca	Cl	Ni	K	S	Si	Ti	V
Zn	0.93	0.40	0.18	0.16	0.02	0.19	0.60	0.49	0.46	0.13	0.54	0.60	0.25	0.20	0.24	0.38
V	0.29	0.60	0.28	0.22	0.08	0.25	0.50	0.24	0.39	0.20	0.85	0.28	0.23	0.25	0.21	
Ti	0.20	0.04	0.63	0.09	0.19	0.65	0.37	0.16	0.63	0.08	0.22	0.36	0.30	0.65		
Si	0.12	0.07	0.89	0.07	0.16	0.96	0.43	0.14	0.81	0.07	0.23	0.44	0.27			
S	0.17	0.24	0.16	-0.02	0.95	0.34	0.34	0.38	0.33	-0.02	0.26	0.38				
K	0.56	0.39	0.36	0.14	0.20	0.44	0.47	0.52	0.52	0.11	0.39					
Ni	0.49	0.67	0.25	0.26	0.06	0.22	0.55	0.37	0.43	0.24						
Cl	0.10	0.09	0.03	0.94	-0.07	0.09	0.12	0.14	0.20							
Ca	0.42	0.13	0.87	0.20	0.17	0.78	0.66	0.28								
Br	0.48	0.38	0.09	0.17	0.25	0.15	0.36									
BC	0.61	0.22	0.53	0.12	0.20	0.44										
Al	0.11	0.05	0.86	0.08	0.24											
Regional	-0.04	0.02	0.06	-0.08												
Sea salt	0.13	0.13	0.01													
Road dust	0.12	0.00														
Oil	0.24															

**Supplemental Material, Figure S1.** Location of monitors and counties in Connecticut and Massachusetts.



**Supplemental Material, Figure S2.** Percent change in risk of respiratory hospital admissions per IQR increase in exposure to PM<sub>2.5</sub> chlorine, with adjustment by other PM<sub>2.5</sub> components. Note: The point represents the central estimate; the horizontal line represents the 95% confidence interval.

