

Supplemental Table 4

SNP position	r^2	r^2 p-value	
3581	0.613	0.001	* linked to <i>ICAM-1</i> ^{Kilifi}
-9	0.578	0.001	* linked to <i>ICAM-1</i> ^{Kilifi}
3704	0.565	0.001	<i>ICAM-1</i> ^{Kilifi}
13081	0.525	0.002	Top 5% of distributions
13416	0.376	0.015	
15387	0.314	0.030	
701	0.313	0.030	
13933	0.313	0.030	
-333	0.300	0.035	
596	0.300	0.035	
14500	0.268	0.048	
15567	0.248	0.059	
175	0.230	0.071	
13529	0.230	0.071	
3829	0.220	0.078	
-1067	0.217	0.080	
15197	0.217	0.080	
13632	0.210	0.086	
13082	0.207	0.089	
445	0.185	0.109	
13890	0.165	0.133	
13611	0.154	0.148	
15297	0.125	0.196	
3907	0.085	0.292	
15295	0.083	0.297	
13847	0.056	0.397	
-805	0.042	0.461	
13653	0.032	0.527	

Manuscript Information:

Patterns of Nucleotide and Haplotype Diversity at ICAM-1 Across Global Human Populations with Varying Levels of Malaria Exposure

Submitted to: Human Genetics

Felicia Gomez, Gil Tomas, Wen-Ya Ko , Alessia Ranciaro, Alain Froment, Muntaser Ibrahim, Godfrey Lema, Thomas B. Nyambo, Sabah A. Omar, Charles Wambebe, Jibril B. Hirbo, Jorge Rocha, and Sarah A. Tishkoff*,

1 Department of Genetics and Biology, School of Medicine and School of Arts and Sciences,
University of Pennsylvania

*Correspondence: tishkoff@mail.med.upenn.edu