

Supplementary Table S1. Relationship between *ADAM8* mRNA (continuous log ratio levels) and disease-free survival (top table) and overall survival (bottom table) in 295 breast cancer patients. Data was analysed using a multivariate Cox model. CI; Confidence Interval.

	Wald test <i>P</i>-value	Relative Risk	95% CI
Number of positive nodes	0.238	1.057	0.964 – 1.159
ER Category ¹	0.901	0.970	0.600 – 1.568
Size (≤ 2cm, > 2 cm)	0.044	1.537	1.011 – 2.336
Tumour Grade	0.004	1.575	1.152 – 2.154
HER-2 Status ²	0.248	1.333	0.879 – 2.171
ADAM-8	0.001	4.470	1.877 – 10.644

	Wald test <i>P</i>-value	Relative Risk	95% CI
ER Category ¹	0.055	0.607	0.365 – 1.011
Tumour size (≤, > 2cm)	0.069	1.558	0.966 – 2.513
Grade Category	0.002	1.860	1.259 – 2.748
Age (yrs)	0.047	0.962	0.926 – 0.999
HER-2 Status ²	0.132	1.498	0.886 – 2.535
ADAM-8	0.052	2.707	0.992 – 7.384

¹ Estrogen Receptor alpha expression, as measured by microarray (van de Vijver et al, 2002)

² HER2 status, as determined by microarray (van de Vijver et al, 2002)