
SUPPORTING INFORMATION

Identification of the first small molecule ligand of the neuronal receptor sortilin; and structure determination of the receptor-ligand complex

Authors

Jacob Lauwing Andersen^{a*}, Tenna Juul Schrøder^b, Søren Christensen^b, Dorthe Strandbygård^a, Lone Tjener Pallesen^c, Maria Marta García-Alai^a, Samsa Lindberg^b, Morten Langgård^b, Jørgen Calí Eskildsen^b, Laurent David^b, Lena Tagmose^b, Klaus Bæk Simonsen^b, Philip James Maltas^b, Lars Christian Biilmann Rønn^b, Inge E. M. de Jong^b, Ibrahim John Malik^b, Jan Egebjerg^b, Jens-Jacob Karlsson^b, Srinivas Uppalanchi^d, Durga Rao Sakumudi^d, Pradheep Eradi^d, Steven P. Watson^{b*} and Søren Thirup^{a*}

^aThe Lundbeck Foundation Research Centre MIND, Department of Molecular Biology and Genetics, Aarhus University, Gustav Wieds Vej 10C, Aarhus C, 8000, Denmark

^bNeuroscience Drug Discovery, H. Lundbeck A/S, Ottiliavej 9, Valby, 2500, Denmark

^cThe Lundbeck Foundation Research Centre MIND, Department of Biomedicine, Aarhus University, Ole Worms Allé 3, Aarhus C, 8000, Denmark

^dMedicinal Chemistry, GVK BioScience, Plot No. 28 A, IDA Nacharam, Hyderabad, 500076, India

Correspondence email: jla@mb.au.dk; stwa@Lundbeck.com; sth@mb.au.dk




Figure S1: Overview of crystal packing. Differences in crystal packing between the sortilin-AF40431 complex (left figures with the β -propeller in grey and 10CC-domain in green) and the sortilin-neurotensin complex (right figures with the β -propeller in orange and 10CC-domain in green). **A**) Overview of the crystal packing with the symmetry related molecules (within a distance of 5 Å) in lighter colours. **B**) Side view displaying the off-centre packing in the sortilin-AF40431 complex crystal form. **C**) Top view displaying the off-centre packing in the sortilin-AF40431 complex crystal form.