

CONSORT 2010 checklist of information to include when reporting a randomised trial*

| Section/Topic | Item No | Checklist item | Reported on page No | |
|--------------------------|------------|---|---|--|
| Title and abstract | | | _ | |
| | 1a | Identification as a randomised trial in the title | Title | |
| | 1b | Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts | s) Abstract | |
| ntroduction | | | | |
| Background and | 2a | Scientific background and explanation of rationale | Introduction | |
| bjectives | 2b | Specific objectives or hypotheses | Introduction | |
| Methods | | | M&M | |
| rial design | 3a | Description of trial design (such as parallel, factorial) including allocation ratio | | |
| | 3b | Important changes to methods after trial commencement (such as eligibility criteria), with reasons | none | |
| Participants | 4a | Eligibility criteria for participants | N 4 0 N 4 | |
| | 4b | Settings and locations where the data were collected | M&M | |
| nterventions | 5 | The interventions for each group with sufficient details to allow replication, including how and when they we actually administered | re M&M | |
| Outcomes | 6a | Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed | Table 1 | |
| | 6b | Any changes to trial outcomes after the trial commenced, with reasons | none | |
| Sample size | 7a | How sample size was determined | M&M, stat. analy | |
| | 7b | When applicable, explanation of any interim analyses and stopping guidelines | none | |
| Randomisation: | | N.S.N. | M, meth. against bias | |
| Sequence | 8a | Method used to generate the random allocation sequence | wi, metri. against bias | |
| generation | 8b | Type of randomisation; details of any restriction (such as blocking and block size) | M ma th against his s | |
| Allocation | 9 | Mechanism used to implement the random allocation sequence (such as sequentially numbered containers | s), | |
| concealment mechanism | | wick | M, meth. against bias, bered containers | |
| Implementation | 10 | Who generated the random allocation sequence, who enrolled participants, and who assigned participants | d participants to | |
| Blinding | 11a | If done, who was blinded after assignment to interventions (for example, participants, care providers, those | M, meth. against bias | |

CONSORT 2010 checklist Page 1

| | | assessing outcomes) and how | |
|---------------------|-----|--|-------------------------|
| | 11b | If relevant, description of the similarity of interventions | |
| Statistical methods | 12a | Statistical methods used to compare groups for primary and secondary outcomes M&I | Л, stat. analysis |
| | 12b | Methods for additional analyses, such as subgroup analyses and adjusted analyses M&I | M, stat. analysis |
| Results | | | |
| Participant flow (a | 13a | For each group, the numbers of participants who were randomly assigned, received intended treatment, an | |
| diagram is strongly | | were analysed for the primary outcome | Fig. 1, flow-chart |
| recommended) | 13b | For each group, losses and exclusions after randomisation, together with reasons | Fig 1 and results |
| Recruitment | 14a | Dates defining the periods of recruitment and follow-up | M&M and Results |
| | 14b | Why the trial ended or was stopped | |
| Baseline data | 15 | A table showing baseline demographic and clinical characteristics for each group | Table 2 |
| Numbers analysed | 16 | For each group, number of participants (denominator) included in each analysis and whether the analysis v | as /as |
| | | by original assigned groups | Fig. 1 |
| Outcomes and | 17a | For each primary and secondary outcome, results for each group, and the estimated effect size and its | aaa Tablaa |
| estimation | | precision (such as 95% confidence interval) | see Tables |
| | 17b | For binary outcomes, presentation of both absolute and relative effect sizes is recommended | see Tables |
| Ancillary analyses | 18 | Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishir pre-specified from exploratory | see Results |
| Harms | 19 | All important harms or unintended effects in each group (for specific guidance see CONSORT for harms) | |
| Discussion | | | |
| Limitations | 20 | Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses | Discussion, Limitations |
| Generalisability | 21 | Generalisability (external validity, applicability) of the trial findings | Discussion |
| Interpretation | 22 | Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence | e <u>Discussion</u> |
| Other information | | | |
| Registration | 23 | Registration number and name of trial registry | Abstract |
| Protocol | 24 | Where the full trial protocol can be accessed, if available | Abstract |
| Funding | 25 | Sources of funding and other support (such as supply of drugs), role of funders nor | e support or funding |

^{*}We strongly recommend reading this statement in conjunction with the CONSORT 2010 Explanation and Elaboration for important clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org.

CONSORT 2010 checklist Page 2