

Table S1. Multivariable adjusted hazard ratios (95% confidence intervals)* for proportional hazards model for other outcomes.

Risk factors	Category	eGFR<15	eGFR halving and <15	eGFR drop 20	eGFR halving or drop 20	eGFR drop 25% and change CKD stage
Age	45-64	0.70 (0.57, 0.87)	0.65 (0.53, 0.81)	0.92 (0.76, 1.11)	0.87 (0.72, 1.04)	1.04 (0.89, 1.23)
	65+	0.62 (0.48, 0.79)	0.55 (0.42, 0.70)	0.78 (0.63, 0.98)	0.74 (0.60, 0.93)	1.09 (0.90, 1.32)
Sex	Female	0.97 (0.83, 1.13)	0.90 (0.77, 1.05)	0.96 (0.84, 1.10)	0.96 (0.84, 1.09)	1.07 (0.96, 1.20)
Race	Black	1.57 (1.33, 1.85)	1.61 (1.36, 1.91)	1.33 (1.15, 1.54)	1.36 (1.18, 1.57)	1.21 (1.07, 1.36)
	Other	1.69 (1.32, 2.16)	1.66 (1.29, 2.14)	1.49 (1.20, 1.85)	1.44 (1.16, 1.78)	1.26 (1.06, 1.50)
Diabetes	Yes	1.27 (1.07, 1.49)	1.38 (1.17, 1.64)	1.31 (1.14, 1.51)	1.35 (1.18, 1.56)	1.25 (1.11, 1.41)
eGFR (ml/min/1.73m ²)	45-<60	3.52 (1.75, 7.07)	3.57 (1.78, 7.17)	0.81 (0.64, 1.02)	0.81 (0.64, 1.03)	1.55 (1.26, 1.91)
	30-<45	10.95 (5.58, 21.51)	11.47 (5.84, 22.54)	0.66 (0.52, 0.85)	0.76 (0.60, 0.97)	1.63 (1.31, 2.01)
	<30	43.61 (22.06, 86.21)	33.39 (16.86, 66.13)	1.22 (0.94, 1.58)	1.54 (1.19, 1.98)	1.59 (1.26, 2.01)
24 hour urine protein (g/day)	0.10-<0.50	1.73 (1.27, 2.37)	1.91 (1.37, 2.66)	1.65 (1.32, 2.06)	1.61 (1.29, 2.00)	1.60 (1.36, 1.88)
	0.50-<1.50	4.83 (3.57, 6.53)	4.77 (3.45, 6.60)	3.86 (3.07, 4.85)	3.84 (3.07, 4.81)	3.58 (3.01, 4.26)
	1.50+	11.19 (8.32, 15.06)	11.64 (8.48, 15.99)	9.36 (7.47, 11.73)	9.09 (7.29, 11.32)	7.32 (6.14, 8.71)

* All models were adjusted for ABI, uric acid, history of CVD, BMI, hypertension, hemoglobin, education and smoking status at baseline.