

Nitric Oxide Inhibits Hetero-adhesion of Cancer Cells to Endothelial Cells: Restraining Circulating Tumor Cells from Initiating Metastatic Cascade

Yusheng Lu¹, Ting Yu¹, Haiyan Liang¹, Jichuang Wang¹, Jingjing Xie¹, Jingwei Shao^{1*},
Yu Gao¹, Suhong Yu¹, Shuming Chen², Lie Wang², Lee Jia^{1*}

¹ Cancer Metastasis Alert and Prevention Center, College of Chemistry and Chemical Engineering, Fuzhou University, Fuzhou 350002, China.

² Surgery Department, Fuzhou General Hospital, Fuzhou, 350025, China.

*Correspondence should be addressed to Lee Jia, CMAPC, Fuzhou University, 523 Industry Road, Science Building, 3FL. Fuzhou, Fujian 350002, China. Phone: 086-591-8357-6912. Email: pharmlink@gmail.com or shaojingwei@fzu.edu.cn

Supplementary Table S1. Percent changes in E-selectin, ICAM-1 and VCAM-1 expression by HUVECs stimulated by various cytokines in comparison with isotype controls.

	Untreated	ECGS (100 µg/mL)	Fibronectin (100 ng/mL)	IL-6 (10 ng/mL)	IFN-γ (10 ng/mL)	IL-1β (1 ng/mL)	TNF-β (10 ng/mL)	TNF-α (10 ng/mL)
CD62E	1.3 ± 0.3%	0.7 ± 0.3%	1.3 ± 0.4%	2.9 ± 0.3%	1.1 ± 0.1%	64.2 ± 3.3%	29.0 ± 2.0%	75.2 ± 3.6%
CD54	66.0 ± 1.8%	80.6 ± 1.4%	78.1 ± 1.4%	80.7 ± 1.0%	93.5 ± 1.0%	99.4 ± 0.2%	99.5 ± 0.2%	99.5 ± 0.2%
CD106	15.3 ± 1.2%	15.4 ± 1.2%	35.2 ± 1.5%	33.4 ± 1.8%	37.0 ± 1.6%	92.8 ± 1.8%	74.9 ± 1.2%	93.7 ± 0.9%

Values are mean ± SEM (n=3).