

Table S1: *S. pombe* strains used in this study

Strain	Genotype	Source
JP3	<i>h</i> ⁻ 972	Lab stock
JP57	<i>h</i> ⁻ <i>plo1.S402A</i>	Lab stock
JP350	<i>h</i> ⁺	Lab stock
JP567	<i>h</i> ⁻ <i>cdc2-1w</i>	(Thuriaux et al., 1978)
JP568	<i>h</i> ⁻ <i>cdc2-3w</i>	(Fantes, 1981)
JP1377	<i>sin1::kanMx6</i>	<i>sin1::kanMx6</i> from (Ikeda et al., 2008)
JP1491	<i>tor1::ura4⁺ ura4-D18</i>	<i>tor1::ura4⁺</i> from (Matsuo et al., 2003)
JP2337	<i>cdr2::ura4⁺ ura4-D18</i>	<i>cdr2::ura4⁺</i> from (Bimbo et al., 2005)
JP1662	<i>h</i> ⁺ <i>nmt81-GFPwee1</i>	(Moseley et al., 2009)
JP1670	<i>h</i> ⁺ <i>cdc25:12myc-ura4⁺ ura4-D18</i>	<i>cdc25:12myc.ura4⁺</i> from (Lopez-Girona et al., 1999)
JP1669	<i>h</i> ⁺ <i>tor2-G2040D</i>	This study
JP1731	<i>h</i> ⁻ <i>maf1pk-kanMx6</i>	Lab stock
JP1763	<i>h</i> ⁻ <i>maf1pk-kanMx6 tor2-G2040D</i>	This study
JP2081	<i>tor2-G2040D nmt81-GFPwee1</i>	This study
JP2089	<i>h</i> ⁺ <i>tor2::ura4⁺ ura4-d18 leu1-32</i> <i>pREP42NAT-tor2</i>	This study <i>tor2::ura4⁺</i> from (Weisman et al., 2007)

JP2090	<i>h⁺ tor2::ura4+ ura4-d18 leu1-32 pREP42NAT-tor2-G2040D</i>	This study
JP2177	<i>cdr2::ura4⁺ nmt81-GFPwee1 ura4-d18</i>	This study
JP2377	<i>plo1.S402A nmt81-GFPwee1</i>	This study
JP2363	<i>h⁺ cdr2::ura4⁺ plo1.S402A ura4.D18</i>	This study
JP2405	<i>cdr2::ura4⁺ plo1.S402A nmt81-GFPwee1</i>	This study
JP2411	<i>h⁻ cdc2::ura4⁺ leu1.32 ura4.D18 mns21L-cdc2.F15 integrated</i>	Paul Nurse D832

References

- Bimbo, A., Y. Jia, S.L. Poh, R.K. Karuturi, N. den Elzen, X. Peng, L. Zheng, M. O'Connell, E.T. Liu, M.K. Balasubramanian, and J. Liu. 2005. Systematic deletion analysis of fission yeast protein kinases. *Eukaryotic cell.* 4:799-813.
- Fantes, P.A. 1981. ISOLATION OF CELL-SIZE MUTANTS OF A FISSION YEAST BY A NEW SELECTIVE METHOD - CHARACTERIZATION OF MUTANTS AND IMPLICATIONS FOR DIVISION CONTROL MECHANISMS. *Journal of Bacteriology.* 146:746-754.
- Ikeda, K., S. Morigasaki, H. Tatebe, F. Tamanoi, and K. Shiozaki. 2008. Fission yeast TOR complex 2 activates the AGC-family Gad8 kinase essential for stress resistance and cell cycle control. *Cell Cycle.* 7:358-364.
- Lopez-Girona, A., B. Furnari, O. Mondesert, and P. Russell. 1999. Nuclear localization of Cdc25 is regulated by DNA damage and a 14-3-3 protein. *Nature.* 397:172-175.
- Matsuo, T., Y. Kubo, Y. Watanabe, and M. Yamamoto. 2003. Schizosaccharomyces pombe AGC family kinase Gad8p forms a conserved signaling module with TOR and PDK1-like kinases. *Embo Journal.* 22:3073-3083.
- Moseley, J.B., A. Mayeux, A. Paoletti, and P. Nurse. 2009. A spatial gradient coordinates cell size and mitotic entry in fission yeast. *Nature.* 459:857-U858.
- Thuriaux, P., P. Nurse, and B. Carter. 1978. MUTANTS ALTERED IN CONTROL COORDINATING CELL-DIVISION WITH CELL-GROWTH IN FISSION YEAST

SCHIZOSACCHAROMYCES-POMBE. *Molecular & General Genetics*. 161:215-220.

Weisman, R., I. Roitburg, M. Schonbrun, R. Harari, and M. Kupiec. 2007. Opposite effects of Tor1 and Tor2 on nitrogen starvation responses in fission yeast. *Genetics*. 175:1153-1162.

|