

Supplementary table 1. Antifungal susceptibility (in mg/L) of rare species according EUCAST protocol (incidence <1%)

Species (N)	Value	MIC (mg/L)								
		AmB	Flucytosine	FLC	ITZ	VOR	POS	CAS	MIC	AND
<i>C. orthopsilosis</i> (n=7)	GM	0.06	0.12	0.5	0.02	0.018	0.015	0.906	0.305	0.37
	Mode	0.06	0.12	0.5	0.015	0.015	0.015	1	0.25	0.25
	Range	0.03-0.12	0.12 – 0.12	0.5 – 0.5	0.015 – 0.03	0.015 – 0.015	0.015 – 0.015	0.5 – 1	0.25 – 0.5	0.25 – 0.5
<i>C. neoformans</i> ^a (n=5)	GM	0.144	11.3	16	0.043	0.1	0.51	16	16	16
	Mode	0.12	4	8	0.015	0.015	0.03	32	32	32
	Range	0.12 -0.25	4 – 32	8 – >64	0.015 – 0.5	0.015 – 4	0.015 – 0.5	16 – >16	>16 – >16	>16 – >16
<i>C. kefir</i> (n=4)	GM	0.35	0.17	0.21	0.019	0.015	0.021	0.247	0.03	0.03
	Mode	0.25	0.12	0.12	0.015	0.015	0.015	0.25	0.03	0.03
	Range	0.12 -2	0.12 – 0.5	0.12 – 0.5	0.015 – 0.03	0.015 – 0.015	0.015 – 0.03	0.12 – 0.5	0.03 – 0.03	0.03 – 0.03
<i>C. dubliniensis</i> (N=4)	GM	0.036	0.12	0.17	0.021	0.015	0.015	0.35	0.03	0.03
	Mode	0.03	0.12	0.12	0.015	0.015	0.015	0.25	0.03	0.03
	Range	0.03 - 0.06	0.12 – 0.12	0.12 – 0.25	0.12 – 0.25	0.015 – 0.015	0.015 – 0.015	0.25 – 0.5	0.03 - 0.03	0.03 – 0.03
<i>C. lypolytica</i> ^a (n=3)	GM	0.32	64	12.7	0.4	0.06	1	6.3	3.2	2
	Range	0.25 - 0.5	32 – >64	2 - >64	0.12 – 2	0.015 – 0.25	0.25 – >8	2 - >16	0.5 - >16	0.5 - >16
<i>D. capitatus</i> ^a (n=3)	GM	0.31	0.15	50.8	0.63	3.17	0.8	16	8	6.35
	Range	0.12 - 0.5	0.12 - 0.25	32 - >64	0.25 - 2	1 - >8	0.5 - 2	16 - >16	0.5 - >16	2 - >16
<i>T. asahii</i> ^a (n=3)	GM	2	11.3	2	0.17	0.25	0.25	16	16	11.3
	Rango	2 - 2	8 - 16	1 - 4	0.12 - 0.25	0.25 - 0.25	0.25 - 0.25	16 - >16	>16 ->16	4 - >16
<i>C. metapsilosis</i> N=2	MIC (1)	0.06	0.12	1	0.015	0.03	0.015	1	0.25	0.12
	MIC (2)	0.03	0.12	1	0.03	0.03	0.015	0.5	0.25	0.12
<i>P. anomala</i> N=2	MIC (1)	0.03	0.12	2	0.06	0.06	0.06	0.25	0.03	0.03
	MIC (2) ^a	0.12	0.12	4	0.12	0.12	0.06	0.25	0.03	0.03
<i>R. mucilaginosa</i> ^a	MIC (1)	0.25	1	>64	1	>8	2	16	>16	>16
	MIC (2)	0.25	0.25	>64	4	8	4	16	>16	>16
<i>C. intermedia</i> *	MIC	0.03	64	0.25	0.015	0.015	0.015	0.25	0.06	0.03
<i>K. ohmeri</i> *	MIC	0.03	0.12	8	0.06	0.06	0.015	>16	0.12	1
<i>L. elongisporus</i> *	MIC	0.03	0.5	0.25	0.06	0.015	0.03	0.25	0.03	0.03
<i>C. nivariensis</i> *	MIC	0.25	1	2	2	4	0.06	0.25	0.03	0.03
<i>P. fabianii</i> *	MIC	0.06	0.12	1	0.03	0.015	0.06	0.25	0.03	0.03

<i>M. pulcherrima*</i>	MIC	0.03	0.12	0.25	0.015	0.015	0.015	0.5	0.06	0.03
------------------------	-----	------	------	------	-------	-------	-------	-----	------	------

640

641 MIC values were obtained after 24 h of incubation according to the EUCAST protocol, except in those isolates that presented low
 642 growth or non-fermentative yeast. In these last cases (^a), MIC values were obtained using the EUCAST recommendations for non-
 643 fermentative yeasts (30°C incubation, shaking and 48 h growth). In the cases where n=2, both MIC values are shown. * n=1.

644

Supplementary table 2. Antifungal susceptibility (in mg/L) of rare species according CLSI protocol (incidence <1%)

Species (N)	Value	MIC (mg/L)						
		AmB	FLC	VOR	POS	CAS	MIC	AND
<i>C. orthopsilosis</i> (n=7)	GM	0.12	0.41	0.01	0.016	0.27	0.33	0.74
	Mode	0.12	1	0.003 ^c	0.015	0.25	0.25	1
	Range	0.12 - 0.12	0.062 - 1	0.003 – 0.3	0.003 – 0.06	0.12 – 0.5	0.12 – 1	0.5 – 1
<i>C. neoformans</i> ^a (n=5)	GM	0.17	1.68	0.018	0.009	>2	>2	>2
	Mode	0.06	1	0.03	0.003	>2	>2	>2
	Range	0.06 - 0.5	1 - 4	0.007 – 0.03	0.003 – 0.06	>2 - >2	>2 - >2	>2 - >2
<i>C. kefyr</i> (n=4)	GM	0.71	0.5	0.008	0.04	0.09	0.05	0.05
	Mode	0.5	0.5	0.007	0.03 ^c	0.12	0.06	0.03
	Range	0.25 - >2	0.25 - 1	0.007 – 0.015	0.03 – 0.06	0.03 – 0.12	0.03 – 0.06	0.03 – 0.12
<i>C. dubliniensis</i> (n=4)	GM	0.06	0.42	0.004	0.06	0.15	0.009	0.018
	Mode	0.06	0.5	0.007	0.015	0.12	NA	0.03
	Range	0.015 – 0.25	0.12 - 1	0.0017 – 0.007	0.0017 – 0.015	0.12 – 0.25	0.0017 – 0.03	0.007 – 0.03
<i>C. lypolytica</i> ^b (n=3)	GM	0.25	2.5	0.09	0.12	0.06	0.5	0.027
	Range	0.12 – 0.5	0.25 - >64	0.015 – >2	0.015 – >2	0.007 – 0.5	0.5 – 0.5	0.003 – 0.25
<i>D. capitatus</i>	GM	0.5	64	1.26	0.4	>2	2	>2
	Range	0.25 - 1	32 - >64	0.5 – >2	0.25 – 0.5	>2 - >2	2 - 2	2 - >2
<i>T. asahii</i>	GM	1	0.5	0.015	0.06	>2	2	>2
	Range	1 – 1	0.5 – 0.5	0.015 – 0.015	0.06 – 0.06	>2 - >2	2 - 2	>2 - >2
<i>C. metapsilosis</i> (n=2)	MIC (1)	0.12	0.12	0.003	0.003	0.25	0.25	0.5
	MIC (2)	0.12	1	0.015	0.03	0.03	0.03	0.12
<i>P. anomala</i>	MIC (1)	0.5 ^b	0.12	0.015	0.06	0.12	0.03	0.007
	MIC (2)	0.12	1	0.007	0.015	0.12	0.003	0.0017
<i>R. mucilaginosa</i>	MIC (1) ^b	0.25	32	0.5	0.06	>2	>2	>2
	MIC (2) ^a	0.5	>64	1	1	>2	>2	>2
<i>C. intermedia</i> *	MIC	0.06	1	0.003	0.003	0.12	0.06	0.03
<i>K. ohmeri</i> *	MIC	0.06	1	0.03	0.06	0.5	0.06	0.007
<i>L. elongisporus</i> *	MIC	0.03	0.12	0.0017	0.007	0.06	0.015	0.015
<i>C. nivariensis</i> *	MIC	0.5	4	0.03	0.12	0.25	0.015	0.03
<i>P. fabianii</i> *	MIC	0.12	2	0.03	0.06	0.12	0.007	0.007

<i>M. pulcherrima</i> * ^b	MIC	0.5	0.12	0.0017	0.0017	0.015	0.007	0.007
--------------------------------------	-----	-----	------	--------	--------	-------	-------	-------

646
647 Susceptibility data was obtained after 24 hours of incubation. In some cases, since the isolates did not show enough growth at 24
648 hours, the MICs were obtained after 48 (^b) or 72 (^a) hours of incubation. ^c When multiple modes available, the smallest value is
649 shown. **NA**, Mode not available due to the low number of isolates. When n=2, both MICs are shown. *, n=1