

Supporting Information

Figure S1. The k^3 -weighted polarized Mn EXAFS spectra. Inset shows the crystal setting with respect to the incident X-ray \mathbf{e} -vector and the two rotation axes, ϕ_{crystal} and χ_{crystal} . The solid line (red), X-ray \mathbf{e} -vector parallel to the crystal b -axis ($\phi_{\text{crystal}} = 90^\circ$, $\chi_{\text{crystal}} = 0^\circ$); dotted line (green), X-ray \mathbf{e} -vector parallel to the c -axis ($\phi_{\text{crystal}} = 0^\circ$, $\chi_{\text{crystal}} = 0^\circ$); dash-dot line (blue), X-ray \mathbf{e} -vector parallel to the a -axis ($\phi_{\text{crystal}} = 0^\circ$, $\chi_{\text{crystal}} = 90^\circ$).




Figure 2S. Fourier transforms of the k^3 -weighted polarized Mn EXAFS spectra. The solid line (red), X-ray \mathbf{e} -vector parallel to the crystal b -axis ($\phi_{\text{crystal}} = 90^\circ$, $\chi_{\text{crystal}} = 0^\circ$); dotted line (green), X-ray \mathbf{e} -vector parallel to the c -axis ($\phi_{\text{crystal}} = 0^\circ$, $\chi_{\text{crystal}} = 0^\circ$); dash-dot line (blue), X-ray \mathbf{e} -vector parallel to the a -axis ($\phi_{\text{crystal}} = 0^\circ$, $\chi_{\text{crystal}} = 90^\circ$). At $(\phi_{\text{crystal}}, \chi_{\text{crystal}}) = (0^\circ, 0^\circ)$, the Mn-Mn vector (at $\sim 2.7 \text{ \AA}$, the second Fourier peak) amplitude was maximal, while it was close to the noise level at $(\phi_{\text{crystal}}, \chi_{\text{crystal}}) = (90^\circ, 0^\circ)$ orientation. The Mn-ligand vector amplitude at $\sim 2 \text{ \AA}$ was also minimal at the $(\phi_{\text{crystal}}, \chi_{\text{crystal}}) = (90^\circ, 0^\circ)$ orientation.

