


**Risk Factors for Development of Cholestatic Drug-induced Liver Injury:  
Inhibition of Hepatic Basolateral Bile Acid Transporters MRP3 and  
MRP4**

Kathleen Köck, Brian C. Ferslew, Ida Netterberg, Kyunghee Yang, Thomas J. Urban, Peter W. Swaan, Paul W. Stewart, and Kim L. R. Brouwer

**Drug Metabolism and Disposition**



**Supplemental Figure 1.** Schematic of the high-throughput assay to study MRP3 and MRP4 inhibition.

**Supplementary Table 1. Human protein binding, C<sub>max</sub>, unbound C<sub>max</sub> (C<sub>max,u</sub>), maximum daily dose (MDD) in man, main route of excretion, extent of liver metabolism and involvement of phase II metabolism for the 88 investigated non-cholestatic and cholestatic compounds.**

Drug	Protein binding (%)	C <sub>max</sub> (µg/ ml)	C <sub>max,u</sub> (µM) <sup>a</sup>	MDD in man (mg/ day) <sup>b</sup>	Standard dose (mg/day) <sup>c</sup>	Main Route of Excretion <sup>d</sup>	Liver Metabolism <sup>e</sup>	Phase II Metabolism
<b>Non-cholestatic BSEP non-inhibitors</b>								
5-Fluorouracil	8.0 <sup>2</sup>	17.7 <sup>1</sup>	125	4420 iv <sup>1</sup>	2380	Urinary <sup>19</sup>	Extensive <sup>20</sup>	No
Alprenolol	80.0 <sup>2</sup>	0.14 <sup>3</sup>	0.11	800 <sup>1</sup>	300	Urinary	Extensive	No
Antipyrine	10.0	25 <sup>3</sup>	120	800	800	N/A	N/A	N/A
Aspirin	80.0 <sup>1</sup>	200 <sup>3</sup>	222	4000 <sup>1</sup>	2037.5	Urinary	Extensive	No
Caffeine	36.0 <sup>1</sup>	17.3 <sup>1</sup>	57	1200 <sup>1</sup>	632.5	Urinary	Extensive	No
Chlorpheniramine	70.0 <sup>2</sup>	0.08 <sup>2</sup>	0.083	24 <sup>1</sup>	24	Urinary	Extensive	No
Clopamide	46.0 <sup>1</sup>	0.18 <sup>21</sup>	0.27	40 <sup>1</sup>	22.5	Urinary	N/A	No
Dexamethasone	77.0 <sup>18</sup>	0.10 <sup>22</sup>	0.075	24 <sup>1</sup>	4.88	Urinary	Extensive <sup>23</sup>	Yes <sup>24</sup>
Diphenhydramine	76.0 <sup>1</sup>	0.23	0.22	300 <sup>1</sup>	200	Urinary	Extensive	Yes
Doxorubicin	74.0 <sup>1</sup>	0.95	0.45	75/ m <sup>2</sup> /3 week iv <sup>1</sup>	81	Biliary	Extensive	No

## DMD #54304

Etoposide	96.0 <sup>2</sup>	4.7	0.32	2 g/m <sup>2</sup> / 24 h iv <sup>1</sup>	85	Urinary	Extensive	No
Fluorescein	85.0 <sup>1</sup>	35 <sup>1</sup>	15.8	500 iv <sup>1</sup>	500	Urinary	Extensive	Yes
Metoclopramide	30.0 <sup>1</sup>	0.1 <sup>1</sup>	0.23	60 <sup>6</sup>	370	Urinary	Non-significant	Yes
Nadolol	28.0 <sup>1</sup>	0.25 <sup>3</sup>	0.58	320 <sup>1</sup>	45	Urinary	Non-significant	No
Naloxone	45.9 <sup>25</sup>	0.03 <sup>3</sup>	0.050	3 <sup>1</sup>	1.2	Urinary	Extensive	Yes
Phenformin	12.0 <sup>1</sup>	0.15 <sup>1</sup>	0.65	2550 <sup>1</sup>	81.25	Urinary	Non-significant	No
Probenecid	80.0 <sup>1</sup>	150 <sup>3</sup>	100	2000 <sup>1</sup>	1125	Urinary	Extensive	Yes
Quinidine	80.0 <sup>1</sup>	8 <sup>1</sup>	4.9	1600 <sup>6</sup>	1200	Urinary	Extensive	No
Tacrine	55.0 <sup>1</sup>	0.016 <sup>5</sup>	0.036	160 <sup>1</sup>	40	Urinary <sup>26</sup>	Extensive	No
Terbutaline	20 <sup>27</sup>	0.03 <sup>3</sup>	0.11	15 <sup>1</sup>	11.25	Urinary	Non-significant	Yes
Theophylline	40.0 <sup>1</sup>	15 <sup>1</sup>	50	900 <sup>1</sup>	450	Urinary	Extensive	No
Timolol	10.0 <sup>1</sup>	0.05 <sup>3</sup>	0.14	60 <sup>1</sup>	35	Urinary	Extensive	No
Triamterene	55.0 <sup>1</sup>	0.1 <sup>3</sup>	0.18	300 <sup>1</sup>	75	Urinary	Extensive	Yes
Vinblastine	98.0 <sup>1</sup>	0.001 <sup>28</sup>	0.00002	11.1/m <sup>2</sup> iv <sup>1</sup>	12.6	Biliary	Extensive	No

## Cholestatic BSEP non-inhibitors

Bezafibrate	95.0 <sup>1</sup>	14.3 <sup>1</sup>	2.0	600 <sup>1</sup>	500	Urinary	N/A	Yes
Carbamazepine	74.0 <sup>2</sup>	12 <sup>3</sup>	13	1600 <sup>1</sup>	1000	Urinary	Extensive	No

## DMD #54304

Chloramphenicol	50.0 <sup>1</sup>	25 <sup>3</sup>	39	7000 <sup>1</sup>	5250	Urinary	Extensive	No
Chlorpromazine	90.0 <sup>1</sup>	0.3 <sup>1</sup>	0.094	2000 <sup>1</sup>	400	Urinary (P.I.)	Extensive	No
Chlorpropamide	60.0 <sup>1</sup>	150 <sup>3</sup>	217	750 <sup>1</sup>	300	Urinary	Extensive	No
Cimetidine	13.0 <sup>1</sup>	3.9 <sup>1</sup>	13	1600 <sup>1</sup>	800	Urinary	Extensive	Yes
Desipramine	73.0 <sup>1</sup>	0.5 <sup>3</sup>	0.45	300 <sup>1</sup>	150	Urinary	Extensive	Yes
D-Penicillamine	80.0 <sup>6</sup>	5.6 <sup>3</sup>	7.5	4000 <sup>1</sup>	187.5	Urinary (P.I.)	Extensive	No
Famotidine	10.0 <sup>1</sup>	104 <sup>1</sup>	277	640 <sup>1</sup>	360	Urinary	Non-significant	Yes
Fluoxetine	94.0 <sup>2</sup>	0.477 <sup>1</sup>	0.09	80 <sup>1</sup>	50	Urinary	Extensive	No
Furosemide	91.0 <sup>1</sup>	6 <sup>3</sup>	1.6	600 <sup>1</sup>	90	Urinary	Non-significant	No
Haloperidol	90.0 <sup>1</sup>	0.02 <sup>2</sup>	0.006	15 <sup>1</sup>	1.25	Urinary	N/A	Yes
Ibuprofen	99.0 <sup>2</sup>	57 <sup>1</sup>	2.8	3200 <sup>1</sup>	1000	Urinary	Extensive	Yes
Maprotiline	88.0 <sup>1</sup>	0.25 <sup>3</sup>	0.11	225 <sup>1</sup>	112.5	Urinary	N/A	No
Metformin	0.0 <sup>2</sup>	1 <sup>3</sup>	8	2550 <sup>1</sup>	1250	Urinary	Non-significant	No
Nitrofurantoin	62.0 <sup>2</sup>	5 <sup>3</sup>	8.0	400 <sup>1</sup>	300	Urinary (P.I.)	N/A	No
Nortriptyline	86.0 <sup>1</sup>	0.2 <sup>3</sup>	0.11	100 <sup>1</sup>	87.5	Urinary	Extensive	No
Promethazine	93.0 <sup>1</sup>	0.4 <sup>3</sup>	0.10	150 <sup>1</sup>	37.5	Urinary	Extensive	No
Quinine	69.0 <sup>1</sup>	10 <sup>2</sup>	10	2600 <sup>1</sup>	1800	Urinary	Extensive	Yes

## DMD #54304

Ranitidine	15.0 <sup>2</sup>	1 <sup>3</sup>	2.7	300 <sup>1</sup>	150	Urinary	Extensive	No
Sulfasalazine	99.3 <sup>1</sup>	30 <sup>3</sup>	0.53	4000 <sup>1</sup>	2000	Urinary	N/A	No
Sulindac	93.0 <sup>1</sup>	6 <sup>3</sup>	1.2	400 <sup>1</sup>	350	Urinary	Extensive	Yes
Tamoxifen	98.0 <sup>2</sup>	0.12 <sup>2</sup>	0.006	40 <sup>1</sup>	30	Biliary	Extensive	No
Tolbutamide	80.0 <sup>1</sup>	100 <sup>3</sup>	74	3000 <sup>1</sup>	1125	Urinary	Extensive	No
Trimethoprim	37.0 <sup>2</sup>	2.5 <sup>2</sup>	5.4	1050 <sup>1</sup>	150	Urinary	Non-significant	No
Verapamil	90.0 <sup>2</sup>	0.27 <sup>2</sup>	0.06	480 <sup>1</sup>	280	Urinary	Extensive	No

## Non-cholestatic BSEP-inhibitors

Alpidem	99.1 <sup>4</sup>	0.27 <sup>4</sup>	0.006	150	150	Biliary <sup>8</sup>	Extensive <sup>8</sup>	Yes
Benzbromarone	99.0 <sup>9</sup>	3.35 <sup>10</sup>	0.079	200 <sup>1</sup>	125	Biliary <sup>11</sup>	Extensive	Yes
Buspirone	86.0 <sup>1</sup>	0.004	0.002	90 <sup>1</sup>	25	Urinary	Extensive	No
Clobetasol propionate	97.5 <sup>4</sup>	0.02 <sup>4</sup>	0.001	50 g/week topical <sup>1</sup>		Urinary	N/A	N/A
Finasteride	90.0 <sup>2</sup>	0.37 <sup>2</sup>	0.099	5 <sup>1</sup>	3	Biliary <sup>12</sup>	Extensive	No
Flupirtine	80.0 <sup>13</sup>	1.5 <sup>3</sup>	0.99	600 <sup>1</sup>	450	Urinary <sup>14</sup>	N/A	Yes <sup>15</sup>
Glafenine	96.7 <sup>4</sup>	0.7 <sup>5</sup>	0.062	1200 <sup>4</sup>	800	Urinary <sup>16</sup>	Extensive <sup>16</sup>	Yes <sup>17</sup>
Lopinavir	98.0 <sup>6</sup>	11.8 <sup>1</sup>	0.38	800 <sup>6</sup>	800	Biliary	Extensive	No
Mibepradil	99.5 <sup>1</sup>	0.3 <sup>1</sup>	0.003	100 <sup>1</sup>	50	Biliary	Extensive	No

## DMD #54304

Oxybutynin	99.0 <sup>1</sup>	0.001 <sup>1</sup>	0.00028	30 <sup>1</sup>	7.5	Biliary <sup>11</sup>	Extensive	No
Praziquantel	80.0 <sup>2</sup>	2.2 <sup>1</sup>	1.4	5250 <sup>1</sup>	5250	Urinary	Extensive	No
Primaquine	77.0 <sup>18</sup>	0.2 <sup>3</sup>	0.18	30 <sup>1</sup>	30	Biliary <sup>11</sup>	Extensive	N/A
Sorafenib	99.5 <sup>1</sup>	9.9 <sup>1</sup>	0.106	800 <sup>1</sup>	800	Biliary	Extensive	Yes
Taxol	88.0 <sup>2</sup>	1 <sup>3</sup>	0.14	250/m <sup>2</sup> /3 week iv <sup>6</sup>	217	Biliary	Extensive	No
Tolcapone	99.9 <sup>2</sup>	13 <sup>3</sup>	0.048	1200 <sup>1</sup>	450	Urinary	Extensive	Yes
Valinomycin	N/A	N/A		N/A	N/A	N/A	N/A	N/A

**Cholestatic BSEP-inhibitors**

Acitretin	99.0 <sup>1</sup>	0.42 <sup>1</sup>	0.013	75 <sup>1</sup>	37.5	Urinary/biliary	Extensive	Yes
Clozapine	95.0 <sup>2</sup>	0.8 <sup>3</sup>	0.12	900 <sup>1</sup>	375	Urinary	Extensive	No
Cyclosporin A	90.0	0.4 <sup>3</sup>	0.03	700 <sup>1</sup>	525	Biliary	Extensive	No
Dicloxacillin	88.0 <sup>1</sup>	91 <sup>2</sup>	23	2000 <sup>1</sup>	1750	Urinary	Non-significant	No
Erythromycin Estolate	95.0 <sup>4</sup>	3.08 <sup>5</sup>	0.15	1000 <sup>4</sup>	1000	Biliary	Non-significant	No
Fenofibrate	99.0 <sup>1</sup>	30 <sup>1</sup>	0.83	160 <sup>1</sup>	100	Urinary	N/A	Yes
Fluvastatin	98.0 <sup>1</sup>	0.42 <sup>1</sup>	0.02	80 <sup>1</sup>	50	Biliary	Extensive	Yes
Glyburide	99.0 <sup>1</sup>	0.2 <sup>1</sup>	0.004	20 <sup>1</sup>	10.6	Urinary/biliary	Extensive	No
Indinavir	60.0 <sup>1</sup>	0.013 <sup>1</sup>	0.008	2400 <sup>1</sup>	2100	Biliary	Extensive	Yes

## DMD #54304

Indomethacin	90.0 <sup>2</sup>	3 <sup>3</sup>	0.84	200 <sup>1</sup>	125	Urinary	Extensive	No
Nifedipine	90.0 <sup>1</sup>	0.1 <sup>3</sup>	0.029	180 <sup>6</sup>	45	Urinary	Extensive	No
Nitrendipine	97.0 <sup>1</sup>	0.05 <sup>3</sup>	0.004	40 <sup>1</sup>	25	Urinary	Extensive	Yes
19-Norethindrone	61.0 <sup>1</sup>	0.01 <sup>5</sup>	0.013	5 <sup>6</sup>	2.7	Urinary	Extensive	Yes
Omeprazole	95.0 <sup>2</sup>	4 <sup>3</sup>	0.58	360 <sup>1</sup>	30	Urinary	Extensive	No
Pioglitazone	99.0 <sup>1</sup>	1.5 <sup>2</sup>	0.042	45 <sup>1</sup>	22.5	Biliary (P.I.)	Extensive	No
Rifampicin	60.0 <sup>1</sup>	10 <sup>3</sup>	4.9	600 <sup>1</sup>	450	Biliary	Extensive	No
Rifamycin SV	80.0 <sup>4</sup>	11 <sup>4</sup>	3.2	600 <sup>4</sup>	450	Biliary	Extensive	No
Ritonavir	98.0 <sup>1</sup>	11.2 <sup>1</sup>	0.31	1200 <sup>1</sup>	600	Biliary	Extensive	No
Rosiglitazone	99.8 <sup>1</sup>	0.275 <sup>2</sup>	0.0015	8 <sup>1</sup>	4	Urinary	Extensive	Yes
Simvastatin	94.0 <sup>2</sup>	0.06 <sup>7</sup>	0.008	80 <sup>1</sup>	22.5	Biliary	Extensive	No
Ticlopidine	98.0 <sup>2</sup>	2.1 <sup>5</sup>	0.14	500 <sup>1</sup>	500	Urinary	Extensive	No
Troglitazone	99.0 <sup>1</sup>	2.8 <sup>5</sup>	0.064	600 <sup>1</sup>	400	Biliary	Extensive	Yes

<sup>a</sup> Calculated from C<sub>max</sub> and protein binding<sup>b</sup> Administered orally if not stated otherwise.<sup>c</sup> If a dose range was recommended (e.g. for different indications), the average of minimum and maximum recommended daily dose was calculated. If dosing was given as mg/kg or mg/m<sup>2</sup>, the maximum and average dose was calculated based on an adult with 70 kg and a body surface area of 1.7.<sup>d</sup> Compounds were categorized as urinary vs. biliary if >50% of the reported elimination occurred by a given pathway.

<sup>e</sup> Compounds with ≥50% of hepatic metabolism were categorized as “extensive”; compounds < 50% were categorized into the “non-significant” group.

N/A, Not Available, P.I., packaging insert

- <sup>1</sup> Micromedex®. Version 2.0 (accessed January 2012). In. Greenwood Village, Colo: Thomson Reuters (Healthcare).
- <sup>2</sup> Hardman JG, Limbird LE. Goodman and Gilman's the pharmacological basis of therapeutics. , 2001.
- <sup>3</sup> Regenthal R, Krueger M, Koeppel C, Preiss R. Drug levels: therapeutic and toxic serum/plasma concentrations of common drugs. *J Clin Monit Comput* 1999;15:529-544.
- <sup>4</sup> Dawson S, Stahl S, Paul N, Barber J, Kenna JG. In vitro inhibition of the bile salt export pump correlates with risk of cholestatic drug-induced liver injury in humans. *Drug Metab Dispos* 2012;40:130-138.
- <sup>5</sup> Xu JJ, Henstock PV, Dunn MC, Smith AR, Chabot JR, de Graaf D. Cellular imaging predictions of clinical drug-induced liver injury. *Toxicol Sci* 2008;105:97-105.
- <sup>6</sup> Lexi-Comp. <http://www.lexi.com/> (accessed January 2012). In. Hudson, Ohio: Lexi-Comp Inc.
- <sup>7</sup> Kyrklund C, Backman JT, Kivistö KT, Neuvonen M, Laitila J, Neuvonen PJ. Rifampin greatly reduces plasma simvastatin and simvastatin acid concentrations. *Clin Pharmacol Ther* 2000;68:592-597.
- <sup>8</sup> Durand A, Thenot JP, Bianchetti G, Morselli PL. Comparative pharmacokinetic profile of two imidazopyridine drugs: zolpidem and alpidem. *Drug Metab Rev* 1992;24:239-266.
- <sup>9</sup> Lee MH, Graham GG, Williams KM, Day RO. A benefit-risk assessment of benz bromarone in the treatment of gout. Was its withdrawal from the market in the best interest of patients? *Drug Saf* 2008;31:643-665.
- <sup>10</sup> de Vries JX, Walter-Sack I, Ittensohn A, Weber E, Empl H, Gresser U, Zollner N. Benz bromarone hydroxylation in man: defective formation of the 6-hydroxybenz bromarone metabolite. *Clin Investig* 1993;71:947-952.

- <sup>11</sup> Morgan RE, Trauner M, van Staden CJ, Lee PH, Ramachandran B, Eschenberg M, Afshari CA, et al. Interference with bile salt export pump function is a susceptibility factor for human liver injury in drug development. *Toxicol Sci* 2010;118:485-500.
- <sup>12</sup> Lundahl A, Hedeland M, Bondesson U, Knutson L, Lennernas H. The effect of St. John's wort on the pharmacokinetics, metabolism and biliary excretion of finasteride and its metabolites in healthy men. *Eur J Pharm Sci* 2009;36:433-443.
- <sup>13</sup> Devulder J. Flupirtine in pain management: pharmacological properties and clinical use. *CNS Drugs* 2010;24:867-881.
- <sup>14</sup> Hlavica P, Niebch G. [Pharmacokinetics and biotransformation of the analgesic flupirtine in humans]. *Arzneimittelforschung* 1985;35:67-74.
- <sup>15</sup> Methling K, Reszka P, Lalk M, Vrana O, Scheuch E, Siegmund W, Terhaag B, et al. Investigation of the in vitro metabolism of the analgesic flupirtine. *Drug Metab Dispos* 2009;37:479-493.
- <sup>16</sup> Pottier J, Busigny M, Raynaud JP. Biotransformations of glafenine in the rat and in man. *Eur J Drug Metab Pharmacokinet* 1979;4:109-115.
- <sup>17</sup> Wen B, Moore DJ. Bioactivation of glafenine by human liver microsomes and peroxidases: identification of electrophilic iminoquinone species and GSH conjugates. *Drug Metab Dispos* 2011;39:1511-1521.
- <sup>18</sup> Peets EA, Staub M, Symchowicz S. Plasma binding of betamethasone-3H, dexamethasone-3H, and cortisol-14C--a comparative study. *Biochem Pharmacol* 1969;18:1655-1663.
- <sup>19</sup> Heggie GD, Sommadossi JP, Cross DS, Huster WJ, Diasio RB. Clinical pharmacokinetics of 5-fluorouracil and its metabolites in plasma, urine, and bile. *Cancer Res* 1987;47:2203-2206.
- <sup>20</sup> Diasio RB, Harris BE. Clinical pharmacology of 5-fluorouracil. *Clin Pharmacokinet* 1989;16:215-237.
- <sup>21</sup> McNeil JJ, Conway EL, Drummer OH, Howes LG, Christophidis N, Louis WJ. Clopamide: plasma concentrations and diuretic effect in humans. *Clin Pharmacol Ther* 1987;42:299-304.
- <sup>22</sup> Harahap Y, Sasongko L, Prasaja B, Indriati E, Lusthom W, Lipin. Comparative bioavailability of two dexamethasone tablet formulations in Indonesian healthy volunteers. *Arzneimittelforschung* 2009;59:191-194.
- <sup>23</sup> Seutter E. Metabolism of systematically given corticosteroids. *Dermatologica* 1975;151:129-134.

- <sup>24</sup> Antignac JP, Le Bizec B, Monteau F, Andre F. Study of natural and artificial corticosteroid phase II metabolites in bovine urine using HPLC-MS/MS. *Steroids* 2002;67:873-882.
- <sup>25</sup> Asali LA, Brown KF. Naloxone protein binding in adult and foetal plasma. *Eur J Clin Pharmacol* 1984;27:459-463.
- <sup>26</sup> Madden S, Spaldin V, Park BK. Clinical pharmacokinetics of tacrine. *Clin Pharmacokinet* 1995;28:449-457.
- <sup>27</sup> Lombardo F, Obach RS, Shalaeva MY, Gao F. Prediction of volume of distribution values in humans for neutral and basic drugs using physicochemical measurements and plasma protein binding data. *J Med Chem* 2002;45:2867-2876.
- <sup>28</sup> Ramirez J, Ogan K, Ratain MJ. Determination of vinca alkaloids in human plasma by liquid chromatography/atmospheric pressure chemical ionization mass spectrometry. *Cancer Chemother Pharmacol* 1997;39:286-290.

**Supplemental Table 2. Relationship between the type of liver injury and the extent of metabolism.** Compounds with  $\geq 50\%$  hepatic metabolism were categorized as extensive, whereas compounds with  $< 50\%$  hepatic metabolism or no hepatic metabolism were categorized as not extensive. Compounds with no information on the extent of hepatic metabolism ( $n=11$ ) were excluded from the analysis. Odds ratio estimates are shown with their 95% CIs.

<b>All compounds (n=77)</b>			
	Cholestatic	Non-cholestatic	Odds Ratio
Extensive	36	31	
Not extensive	6	4	0.77 [0.15, 3.62]
<b>BSEP non-inhibitors (n=43)</b>			
	Cholestatic	Non-cholestatic	Odds Ratio
Extensive	17	18	
Not extensive	4	4	0.94 [0.15, 5.96]
<b>BSEP inhibitors (n=34)</b>			
	Cholestatic	Non-cholestatic	Odds Ratio
Extensive	19	13	
Not extensive	2	0	0.00 [0.00, 5.61]

**Supplementary Table 3. Relationship between the type of liver injury and the primary route of excretion.** Compounds were categorized as urinary or biliary if >50% of the reported elimination occurred by a given pathway. Four compounds were excluded because the route of excretion was not available, or the same extent of biliary and urinary excretion was reported. Odds ratios with 95% confidence intervals are shown.

<b>All compounds (n=84)</b>			
	Cholestatic	Non cholestatic	Odds Ratio
Biliary excretion	11	11	
Urinary excretion	35	26	0.74 [0.25, 2.22]
<b>BSEP non-inhibitors (n=49)</b>			
	Cholestatic	Non cholestatic	Odds Ratio
Biliary excretion	1	2	
Urinary excretion	25	21	0.42 [0.01, 8.75]
<b>BSEP inhibitors (n=35)</b>			
	Cholestatic	Non cholestatic	Odds Ratio
Biliary excretion	10	9	
Urinary excretion	10	6	0.67 [0.14, 3.13]

**Supplementary Table 4. Association between maximal unbound concentration C<sub>max,u</sub>, maximal daily dose (MDD), average daily dose (Avg. dose), MRP4 and BSEP inhibition for non-cholestatic and cholestatic drugs.** Valinomycin was excluded from the analysis due to missing information on C<sub>max</sub> and dose. Data are presented as median values.

<b>TYPE OF LIVER INJURY</b>			
<b>All compounds (n=87)</b>			
	C <sub>max,u</sub> ( $\mu$ M)	MDD (mg/day)	Avg. dose (mg/day)
Non-cholestatic (n = 39)	0.27	310	132.5
Cholestatic (n = 48)	2.05	490	300
<b>BSEP INHIBITION</b>			
<b>All compounds (n=87)</b>			
	C <sub>max,u</sub> ( $\mu$ M)	MDD (mg/day)	Avg. dose (mg/day)
Non-cholestatic BSEP non-inhibitors (n=24)	0.30	310	85
Cholestatic BSEP non-inhibitors (n=26)	2.34	620	300
Non-cholestatic BSEP inhibitors (n=15)	0.08	312.5	217
Cholestatic BSEP inhibitors (n=22)	0.05	280	250
<b>MRP4 INHIBITION</b>			
<b>All compounds (n= 87)</b>			
	C <sub>max,u</sub> ( $\mu$ M)	MDD (mg/day)	Avg. dose (mg/day)
Non-cholestatic MRP4 non-inhibitors (n=31)	0.25	240	81
Cholestatic MRP4 non-inhibitors (n=11)	13.2	1600	800
Non-cholestatic MRP4 inhibitors (n=8)	0.10	550	200
Cholestatic MRP4 inhibitors (n=37)	0.11	360	215
<b>BSEP non-inhibitors (n=50)</b>			
	C <sub>max,u</sub> ( $\mu$ M)	MDD (mg/day)	Avg. dose (mg/day)
Non-cholestatic MRP4 non-inhibitors (n=20)	0.27	160	81
Cholestatic MRP4 non-inhibitors (n=10)	13.3	1600	580
Non-cholestatic MRP4 inhibitors (n=4)	0.27	950	143
Cholestatic MRP4 inhibitors (n=16)	0.28	400	280
<b>BSEP inhibitors (n=37)</b>			
	C <sub>max,u</sub> ( $\mu$ M)	MDD (mg/day)	Avg. dose (mg/day)
Non-cholestatic MRP4 non-inhibitors (n=11)	0.05	200	288
Cholestatic MRP4 non-inhibitors (n=1)	0.01	2400	2100
Non-cholestatic MRP4 inhibitors (n=4)	0.16	257.5	123
Cholestatic MRP4 inhibitors (n=21)	0.06	200	125

If dosing was given as mg/kg or mg/m<sup>2</sup>, the maximum and average dose was calculated based on an adult with 70 kg and a body surface area of 1.7.