

Table S2. Immunogenic salivary antigens of four different *T. infestans* strains.

| kDa* | IgG antigens | | | | | | | | IgM antigens | | | | | | | |
|------|--------------|---|---------|---|-------|---|------|---|--------------|---|---------|---|-------|---|------|---|
| | Argentina | | Bolivia | | Chile | | Peru | | Argentina | | Bolivia | | Chile | | Peru | |
| | N | A | N | A | N | A | N | A | N | A | N | A | N | A | N | A |
| 91 | x | | | | | | | | x | x | | | | | | |
| 88 | | x | | | | | x | x | x | x | | | | x | | |
| 79 | x | | x | x | x | | | x | | x | x | | | | | |
| 71 | x | x | | | | | x | | | | x | x | | | | |
| 64 | x | | | | x | | | | | | | | x | x | | x |
| 60 | | | | x | | | | | | x | x | | x | x | | |
| 55 | x | | | | | | | | | | | | x | x | | |
| 51 | | | | x | | | | | | | | | | | | |
| 48 | | | | x | | | | | | | | | | | | |
| 43 | | | x | x | | | | x | x | | x | | | | | |
| 40 | x | | | | x | | x | x | x | | | | | x | x | |
| 36 | | | x | x | | x | x | x | x | x | x | x | x | x | x | x |
| 35 | x | x | x | x | x | x | x | x | x | x | x | x | x | x | x | x |
| 33 | x | x | x | x | x | x | x | x | x | x | x | x | x | x | x | x |
| 32 | | | x | x | x | x | x | x | x | x | | | x | x | x | x |
| 30 | x | | x | x | | x | x | x | x | x | | | | | | |
| 28 | x | | | x | | x | | | | | | | | x | | |
| 27 | | | | x | | | | | | | | | | | | |
| 25 | x | x | x | x | x | x | | | x | | | x | x | x | x | x |
| 24 | | | x | x | | | | | | | | | | | | |
| 21 | x | | | x | x | x | | x | | | | x | | x | | |
| 18 | | | | x | | | | | x | | | x | | | | |
| 15 | | | x | x | x | x | | | x | x | | | x | | | |
| 14 | x | | x | x | x | | | | | | | | | x | | |
| 13 | | | | x | x | | | | | | | | | | | |
| 12 | | | | | x | | | | | | | | | | | |
| 8 | | | | | x | | | | | | | | | | | |

* The presented proteins reacted with IgG or IgM antibodies of at least one guinea pig serum, respectively.