

Supporting Table 1. Demographics, BMI, Blood Pressure, and Conscious Level at Study

Entry

Race/Ethnicity - Gender	Number in Group (%)	Age Years ± SD	BMI kg/m² Median and IQR (n)	Systolic BP mm Hg Mean ± SD (n)	Diastolic BP mm Hg Mean ± SD (n)	Coma Grade Mean ± SD (n)
All Subjects	133 (100)	43.8 ± 14.1	28.7 24.6-32.8 (101)	126 ± 20 (133)	68 ± 14 (133)	2.2 ± 1.1 (132)
- Women	94(70.7)	44.7 ± 13.5	29.0 24.2-33.9 (67)	127 ± 21 (94)	68 ± 14 (94)	2.3 ± 1.1 (94)
- Men	39 (29.3)	41.7 ± 15.5	28.4 25.6-31.1 (34)	125 ± 20 (39)	67 ± 14 (39)	1.9 ± 1.9 (38)
White	76 (57.1)	45.4 ± 14.4	28.7 24.3-32.9 (57)	124 ± 21 (76)	66 ± 14 (76)	2.1 ± 1.0 (75)
- Women	55 (72.3)	46.1 ± 13.5	29.0 24.0-33.3 (39)	124 ± 21 (55)	66 ± 14 (55)	2.3 ± 1.1 (55)
- Men	21 (27.6)	43.6 ± 16.7	27.9 26.1-31.1 (18)	124 ± 22 (21)	65 ± 13 (21)	1.8 ± 0.8 (20)
African-American	21 (15.8)	41.6 ± 13.3 (21)	32.5 26.2 -42.2 (15)	135 ± 19 (21)	70 ± 16 (21)	2.3 ± 1.3 (21)
- Women	15 (71.4)	42.3 ± 14.1	34.3 26.0-47.0 (10)	137 ± 20 (15)	72 ± 13 (15)	2.5 ± 1.3 (15)
- Men	6 (28.6)	(15) 39.8 ± 12.2 (6)	29.0 28.1-32.5 (5)	130 ± 17 (6)	64 ± 23 (6)	1.8 ± 1.2 (6)
Hispanic	20 (15.0)	39.1 ± 12.1 (20)	27.1 21.9-32.4 (13)	127 ± 18 (20)	71 ± 12 (20)	2.0 ± 1.0 (20)
- Women	16 (80)	41.6 ± 12.1	28.6 22.5-32.0 (10)	129 ± 18 (16)	71 ± 13 (16)	2.4 ± 1.2 (16)
- Men	4 (20)	(16) 29.3 ± 6.2 (4)	25.3 22.4-28.9 (3)	119 ± 12 (4)	70 ± 7 (4)	2.0 ± 1.2 (4)
Asian	9 (6.8)	41.2 ± 16.6 (9)	24.8 23.1 – 26.9 (7)	119 ± 24 (9)	65 ± 15 (9)	2.4 ± 1.0 (9)
- Women	5 (56)	42.6 ± 18.3 (5)	24.7 23.9 - 25.3 (4)	113 ± 26 (5)	63 ± 15 (5)	2.4 ± 1.1 (5)
- Men	4 (44)	39.5 ± 16.7 (4)	23.9 23.9 - 28.3 (3)	126 ± 23 (4)	68 ± 17 (4)	2.5 ± 1.0 (4)
Native American	3 (2.3)	43.7 ± 18 (3)	29.8 25.2-41.1* (3)	132 ± 22 (3)	79 ± 3 (3)	2.3 ± 0.7 (6)
- Women	0 (0)	-	-	-	-	-
- Men	3 (100)	43.7 ± 18 (3)	29.8 25.2-41.1 (3)	132 ± 22 (3)	79 ± 3 (3)	2.3 ± 0.7 (6)
Hawaiian/Pacific- Islander	1 (0.8)	64	30.0	140 (1)	70 (1)	2 (1)
- Women	0 (0)	-	-	-	-	-
- Men	1 (100)	64	30.0	140 (1)	70 (1)	2 (1)

Other	3 (2.3)	51.7 ± 12.7 (3)	29.0 (3) 28.3 – 39.0* (3)	131 ± 4 (3)	68 ± 6 (3)	2.0 ± 1.4 (3)
- Women	3 (100)	51.7 ± 12.7 (3)	29.0 (3) 28.3 - 39.0* (3)	131 ± 4 (3)	68 ± 6 (3)	2.0 ± 1.4 (3)
- Men	0 (0)	-	-	-	-	-

Notes: (n) = Number of subjects per group (which may vary where data are missing)

BMI = Body Mass Index

BP = Blood pressure

SD = Standard deviation

IQR = Interquartile range

*range for 3 subjects