

SUPPORTING INFORMATION FOR MANUSCRIPT TITLED
“Chemoenzymatic Synthesis of Oligohyaluronan-Lipid Conjugates”

Dipali Ruhela, Saul Kivimäe and Francis C. Szoka, Jr.*

Departments of Bioengineering, Therapeutic Sciences and Pharmaceutical Chemistry, University of California San Francisco, 513 Parnassus Ave, San Francisco, CA 94143-0912

Table of contents

FIGURE S1: MALDI-TOF (negative mode) profile of HA-DiPhPE digested over different time intervals with hyaluronidase

FIGURE S2: MALDI-TOF (negative mode) analysis of aqueous layer obtained after Phospholipase digestion of HA_n-DiPhPE

Figure S1. MALDI-TOF (negative mode) profile of HA-DiPhPE digested over different time intervals with hyaluronidase






Figure S2. MALDI-TOF (negative mode) analysis of aqueous layer obtained after Phospholipase digestion of HA_n-DiPhPE



Parent HAn-DiPhPE (1)	Observed MW of HA (2)	Expected MW of HA (2)
HA ₁₀ -DiPhPE	1938	1939 (M-H ₂ O)
HA ₁₀ -DiPhPE	1957	1957 (M)
HA ₁₀ -(DiPhPE) ₂	1981	1982 (M-H ₂ O)
HA ₁₂ -DiPhPE	2321	2318 (M-H ₂ O)
HA ₁₂ -DiPhPE	2338	2336 (M)
HA ₁₂ -(DiPhPE) ₂	2360	2361 (M)
HA ₁₄ -DiPhPE	2703	2698 (M-H ₂ O)
HA ₁₄ -(DiPhPE) ₂	2741	2741 (M-H ₂ O)
HA ₁₆ -DiPhPE	3092	3095 (M)



Representative MALDI profile