

**Table S2** Correlation of n-terminal pro-brain natriuretic peptide (NT-proBNP) with hemodynamic parameters in PH patients with and without renal insufficiency.

<b>Parameters</b>	<b>NT-proBNP [ng/L]</b>	<b>Normalized NT- proBNP ratio</b>
<b>Patients with GFR &gt;60 [ml/min/1.73 m<sup>2</sup>] (n=56)</b>		
Mean PAP [mm Hg]	0.454; <0.001	0.404; 0.002
RAP [mm Hg]	0.244; 0.2	0.245; 0.2
PVR [dyne*s*cm <sup>5</sup> ]	0.513; <0.001	0.474; <0.001
CI [L/min*m <sup>2</sup> ]	-0.370; 0.006	-0.386; 0.004
<b>Patients with GFR ≤60 [ml/min/1.73 m<sup>2</sup>] (n=40)</b>		
Mean PAP [mm Hg]	-0.057; 0.74	0.006; 0.97
RAP [mm Hg]	0.410; 0.016	0.385; 0.018
PVR [dyne*s*cm <sup>5</sup> ]	0.343; 0.044	0.318; 0.052
CI [L/min*m <sup>2</sup> ]	-0.457; 0.009	-0.452; 0.007

Data are presented as R-value; p-value of Spearman's rank correlation.