


Table S1. Incidence of DAIDS Adverse Events

	LPV/r	NNRTI	p-value
Altered mental status	0.00 (0)	0.55 (1)	n/a
Anemia	2.55 (5)	4.99 (9)	0.39
Diarrhea	0.51 (1)	1.11 (2)	n/a
Respiratory Distress	2.04 (4)	2.22 (4)	0.95
Elevated ALT	1.53 (3)	3.32 (6)	0.26
Elevated AST	0.51 (1)	0.55 (1)	0.95
Leukopenia	0.51 (1)	1.11 (2)	0.51
Low serum glucose	0.00 (0)	0.55 (1)	n/a
Neutropenia	24.00 (47)	28.26 (51)	0.42
Severe malnutrition	0.00 (0)	0.55 (1)	n/a
Stevens Johnson Syndrome	0.00 (0)	0.55 (1)	n/a
Thrombocytopenia	8.17 (16)	3.87 (7)	0.14
Unintentional weight Loss	0.51 (1)	0.55 (1)	n/a

Incidence per 100-person-years (# of events) of Division of HIV AIDS Grade III or IV adverse events¹⁰; P-value for comparison between arms using negative binomial regression; n/a comparison not possible.

Figure S1



Number at risk		0	24	48	72	96
arm = NNRTI		27	26	24	22	21
arm = LPV/r		27	27	26	25	24

Figure S2a




Figure S2b

