

Matsumoto et al., Table S2

Primary antibodies	Application(s) and working dilution(s)
Anti- β -catenin (Cell Signaling)	IB 1:5,000; IF 1:500
Anti-phospho-LRP6 (Ser1490) (Cell Signaling)	IB 1:1,000
Anti-active (cleaved) caspase-3 (Cell Signaling)	IF 1: 250
Anti-Axin1 (Cell signaling)	IB 1:1,000
Anti-Dvl2 (Cell Signaling)	IB 1:1,000
Anti-phospho-myosin light chain 2 (Thr18/Ser19) (Cell Signaling)	IF 1:200
Anti-RhoA (Cell Signaling)	IB 1:1,000
Anti-Arf6 (Cell Signaling)	IB 1:1,000
Anti-YAP/TAZ (Cell Signaling)	IB 1:2,000; IF 1:200
Anti-phospho-Ezrin (Thr567) (Abcam)	IF 1:200
Anti-Ezrin (Abcam)	IF 1:200
Anti-LRP6 (Abcam)	IB 1:1,000
Anti-Arl4c (Abcam)	IF 1:100; IHC 1:100
Anti-Cytokeratin8 (Abcam)	IF 1:200
Anti- β -tubulin (Sigma)	IB 1:5,000
Anti-Cytokeratin,pan (Sigma)	IF 1:100
Anti-NCAM (Sigma)	IF 1:100
Anti-Rac1 (BD Biosciences)	IB 1:5,000
Anti-FAK (BD Biosciences)	IB 1:2,500
Anti- β -catenin (BD Biosciences)	IB 1:2,500
Anti-E-cadherin (BD Biosciences)	IF 1:250
Anti-HSP90 (BD Biosciences)	IB 1:5000
Anti-Clathrin (BD Biosciences)	IB 1:5000
Anti-phospho-FAK (Tyr397) (Invitrogen)	IB 1:1,000
Anti-Dvl3 (Santa Cruz)	IB 1:1,000
Anti-PKC ζ (Santa Cruz)	IF 1:200
Anti-Ets1 (C-20) (Santa Cruz)	IB 1:1,000
Anti-Ets1 (C-4) (Santa Cruz)	IB 1:1,000
Anti-mAG1 (MBL)	IF 1:500
Anti-mKO2 (MBL)	IF 1:500
Anti-Arl4c (Proteintech)	IB 1:100
Anti-Tcf3/4 (Millipore)	IB 1:1,000
Anti-Dvl1 (DIX) (generated in rabbits by immunization)	IB 1:5,000

Supplementary Table S2 Primary antibodies used in this study. IB, immunoblotting; IF, immunofluorescence; IHC, immunohistochemistry.