


Supplementary information

Combined staurosporine and retinoic acid induces differentiation in retinoic acid resistant acute promyelocytic leukemia cell lines


Dong-zheng Ge^{1,2}, Yan Sheng¹, Xun Cai¹

¹Shanghai Institute of Hematology and State Key Laboratory of Medical Genomics, Rui-jin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China; ²Department of Hematology, Peace Hospital of Changzhi Medical Colledge, Changzhi 046000, China.




Supplemental Figure S1.

Full size blots of the immunoblot detection shown in Figure 2. The left panel were full size blots of NB4-R1 cells, the right panel were those of NB4-R2 cells.* indicates non-specific bands. β -actin was probed in the same membrane of PML-RAR α .




Supplemental Figure S1 (continued) .

Full size blots of the immunoblot detection shown in Figure 2. The left panel were full size blots of NB4-R1 cells, the right panel were those of NB4-R2 cells.



Supplemental Figure S2.

Full size blots of the immunoblot detection shown in Figure 3. The left panel were full size blots of NB4-R1 cells, the right panel were those of NB4-R2 cells.



Supplemental Figure S3.

Full size blots of the immunoblot detection shown in Figure 4. The left panel were full size blots of NB4-R1 cells, the rightpanel were those of NB4-R2 cells.* indicates non-specific bands.