

## Local Gyrfication Index in Probands with Psychotic Disorders and Their First-Degree Relatives

### *Supplemental Information*

**Table S1.** Handedness information of included participants

	Controls			Psychotic Bipolar Disorder			Schizoaffective Disorder			Schizophrenia			Relatives		
	L	R	A	L	R	A	L	R	A	L	R	A	L	R	A
Handedness	30	207	2	18	121	1	6	80	4	18	134	3	19	277	3
Distribution <sup>a</sup>	13%	87%	1%	13%	86%	1%	7%	89%	4%	12%	86%	2%	6%	93%	1%

L, left-handed; R, right-handed; A, ambidextrous.

<sup>a</sup> 8 participants were unknown.

**Table S2.** Distribution of relatives by their own diagnosis and their related proband's diagnosis

	Proband Diagnosis	Relative's Own Diagnosis				Total
		Axis II Cluster A	Axis I Non-Psychotic	Axis II Cluster B/C	No Diagnosis	
	Schizophrenia	17	53	0	44	114 38%
	Schizoaffective Disorder	13	38	2	32	85 28%
	Psychotic Bipolar Disorder	5	61	2	33	101 34%
	Total	35 12%	152 51%	4 1%	109 36%	300

**Table S3.** Sites of included participants

	<b>Controls</b>	<b>Schizophrenia</b>	<b>Schizoaffective Disorder</b>	<b>Psychotic Bipolar Disorder</b>	<b>Relatives</b>	<b>Total</b>
Wayne State University	33	28	3	25	40	129
Harvard University	26	7	1	7	23	64
Maryland Psychiatric Research Center	44	52	21	23	50	190
University of Chicago, University of Illinois at Chicago	57	30	21	48	97	253
University of Texas Southwestern	49	14	24	23	55	165
Institute of Living, Yale University	34	26	20	15	35	130