

SUPPLEMENTAL TABLE 1
Cytokine responses in acute and convalescent samples of typhoid fever patients

	Mean acute	SEM	Mean convalescent	SEM	Relative change	P value
EGF	17.25	11.32	23.11	23.11	1.34	0.4067
Eotaxin	0.71	0.71	5.52	1.86	7.77	0.0154
FGF-2	0.00	0.00	4.15	3.36	4.15	0.2716
Flt-3 Ligand	4.81	4.40	6.32	3.50	1.32	0.2907
Fractalkin	4.79	4.79	59.70	30.44	12.45	0.0101
G-CSF	1226.19	578.58	2738.51	2074.23	2.23	0.4311
GM-CSF	494.68	180.66	819.55	182.97	1.66	0.2235
GRO	0.00	0.00	OOOR >			
IFN α 2	3.49	3.49	2.33	1.36	0.67	0.4964
IFN γ	572.82	392.51	5482.00	2941.18	9.57	0.0492
IL1A	265.99	124.33	395.55	84.18	1.49	0.1957
IL1B	263.05	73.69	788.21	164.85	3.00	0.0851
IL1Ra	0.00	0.00	21.67	21.00	21.67	0.1913
IL-2	3.26	1.71	17.28	8.65	5.30	0.0935
IL-3	ND		ND			
IL-4	6.28	6.28	9.72	3.97	1.55	0.2513
IL-5	2.79	2.10	58.32	41.57	20.88	0.0562
IL-6	1884.40	1212.60	8079.68	3013.30	4.29	0.1613
IL-7	16.36	8.40	56.25	17.43	3.44	0.1571
IL-8	OOOR >		OOOR >			
IL-9	1.06	1.06	31.34	10.22	29.56	0.0065
IL-10	53.53	38.80	150.40	29.30	2.81	0.1206
IL12 (p40)	47.04	35.88	89.02	16.76	1.89	0.1277
IL12 (p70)	3.52	1.44	8.50	1.67	2.41	0.1215
IL-13	17.54	8.37	198.76	96.51	11.33	0.0104
IL-15	0.42	0.42	3.95	0.75	9.48	0.0130
IL-17	4.81	4.81	5.71	3.12	1.19	0.2051
IP 10	112.81	53.99	516.92	155.47	4.58	0.1057
MCP-1	0.00	0.00	0.00	0.00		
MCP-3	0.00	0.00	279.68	279.68	279.68	0.2113
MDC	0.00	0.00	0.00	0.00		
MIP-1 α	OOOR >		OOOR >			
MIP-1 β	79.66	53.68	2202.62	556.99	27.65	0.0417
sCD40L	6.26	6.26	99.96	31.87	15.96	0.0354
sIL-2Ra	5.03	3.28	67.88	25.71	13.49	0.0561
TGF α	0.00	0.00	3.80	2.20	3.80	0.1005
TNF α	910.43	500.57	978.84	210.93	1.08	0.2625
TNF β	0.82	0.82	27.90	14.31	34.16	0.0366
VEGF	9.10	9.10	48.53	19.94	5.33	0.0718

ND = not detected; OOR > = out of range high.