Supplemental information to accompany

"Aromatic organosulfates in atmospheric aerosols: synthesis, characterization, and abundance"

Sean Staudt¹, Shuvashish Kundu¹, Hans-Joachim Lehmler², Xianran He², Tianqu Cui,³ Ying-Hsuan Lin,³ Kasper Kristensen,⁴ Marianne Glasius,⁴ Xiaolu Zhang,^{5,#} Rodney J. Weber,⁵ Jason D. Surratt³, Elizabeth A. Stone^{1,*}

¹Department of Chemistry, University of Iowa, Iowa City, IA 52242, United States

²Department of Occupational and Environmental Health, College of Public Health, University of Iowa, Iowa City, IA 52242, USA

³Department of Environmental Sciences and Engineering, Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA

⁴Department of Chemistry and iNANO, Aarhus University, 8000 Aarhus C, Denmark

⁵School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332, USA

[#]Present address: Department of Civil and Environmental Engineering, University of California, Davis, CA 95616, USA

*Corresponding author phone: +1-319-384-1863, fax: +1-319-335-1270;

email: <u>betsy-stone@uiowa.edu</u>

Figure S1. Experimental time profiles for the toluene SOA experiments conducted in the presence of either neutral (blue) or acidified (red) sulfate seed aerosol. This figure demonstrates that the concentrations of gases and aerosol are similar between both experiments conducted on the same day within the UNC dual outdoor smog chamber facility.

