

**Table S2**  
**Vectors used for plasmid generation**

Vector	purpose	Source
pGEX-4T	N-terminal GST fusions for expression in / purification from <i>E. coli</i>	GE Healthcare Life Sciences, Pittsburg, PA, USA
pMAL-c2	N-terminal MBP fusions for expression in / purification from <i>E. coli</i>	New England Biolabs, Ipswich, MA, USA
pEG202	Yeast-two hybrid bait vector	Golemis et al. 2008
pJG4-5	Yeast-two hybrid prey vector	Golemis et al. 2008
pJLSmart	GW entry vector for expression in eukaryotes; N- and C-terminal fusions	Mathieu et al. 2007
pJM51	modified pJLSmart GW entry vector: contains Shine Dalgarno sequence for expression in prokaryotes	Cheng et al. 2011
pGWB417	GW destination vector for transient protein expression in <i>N. benthamiana</i> . Adds C-terminal 4x Myc tag	Nakagawa et al. 2007
pCPP5372	GW destination vector for effector expression in <i>Pst.</i> Contains hrp promoter and C-terminal HA tag.	Oh et al. 2007
pBTEX:PRF	Co-expression of tomato PRF with Pto in <i>N. benthamiana</i> .	Du et al. 2012
pMAL-Pto	Expression of Pto in <i>E. coli</i>	Loh et al. 1995
pMAL-Fen	Expression of Fen in <i>E. coli</i>	Loh et al. 1995

**Table S2**