

Comparative analyses of lipidomes and transcriptomes reveal a concerted action of multiple defensive systems against photooxidative stress in *Haematococcus pluvialis*

Yunho Gwak¹⁺, Yong-sic Hwang²⁺, Baobei Wang³, Minju Kim¹, Jooyeon Jeong¹, Choul-Gyun Lee⁴, Qiang Hu⁵, Danxiang Han^{3*} and EonSeon Jin^{1*}

¹Department of Life Science, Research Institute for Natural Science, Hanyang University, Seoul, 133-791, South Korea

²Department of Bioscience and Biotechnology, Konkuk University, Seoul, 143-701, South Korea

³College of Technology and Innovation, Arizona State University, Mesa, Arizona 85212, USA

⁴Department of Biotechnology, Institute of Industrial Biotechnology, Inha University, Incheon, 402-751, South Korea

⁵Center for Microalgal Biotechnology and Biofuels, Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, Hubei 430072, China

+Both authors contributed equally to this work

*** Corresponding authors:**

EonSeon Jin: Tel: +82-2-2220-2561; fax: 82-2-2299-2561; e-mail: esjin@hanyang.ac.kr

Danxiang Han: Tel: +1-480-727-5661; email: Danxiang.Han@asu.edu

Fig. 5. Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis of *H. pluvialis* transcripts.




Fig. S6. KEGG pathway analysis of *H. pluvialis*. Macrozooids (A)




Fig. S6. KEGG pathway analysis of *H. pluvialis*. Cysts (B)

