


## Supplemental Materials

Supplemental Figure 1



Supplemental Figure 1. Amygdala ROI. Anatomical ROI of the amygdala were determined in AFNI based on the Talairach and Tournoux (1988) coordinate system.

Supplemental Table 1. Whole brain analyses.

	Hemisphere	Vol (mm <sup>3</sup> )	Talairach coordinates		
			x	y	z
<i>Arousal</i>					
Lingual Gyrus	Bilateral	<b>36451</b>	<b>0.9</b>	<b>-76.6</b>	<b>-5.1</b>
Thalamus	Right	<b>1418</b>	<b>25.7</b>	<b>-29.7</b>	<b>3.4</b>
Dorsolateral PFC	Right	<b>1231</b>	<b>49.2</b>	<b>-14.5</b>	<b>29.9</b>
<i>Valence</i>					
Middle Temporal Gyrus	Left	<b>2631</b>	<b>-50.0</b>	<b>-57.0</b>	<b>-2.2</b>

*Note:* Location, volumes, and coordinates from Talairach and Tournoux (1988) for the center of mass for areas of activation. Significance threshold criteria:  $p < 0.005$  and cluster size of 788 mm<sup>3</sup> (14 voxels of  $3.75 \times 3.75 \times 4.00$  mm dimension) resulted in  $p < 0.05$ . Significance criteria:  $t[20] > 3.16$ ,  $p < 0.05$  (corrected),  $d = 0.69$ .