

Supplementary Table 1: clinicopathological characteristics, PDAC cohort.

Characteristic	All patients (n=26)	Weak asTF (n=5)	Strong asTF (n=4)
Median age, years (range)	65 (41-81)	62.5 (56-69)	57 (56-78)
Sex: M/F	11/15	3/2	3/1
pT category: T1/2/3	4/0/22	1/0/4	0/0/4
pN category: N0/1/2	7/18/1	2/3/0	1/3/0
Histological grade: G1/2/3	1/18/7	0/3/2	0/4/0
Pathological margin: involved/uninvolved	7/19	2/3	1/3
Perilymphatic, perivascular invasion: yes/no	4/21	1/4	1/3
Perineural invasion: yes/no	17/9	3/2	4/0
Recurrence: yes/no	11/15	3/2	1/3
Median time to recurrence, months (range)	16 (10-22)	21 (16-22)	22

Figure legend – Supplementary Figure 1

(a) Quantification of vessel density in the PDAC specimens stratified into weak asTF and strong asTF cohorts. (b) Levels of $\beta 1/\beta 3$ integrins and EGFR on the surface of Pt45P1 and Pt45P1/asTF+ cells (n=3). (c) Total number of microparticles generated by Pt45P1 and Pt45P1/asTF+ cells (n \geq 3). (d) Levels of flTF protein on cell surfaces assessed using on-cell western assay (n=3). (e) Levels of $\beta 1/\beta 3$ integrins on the surface of MP generated by Pt45P1 and Pt45P1/asTF+ cells (n=3).

Supplementary Figure 1

